

THE ROLE OF LAMININ IN DEVELOPMENT, REGENERATION AND INJURIES OF THE NERVOUS SYSTEM

SANNA MURTOMÄKI-REPO

**Institute of Biomedicine
The Brain Laboratory
Department of Anatomy
University of Helsinki**

Academic dissertation

*To be presented, with the permission of the Faculty of Medicine of the University of Helsinki
for public examination in the auditorium of the Institute of Biomedicine, Department of
Anatomy, Siltavuorenpenger 20A, on the 26th of May 2000, at 12 o'clock.*

Helsinki 2000

Supervisor

Docent Päivi Liesi
The Brain Laboratory
Institute of Biomedicine
Department of Anatomy
University of Helsinki

Reviewers

Professor Eero Castren
A. I. V. Institute
University of Kuopio

Docent Ilmo Leivo
Department of Pathology
University of Helsinki

Opponent

Professor Dan Lindholm
Department of Developmental Neuroscience
University of Uppsala
Uppsala, Sweden

ISBN 952-91-2141-5 (nid)
ISBN 952-91-2141-3 (PDF version)
Yliopistopaino
Helsinki 2000

CONTENTS

PREFACE	5
ABBREVIATIONS	6
LIST OF THE ORIGINAL PUBLICATIONS	7
INTRODUCTION	8
REVIEW OF THE LITERATURE	10
1. Laminin	10
1.1. General features of laminins	10
1.2. Nomenclature of laminins	11
1.3. Structure of laminin-1	12
1.4. Biological functions of laminin	16
1.5. Analysis of the functions of laminin-1 using proteolytic fragments and synthetic peptides	19
1.6. Molecular interactions of laminin-1	23
1.7. Laminins in the nervous system	25
2. Nervous system development, degeneration and regeneration	29
2.1. Neuronal differentiation	29
2.1.1. Laminin in neuronal differentiation	31
2.1.2. Neuronal cell lines	32
2.1.3. Teratocarcinoma cells as a model system for neuronal differentiation	33
2.2. Neuronal migration	33
2.2.1. Current models of neuronal migration	33
2.2.2. Molecules involved in neuronal migration	35
2.1.2.a ECM molecules	35
2.1.2.b Cell adhesion molecules	37
2.1.2.c Other mediators of neuronal migration	37
2.2.3. Neuronal migration during cerebellar development	38
2.2.4. The weaver mutant mouse as a model of neuronal migration and death	41
2.3. Neurite outgrowth and axonal targeting	44
2.3.1. Laminin and other extracellular matrix molecules in neurite outgrowth	44
2.3.2. Netrins, semaphorins and ephrins in axon guidance	47
2.3.3. Cell Adhesion molecules	47
2.4. Neuronal degeneration	48
2.4.1. Alzheimer's disease	49
2.4.1.1. Pathophysiology of Alzheimer's disease	49
2.4.1.2. Extracellular matrix molecules in Alzheimer's disease	51
2.5. Nervous system regeneration	52
2.5.1. Regeneration in the peripheral and central nervous systems	52
2.5.2. Attempts to regenerate PNS injuries	54
2.5.2.1. Laminin treatment of the PNS injuries	55
2.5.3. Attempts to regenerate CNS injuries	56

AIM OF THE STUDY	58
MATERIALS AND METHODS	59
RESULTS	69
I Laminin and its neurite outgrowth-promoting domain in the brain in Alzheimer's disease and Down's syndrome patients	69
1.1. Immunochemical characterization of antibodies against γ 1-chain neurite outgrowth promoting peptides of laminin-1	69
1.2. Immunocytochemical studies on of laminin and its neurite outgrowth domain in the Alzheimer's disease brain	69
1.3. Demonstration of laminin-1 mRNA in the Alzheimer's disease brain tissue	70
1.4. Immunoblotting experiments on the Alzheimer's disease brain tissue	71
II Increased proteolytic activity of the granule neurons may contribute to neuronal death in the weaver mouse cerebellum	71
2.1. <i>In vitro</i> studies of weaver cerebellum	71
2.2. <i>In vivo</i> studies of weaver cerebellum	72
III Use of paper for treatment of a peripheral nerve trauma in the rat	73
3.1. Immunocytochemical analysis of nerve regeneration	73
3.2. The twitch tensions of the muscles after treatment	73
3.3. The scores and rate of autotomy	74
IV Neurofilament proteins are constitutively expressed in F9 teratocarcinoma cells	74
4.1. Immunocytochemical analysis of induced F9 cells	74
4.2. Northern blot analysis of the induced F9 cells	75
4.3. - RT-PCR analysis	75
4.4. Cloning of a neuronal F9 cell line	75
DISCUSSION	76
I Laminin and its neurite outgrowth-promoting domain in the brain in Alzheimer's disease and Down's syndrome patients	76
II Increased proteolytic activity of the granule neurons may contribute to neuronal death in the weaver mouse cerebellum	79
III Use of cellulose for treatment of a peripheral nerve trauma in the rat	81
IV Neurofilament proteins are constitutively expressed in F9 teratocarcinoma cells	82
SUMMARY AND CONCLUSIONS	85
REFERENCES	87

PREFACE

The experiments of the present study were carried out at the Institute of Biotechnology, University of Helsinki, and the Institute of Biomedicine, Department of Anatomy, University of Helsinki during the years 1992-2000.

I would like to thank Professor Ismo Virtanen for his encouragement and for putting the facilities of the institute at my disposal.

My warmest thanks to my supervisor Docent Päivi Liesi for providing guidance, support, and encouragement during my work and guiding me to the fascinating world of science.

I would like to thank Professor Eero Castren and Docent Ilmo Leivo, my appointed reviewers, for their constructive criticism of the manuscript.

My sincere thanks to Docent Timo Kauppila for stimulating discussions, to Docent Leila Risteli, Docent Juha Risteli, Dr Ulla-Maija Koivisto, Docent Staffan Johansson, Dr Ekkhart Trenkner, Dr Jerry Wright, Docent Olli Saksela, Docent Erkki Jyväsjärvi, Dr Heikki Mansikka, and Professor Antti Pertovaara for their stimulating collaboration during these years.

I wish to thank Ms Raija Sassi and Mr Reijo Karppinen for their expert help with the artwork, Mr Riku Murtomäki for resolving my numerous computer problems, Dr Timo Laine for discussions and, Mrs Outi Rauanheimo for all the help that made my life easier at the department. I warmly thank the personnel of the Institute of Biomedicine for creating such a pleasant working environment.

I like to thank my parents Mirja and Antti and my parents in law Juhani and Martta for their support and for invaluable time they spent with my children whenever necessary for the work.

Last but not the least, I would like to dedicate this work to my husband Timo and my sons Rasmus and Oskari, and thank them for their patience and love.

This work was partially financed by a grant from the University of Helsinki.

ABBREVIATIONS

A β	amyloid β -peptide
ACT	α -antichymotrypsin
AD	Alzheimer's disease
ALS	amyotrophic lateral sclerosis
AMOG	adhesion molecule on glia
APP	β -amyloid precursor protein
BL	basal laminae
BMP	bone morphogenetic protein
CAM	cell adhesion molecule
CMD	congenital muscular dystrophy
CNS	central nervous system
CO	cytochrome c oxidase
CS	chondroitin sulfate
dbcAMP	dibutyl cyclic adenosine mono phosphate
DCC	deleted in colorectal cancer
GAG	glycosaminoglycan
GFAP	glial acidic fibrillary protein
GIRK	G-protein coupled inwardly rectifying K ⁺ channel gene
DS	dermatan sulfate
ECM	extracellular matrix
EGF	epidermal growth factor
EHS	Engelbreth-Holm-Swarm
FAD	familial Alzheimer's disease
FGF	fibroblast growth factor
HC	hippocampus
HS	heparan sulfate
HSPG	heparan sulfate proteoglycans
KS	keratan sulfate
LSB-	Laemmli sample buffer without β -mercaptoethanol
Ng-CAM	neuron-glia cell adhesion molecule
N-CAM	neural cell adhesion molecules
NF	neurofilament
NMDA	N-methyl-D-aspartate
PGs	proteoglycans
PNS	peripheral nervous system
PS-1	presenilin-1
PS-2	presenilin-2
RA	retinoic acid
RGC	retinal ganglion cell
RMP	resting membrane potential
RPE	retinal pigment epithelial
SAP	serum amyloid P
Shh	Sonic hedgehog
SN	substantia nigra
TGF beta 1	transforming growth factor beta 1
tPA	tissue plasminogen activator
TUJI	neuron specific β -tubulin III isoform

LIST OF ORIGINAL PUBLICATIONS

This thesis is based on the following original papers, referred to in the text by Roman numerals I-IV

- I S. Murtomäki, J. Risteli, L. Risteli, U.-M. Koivisto, S. Johansson and P. Liesi.: Laminin and its neurite outgrowth-promoting domain in the brain in Alzheimer's disease and Down's syndrome patients. *J. Neurosci. Res.* 32:261-273, 1992.
- II. S. Murtomäki, E. Trenkner, J.M. Wright, O. Saksela and P. Liesi: Increased proteolytic activity of the granule neurons may contribute to neuronal death in the weaver mouse cerebellum. *Dev. Biol.* 168, 635-648, 1995.
- III. T. Kauppila, E. Jyväsjärvi, S. Murtomäki, H. Mansikka, A. Pertovaara, I. Virtanen and P. Liesi: Use of paper for treatment of a peripheral nerve trauma in the rat. *Neuroreport.*, 8, 3151-3155, 1997.
- IV S. Murtomäki, I. Virtanen and P. Liesi: Neurofilament proteins are constitutively expressed in F9 teratocarcinoma cells. *Int. J. Dev. Neurosci.*, 17, 829-838, 1999.

INTRODUCTION

Mechanisms of neuronal migration, neurite outgrowth, neuronal degeneration and neuronal regeneration have been the main focus of neurobiological research for the past hundred years. Even though several major principles have emerged, molecular mechanisms of brain development and neuronal injuries are still largely unknown. In recent years, investigators have concentrated on identification of molecules involved in cell-to-cell interactions in the nervous system. Laminin-1, originally isolated from a mouse tumor (so called EHS sarcoma), rich in basement membrane proteins, has been shown to be one of the key molecules in nervous system development and response to trauma. Laminin-1 promotes neurite outgrowth, guides neuronal migration, promotes neuronal regeneration and is involved in neuronal differentiation as well as death mechanisms.

As soon as these nervous system related general functions of laminin-1 were identified, attempts began to 1) determine specific domains of laminin-1 that mediate various functions of this protein and 2) understand the molecular interactions that mediate the different functions of laminin-1. Thusfar, five neurite outgrowth promoting domains have been identified in the laminin-1 molecule: One in domain I of the γ 1-chain, one in the α 1-chain close to the G-domain and three in the G-domain of the α 1-chain. The most studied of these neurite outgrowth domains is a decapeptide derived from the carboxy terminal part of the γ 1-chain of laminin-1. This domain is involved in neuronal migration and axon guidance, and has a dual neurotrophic/neurotoxic role. Synthetic peptides derived from this domain have a neurotoxic effect when applied to neurons at high (micromolar) concentrations whereas low (nanomolar) concentrations promote neurite outgrowth and neuronal survival in both soluble and substrate-bound forms.

After the neurite outgrowth domain of the γ 1-chain of laminin-1 was identified, the A β peptide, a major constituent of plaques in Alzheimer's disease, was also shown to have a concentration dependent dual neurotrophic/neurotoxic effect. This effect was thought to result in neuronal degeneration and/or sprouting of neurites detected in the plaques. Thus, it became relevant to investigate the distribution of laminin-1 and its γ 1-chain peptide in Alzheimer's disease. Neurological mutants with known defects in neuronal survival and migration, such as the weaver mutant mouse, offer suitable model systems to study the molecular mechanism of neuronal death and migration defects. We used both normal and

weaver mutant mice to elucidate the possible roles of laminin-1, its γ 1-chain and tissue plasminogen activator in the migratory failure of the homozygous weaver granule neurons, and in death of the weaver cerebellar granule neurons.

Few attempts have been made to use laminin-1 or its neurite outgrowth promoting γ 1-chain decapeptide in repair of CNS injuries. In PNS injuries, a number of different laminin-1 grafts have been applied. In PNS, both laminin-1 and the γ 1-chain decapeptide allowed neurorraphy and reduced autotomy pain caused by sciatic nerve injuries. However, the laminin-grafts were thick and caused severe compression, if used to repair thin nerves. Therefore, they could only be used on large nerves. We initiated trials to develop more suitable thin materials that could be used to form the back bone of the grafts for attachment of laminin-1 and its biologically active peptides. These grafts could then be tested in attempts to regenerate both peripheral nerve and spinal cord injuries.

Laminin-1 has been shown to induce neuronal differentiation of both early chick neural tube cells and retinal neuroepithelial cells. Neuronal differentiation is finalized by cessation of cell division by the fully differentiated neurons. Molecules and mechanism involved in neuronal differentiation are poorly understood. In recent years, the ability to isolate and maintain neuronal precursor cells in culture has allowed molecular and biochemical analysis of neuronal differentiation. Teratocarcinoma cells have been used as alternative model systems in studies on early neuronal development. For example, the mouse F9 teratocarcinoma cells have been shown to differentiate into neuron when cultured in the presence of both retinoic acid (RA) and dibutyryl cyclic adenosine monophosphate (dbcAMP) in low serum concentrations. However, neuronal differentiation of the F9 cells occurs in a subclass of the cells while a large proportion of the cells still remain undifferentiated. To enable the use of F9 cells in molecular studies on neuronal differentiation, we subcloned the F9 cells into a homogeneous population of cells that expressed neuronal characteristics on regular tissue culture plastics without any inductive agents. This novel cell line will be used in further studies to investigate the mechanisms of neuronal differentiation.

REVIEW OF LITERATURE

1. Laminin

1.1. General features of laminins

Laminins form a growing family of large secretory glycoproteins with diverse functions and cellular distributions. The first, and best characterized laminin molecule, laminin-1, was purified from the Engelbreth-Holm-Swarm tumor (Timpl *et al* 1979). This prototype of laminin consists of three different, but related polypeptide chains, A (400 kDa), B1 (210 kDa) and B2 (200 kDa), all coded by different genes (Beck *et al* 1990; Engel *et al* 1993) (Figure 1). Laminin-1 is a constituent of basement membranes, and antibodies raised against laminin-1 cross-react with laminins from various tissues and species. At present, 14 different laminin chains have been cloned (Table 1) and shown to assemble into 12 different heterotrimeric molecules (Table 2). Currently, there are five different α -chains, three different β -chains and six different γ -chains, including netrins 1-3. The domain assembly of the different laminins is shown in Figure 2.

Table 1. Different chains of laminin and their respective genes in humans

Chain	Gene	Reference
$\alpha 1$	LAMA1	Nissinen <i>et al.</i> , 1991
$\alpha 2$	LAMA2	Vuolteenaho <i>et al.</i> , 1994
$\alpha 3A$	LAMA3A	Ryan <i>et al.</i> , 1994
$\alpha 3B$	LAMA3B	Ryan <i>et al.</i> , 1994
$\alpha 4$	LAMA4	Iivanainen <i>et al.</i> , 1995
$\alpha 5$	LAMA5	Durkin <i>et al.</i> , 1997 (partial)
$\beta 1$	LAMB1	Pikkarainen <i>et al.</i> , 1987
$\beta 2$	LAMB2	Wewer <i>et al.</i> , 1994
$\beta 3$	LAMB3	Gerecke <i>et al.</i> , 1994
$\gamma 1$	LAMC1	Kallunki <i>et al.</i> , 1991
$\gamma 2$	LAMC2	Kallunki <i>et al.</i> , 1992
$\gamma 3$	LAMC3	Koch <i>et al.</i> , 1999
$\gamma 4$ (netrin-1)	LAMC4	Meyerhardt <i>et al.</i> , 1999
$\gamma 5$ (netrin-2)	LAMC5	van Raay <i>et al.</i> , 1997

Table 2. Chain assembly of laminins.

Old name	New name	Chain assembly	Reference
EHS-laminin	laminin-1	$\alpha 1\beta 1\gamma 1$	Timpl <i>et al.</i> , 1979
merosin	laminin-2	$\alpha 2\beta 1\gamma 1$	Ehrig <i>et al.</i> , 1990
S-laminin	laminin-3	$\alpha 1\beta 2\gamma 1$	Engvall <i>et al.</i> , 1990
S-merosin	laminin-4	$\alpha 2\beta 2\gamma 1$	Engvall <i>et al.</i> , 1990
kalinin/ nicein/ epiligrin	laminin-5	$\alpha 3\beta 3\gamma 2$	Rousselle <i>et al.</i> , 1991
K-laminin	laminin-6	$\alpha 3\beta 1\gamma 1$	Marinkovich <i>et al.</i> , 1992
KS-laminin	laminin-7	$\alpha 3\beta 2\gamma 1$	Champlaud <i>et al.</i> , 1996
	laminin-8	$\alpha 4\beta 1\gamma 1$	Miner <i>et al.</i> , 1997
	laminin-9	$\alpha 4\beta 2\gamma 1$	Miner <i>et al.</i> , 1997
	laminin-10	$\alpha 5\beta 1\gamma 1$	Miner <i>et al.</i> , 1997
	laminin-11	$\alpha 5\beta 2\gamma 1$	Miner <i>et al.</i> , 1997
	laminin-12	$\alpha 2\beta 1\gamma 3$	Koch <i>et al.</i> , 1999
netrin-1	laminin-13	$\gamma 4$	Serafini <i>et al.</i> , 1994
netrin-2	laminin-14	$\gamma 5$	Serafini <i>et al.</i> , 1994
netrin-3	laminin-15	$\gamma 6$	Wang <i>et al.</i> , 1999

1.2. Nomenclature of laminins

The new nomenclature of laminins (Burgeson *et al.*, 1994) was established to allow the identification of a growing number of proteins composed of different α , β and γ subunits. The first cloned chains of laminin, A, B1 and B2 chains, were re-named as $\alpha 1$, $\beta 1$ and $\gamma 1$ chains, respectively. The newly identified laminin chains, bearing homology to the first α , β , and γ chains, are then named as α , β and γ 2,3,4 etc. in the chronological order of

publication. The EHS-tumor laminin is re-named as laminin-1 ($\alpha 1\beta 1\gamma 1$). The laminin containing a variant A-chain, merosin, is called laminin-2 ($\alpha 2\beta 1\gamma 1$), and that containing a variant B1 chain, s-laminin, is called laminin-3 ($\alpha 1\beta 2\gamma 1$). S-merosin, a variant of both A- and B-chains, is laminin-4 ($\alpha 2\beta 2\gamma 1$). Kalinin/nicein, a variant of all three chains is laminin-5 ($\alpha 3\beta 3\gamma 2$), k-laminin, another variant of the A chain is laminin-6 ($\alpha 3\beta 1\gamma 1$), and ks-laminin that has both variant A- and B1-chains is laminin-7 ($\alpha 3\beta 2\gamma 1$; Burgeson *et al* 1994).

1.3. Structure of laminin-1

Rotatory shadowing electron microscopy revealed laminin-1 as a cross-shaped molecule with one long and three short arms (Engel *et al.*, 1981). The primary structures of both mouse and human laminin-1 (Pikkarainen *et al.*, 1987; Pikkarainen *et al.*, 1988; Nissinen *et al.*, 1991; Sasaki *et al.*, 1987; Sasaki and Yamada, 1987; Sasaki *et al.*, 1988) are consistent with the ultrastructure of the purified protein (Sasaki *et al.*, 1988). The polypeptide chains of laminin-1 consist of several independent domains. The $\alpha 1$ -chain has nine distinct domains (Sasaki *et al.*, 1988), the $\beta 1$ -chain is composed of seven different domains (Sasaki *et al.*, 1987), and the $\gamma 1$ -chain has six unique domains (Sasaki and Yamada, 1987; Figure 1). The primary structural data suggest that laminin-1 consists of three closely related short arms, each composed of the N-terminal regions of the $\alpha 1$, $\beta 1$ or $\gamma 1$ chains (domains III-VI), one long arm (domains I and II) in which all the three chains associate into one rod like structure, and a globular G-domain present in the $\alpha 1$ -chain only (Figure 1). The three polypeptide chains ($\alpha 1$, $\beta 1$, and $\gamma 1$) of laminin-1 are linked to each other by disulfide bonds in domain II near the center of the laminin cross and in domain I near the globular C-terminal part of the $\alpha 1$ -chain (Engel 1993).

Laminin-1 contains a large number of cysteine-rich motifs in its short arms. These motifs bear a structural homology to epidermal growth factor (EGF; Engel, 1989). The $\alpha 1$ -chain has three cysteine-rich motifs in domains IIIa, IIIb and V (Sasaki *et al.*, 1988). In the $\beta 1$ - and $\gamma 1$ chains two cysteine-rich repeats are located in domains III and V (Sasaki *et al.*, 1987; Sasaki and Yamada, 1987).




Figure 1. The domain structure and proteolytic fragments of laminin-1. The short arms of all the laminin chains have the same domain organization. Each short arm has two kinds of globular domains, VI and IV, which are separated by rod-like domains V and III. According to sequence analysis, domain VI has a mixture of α -helix, β -sheet and random coil structures. Domain VI has several cysteine residues. Domain IV has two unrelated structures, IV typical for most α and γ chains and IV' specific for the β -chains. The IV domains, like the VI domains, have a mixture of α -helix, β -sheet and random coil structures, but no cysteines to allow the formation of disulfide bonds. The α -chain has two IV- domains, IVa and IVb. The rod-like domains V and III have cysteine-rich repeats, that show structural similarity to the EGF-like repeats, with the exception that the laminin repeats have eight cysteines, whereas the EGF-like repeats have only six cysteines. The cysteine repeats of laminin are named "laminin-type cysteine-rich repeats". The long arm is a heterotrimeric rod-like structure composed of all three chains forming domains I and II and a globular G domain composed of α -chain only. In the β -chain there is also an α -domain between domains I and II. The proteolytic enzymes used for limited proteolysis of laminin to produce various proteolytic fragments are pepsin (P), pancreatic elastase (E) trypsin and cathepsin G. The proteolytic fragments are named according to the enzyme used for proteolysis.

Laminin-1 is a highly glycosylated protein with a carbohydrate content of approximately 13-15% (Chung *et al.*, 1979; Arumugham *et al.*, 1986) that accounts for up to 25-30% of its molecular weight (Knibbs *et al.*, 1989). There are 74 potential N-glycosylation sites in the laminin-1 molecule (Beck *et al.*, 1990). A functional role for glycosylation of laminin has been reported in tumor cell adhesion, cell spreading, neurite outgrowth and integrin-laminin interactions (Dennis *et al.*, 1984; Bouzon *et al.*, 1990; Dean *et al.*, 1990; Chandrasekaran *et al.*, 1991; Chammas *et al.*, 1991). Some of these functions have also been reported for synthetic peptides derived from the laminin-1 sequence devoid of the sugar chains (see Table 4 for references). Thus, it is possible that glycosylation may have a regulatory role in multiple functions of laminin-1.

A short sequence (up to 100 to 200 amino acid residues) at the carboxy terminus of each chain is required for the assembly of laminin-1 polypeptide chains into double- and triple-stranded coiled-coil structures (Utani, *et al.*, 1994, 1995b). The β and γ chains first form a dimeric structure followed by a more stable trimer formation with the α chain (Peters *et al.*, 1985; Utani, *et al.*, 1994, 1995b; Nomizu *et al.*, 1996).

Thusfar, primary structures of human $\alpha 1$ - $\alpha 5$ (partial), $\beta 1$ - $\beta 3$ and $\gamma 1$ - $\gamma 5$ chains have been identified (for references see Table1). The primary structures of the $\alpha 1$ - $\alpha 5$, $\beta 1$ - $\beta 3$, $\gamma 1$ - $\gamma 4$ and $\gamma 6$ chains of the rodent laminins are sequenced (Sasaki *et al.*, 1988; Bernier *et al.*, 1994, Galliano *et al.*, 1995; Miner *et al.*, 1997; Liu and Mayne, 1996; Miner *et al.*, 1995; Sasaki *et al.*, 1987; Hunter *et al.*, 1989a; Utani *et al.*, 1995a; Sasaki and Yamada 1987; Sugiyama *et al.*, 1995; Iivanainen *et al.*, 1999; Serafini *et al.*, 1996; Wang *et al.*, 1999). The only laminin gene found to encode for more than a single polypeptide chain is the laminin $\alpha 3$ gene (Ryan *et al.*, 1994). The $\alpha 3$ B-chain lacks domains V and VI (Ryan *et al.*, 1994). In addition to domains V and VI, the $\alpha 3$ A- and $\alpha 4$ -chains miss also domains IVb, IIIb and IVa (Ryan *et al.*, 1994; Iivanainen *et al.*, 1995). The $\beta 3$ -chain does not have a domain IV' found in the $\beta 1$ - and $\beta 2$ chains (Gerecke *et al.*, 1994). The $\gamma 2$ - and $\gamma 3$ -chains are devoid of the globular domain VI (Kallunki, *et al.*, 1992; Koch *et al.*, 1999). The $\gamma 4$ -6 chains (netrins 1-3) differ from the other γ -chains by having only the domains VI and V in addition to the C-terminal domain unrelated to other laminins (Meyerhardt *et al.*, 1999; van Raay, *et al.*, 1997; Wang *et al.*, 1999). It is also believed, although not verified, that netrins are secreted from cells as a single polypeptide chain. The domain organization of different laminin chains is presented in Fig. 2. All variants of the α -, β - and γ -chains have

structural homology within domains I and II, although sequences coding for these domains are only 20-40% conserved (Engvall and Wewer, 1996). The sequence of the domain VI is the most highly conserved amongst the various laminin chains and amongst the same subunits from different species (Engvall and Wewer, 1996). The sequence of the globular IV domain is also highly conserved (Engvall and Wewer, 1996).




Figure 2. The domain (I-IV) organization of different laminin subunits.

1.4. The biological functions of laminin

The importance of laminins in mammalian development is emphasized by the fact that laminin-1 is the first extracellular matrix protein present in between cells of the developing mouse embryo. This was first demonstrated at the morula stage (Leivo *et al.*, 1980), and more recently by using different antibodies, at the 2-cell stage (Cooper and McQueen, 1983; Dziadek and Timpl, 1986). Expression of the specific chains of laminin is spatially and temporarily regulated, which means that each chain is present in a particular tissue at defined periods of time (Aumailley and Krieg, 1996). For example, the laminins in the adult brain are largely components of basement membranes whereas in developing brain laminins exist also in a soluble less polymerized state (Liesi, 1990; Edgar, 1991). The differential expression of the various subunits of laminin has been demonstrated for both developing and adult nervous tissue (Liesi, 1990; Edgar, 1991; Hunter *et al.*, 1992a,b; Jucker *et al.*, 1996; Luckenbill-Edds, 1997; Raabe *et al.*, 1997; Powell *et al.*, 1998; Table 5). The laminins of both *Drosophila* (Fessler *et al.* 1987) and sea urchin (McCarthy *et al.* 1987) resemble mammalian laminin-1 both structurally and functionally. Goldfish and frog central nervous system (CNS) laminins are antigenically close enough to the mammalian laminin-1 to be recognized by antibodies against mouse laminin-1 (Liesi 1985b). Leech laminin β 1-chain shows homology to human, *Drosophila*, and mouse laminin β 1-chains (Luebke *et al.*, 1995). Furthermore, both the mammalian netrin genes and the *unc-6* genes of *C. elegans* code for proteins that are homologues of the γ 1-chain of laminin-1 (Serafini *et al.*, 1994; Wang *et al.*, 1999; Hedgecock *et al.*, 1990; Ishii *et al.*, 1992). These results indicate that laminins are highly conserved molecules throughout evolution. The high phylogenetic preservation of laminins implies that these proteins serve in fundamental roles throughout the species.

Laminins are essential for the architecture of basement membranes by participating in their assembly, and by maintaining the differentiated states of epithelial and endothelial cell layers, intimately associated with the basement membranes (Yurchenco *et al.*, 1992; Yang *et al.*, 1998). Formation of a laminin-1 and type IV collagen network is considered a major event in the supramolecular organization of basement membranes (Timpl and Brown, 1994; Timpl, 1996). Laminin-collagen networks are thought to be stabilized by other molecules, such as nidogen (Dziadek, 1995), heparan sulfate proteoglycans (Timpl, 1994), agrin (Denzer *et al.*, 1995), perlecan (Iozzo *et al.*, 1994), fibulin-1 (Sasaki *et al.*, 1995b), fibulin-2 (Sasaki *et al.*, 1995a), and BM-40 (Hohenester *et al.*, 1996), an anti-adhesive glycoprotein involved in tissue remodeling.

The identified functions of laminins include promotion of attachment and spreading of various cell types, such as hepatocytes (Johansson *et al.*, 1981), fibroblasts (Couchman, *et al.*, 1983), tumor cells (Vlodavsky and Gospodarowicz, 1981; Terranova *et al.*, 1982, 1983; Malinoff *et al.*, 1983), and neuronal cells (Baron van Evercooren, 1982; Liesi *et al.*, 1984a). Laminin-1 also stimulates the growth of Schwann cells (McGarvey *et al.*, 1984), primary renal cortical tubular epithelial cells (Oberley and Steinert, 1983), F9 cells (Rizzino *et al.*, 1980), and bone-marrow-derived macrophages (Ohki and Kohashi, 1994). Laminin-1 can stimulate DNA synthesis and growth of cells possessing receptors that are functionally or topologically associated with the EGF receptors (Panayotou *et al.*, 1989). Laminin-1 also functions in promoting epithelial cell polarity (Martin and Timpl, 1987; Klein *et al.*, 1988), neurite outgrowth (Baron van Evercooren *et al.*, 1982; Liesi *et al.*, 1984a; 1989), axon guidance (Liesi and Silver, 1988; Letourneau *et al.*, 1988), neuronal migration (Liesi 1985a, Liesi *et al.*, 1995), and regeneration of both central and peripheral nervous systems (Hopkins *et al.*, 1985; Liesi 1985b; Madison *et al.*, 1985, 1987; Kauppi, 1993). Laminin-1 induces the formation of capillary-like structures by endothelial cells *in vitro* (Kramer *et al.*, 1989), and influences macrophage development and function by inducing secretion of cytokines, such as IL-6 and TNF-alpha (Armstrong and Chapes, 1994).

Laminins are involved in wound healing, and regeneration of several non-neuronal tissues. E.g. laminin-5 expression is upregulated in skin wounds (Ryan *et al.*, 1994). The $\alpha 3$ -chain (Ryan *et al.*, 1994; Goldfinger *et al.*, 1999) and $\gamma 2$ -chain (Kainulainen *et al.*, 1998) of laminin-5 are shown to be essential for epithelial migration and wound closure. The interaction of hepatocytes and laminin-1 has also been shown to be important in liver regeneration, possibly in growth stimulation of hepatocytes and/or maintenance of hepatocyte-specific functions (Kato *et al.*, 1992; Wewer *et al.*, 1992). During liver regeneration, the level of the $\alpha 1$ -chain expression increases progressively from 12 hours after partial hepatectomy to the end of the regenerative period (Giménez *et al.*, 1995). Expression of the $\beta 1$ - and $\gamma 1$ -chains increases rapidly during the first day after resection of the liver, and decreases gradually reaching normal levels at the end of the regenerative period (Giménez *et al.*, 1995). Furthermore, during the regenerative period of intestinal epithelia, laminin $\beta 1$ - and $\gamma 1$ -chains are expressed differentially indicating their role in the healing process (Goke *et al.*, 1996).

In addition to the function of laminins in normal tissues, laminins and their cellular receptors may have a significant role in tumor cell growth and metastasis (Yamamura *et al.*, 1993; Menard *et al.*, 1998). Laminin-1 has been shown to promote metastatic activity of tumor cells by stimulating their attachment and migration as well as by promoting their degradation of extracellular matrix (Yamamura *et al.*, 1993; Menard *et al.*, 1998). Several synthetic peptides derived from laminin-1 play a role in tumor cell metastasis (see Table 4). E.g. the YIGSR peptide, derived from the short arm of $\beta 1$ chain of laminin-1, has been shown to inhibit tumor cell growth and metastasis (Nomizu *et al.*, 1993; Iwamoto *et al.*, 1996). Another $\beta 1$ -chain peptide, PDSRG, has also been shown to inhibit tumor cell metastasis (Kleinman, *et al.*, 1989). The RGD peptide, derived from the cell recognition site of fibronectin inhibits tumor invasion and metastasis (Saiki *et al.*, 1990; Fujii *et al.*, 1996). This RGD sequence is also identified in the $\alpha 1$ -chain of laminin-1 (Tashiro *et al.*, 1991).

A number of point mutations in the various chains of laminins have been linked to human diseases. For example, mutations in $\alpha 3$, $\beta 3$ and $\gamma 2$ -chains of laminin-5 have been reported in patients with junctional epidermolysis bullosa (McGrath *et al.*, 1995; Aumailley and Grieg, 1996; Aumailley and Smyth, 1998). The dy^{2j} mice with a truncated domain VI of the $\alpha 2$ -chain of laminin-2 lack a structurally identifiable basement membrane in their peripheral nerves (Xu *et al.*, 1994). In humans, mutations in domains VI and I of the laminin-2 $\alpha 2$ -chain (laminin-2) gene have been found in patients with congenital muscular dystrophy (CMD; Helbling-Lecler *et al.*, 1995; Hayashi *et al.*, 1997; Gullberg *et al.*, 1999). Complete absence of the $\alpha 2$ -chain is a characteristic of some congenital muscular dystrophies (Tomé *et al.*, 1994; Gullberg *et al.*, 1999). The $\alpha 2$ -chain deficiency of CMD is also reported to lead to extensive brain abnormalities (Sunada *et al.*, 1995). This result suggests that the $\alpha 2$ -chain of laminin is important in the development of CNS. In *Drosophila*, mutations of the $\alpha 1$ -chain, the only α -chain identified in this species, is embryonically lethal (Henchcliffe *et al.*, 1993), leading to defects in numerous tissues, such as brain, heart, gut and somatic muscles (Yarnitzky and Volk, 1995; Garcia-Alonso *et al.*, 1996).

Attempts to produce knockouts for most chains of laminins have failed, which further emphasizes their fundamental importance for embryonic development. The $\alpha 2$ -chain null mutant mice are characterized by growth retardation and symptoms of severe muscular dystrophy. These animals die by the age of 5 weeks (Miyagoe *et al.*, 1997). In the $\alpha 3$ -chain deficient mice the formation of hemidesmosomes is perturbed and functional interaction between laminin-5 ($\alpha 3\beta 3\gamma 2$) and integrin $\alpha 6\beta 4$ is disrupted (Ryan *et al.*, 1999). The $\alpha 3$ -chain deficient mice die 2-3 days after birth (Ryan *et al.*, 1999). Mice lacking the $\alpha 5$ -chain of laminin have multiple developmental defects, including a failure of closure of the anterior neural tube (Miner *et al.*, 1998). Embryos lacking laminin $\alpha 5$ -chain die late in embryogenesis (Miner *et al.*, 1998). The function of the $\beta 2$ -chain of the laminin-3 has been tested in $\beta 2$ -chain knockout mice (Noakes *et al.*, 1995a, b). Mice lacking the $\beta 2$ -chain show aberrant synaptic differentiation of their neuromuscular junctions and die within 4 weeks after birth (Noakes *et al.*, 1995a). These results suggest that laminin-3 regulates the formation of motor nerve terminals (Noakes *et al.*, 1995a). The $\beta 2$ -chain deficient knockout mice also show impaired glomerular ultrafiltration and kidney function (Noakes *et al.*, 1995b) as well as abnormal retinal development (Libby *et al.*, 1999). Targeted deletions of the $\gamma 1$ -chain in mouse embryonic stem cells cause early embryonic lethality (Smyth *et al.*, 1999). The $\gamma 1$ -chain-deficient embryos lack basement membranes and die by the embryonic day 5.5 (Smyth *et al.*, 1999). Netrin-1 knockout mice exhibit defects in spinal commissural axonal projections and several forebrain commissures and die within a few days (Serafini *et al.*, 1996). Mice lacking a functional netrin-1 receptor (DCC, deleted in colorectal cancer) have similar defects in their axonal projections as shown in netrin-1 knockout mice (Fazeli *et al.*, 1997; Deiner and Sretavan, 1999).

1.5. Analysis of the functions of laminin-1 using proteolytic fragments and synthetic peptides

Several studies have identified specific functional domains of laminin-1 polypeptide chains. Initially such functional domains were identified by isolating proteolytically cleaved and biochemically purified fragments of laminin-1 (Timpl and Martin, 1987; Paulsson, 1992). The location and biological functions of these proteolytically cleaved fragments are summarized in Table 3.

Table 3. Functions of proteolytic fragments of laminin. Abbreviations C, E, P and T indicate the proteolytic enzyme used; C, chymotrypsin; E, elastase; P, pepsin and T, trypsin. See Fig 1. for positions of proteolytic fragments in the laminin-1 molecule.

Proteolytic fragment	Functions of the fragment	Reference
P1	Cell attachment, Type IV collagen binding Nidogen binding Mitogenic	Aumailley <i>et al.</i> , 1987 Paulsson <i>et al.</i> , 1987 Panayotou <i>et al.</i> , 1989
E1	Mitogenic	Panayotou <i>et al.</i> , 1989
E4	Inhibition of Ca^{2+} induced aggregation of laminin	Schittny and Yurchenco, 1990
C1-4	Ca^{2+} dependent polymerization	Bruch <i>et al.</i> , 1989
C8-9	Not detected	Bruch <i>et al.</i> , 1989
E8	Cell attachment Promotion of neurite outgrowth Heparin binding	Aumailley <i>et al.</i> , 1987 Edgar <i>et al.</i> , 1984 Edgar <i>et al.</i> , 1984
T8	Cell attachment	von der Mark, <i>et al.</i> , 1991
E3, C3	Antibodies against 25K inhibit neurite outgrowth Binding of heparin and heparansulfate	Edgar <i>et al.</i> , 1988 Ott <i>et al.</i> , 1982

More accurate information on the specific functional domains of laminin-1 has been obtained using synthetic peptides. The peptides, their functions, and their positions within the laminin-1 molecule are listed in Table 4. Some of the identified peptides exhibit multifunctional properties.

Neurite outgrowth function of laminin-1 has been mapped to five different peptide domains. One is in the domain I of the γ 1-chain (Liesi *et al.*, 1989b), another in the α 1-chain just before the G-domain (Tashiro *et al.*, 1989), and the remaining three in the domain G of the α 1-chain (Skubitz *et al.*, 1991). Other neuronally active peptides have been localized in the β 2-chain of laminin-3 (LRE; Hunter *et al.*, 1989b, 1991).

Table 4. Some of the biologically active synthetic peptides of laminins and their functions. Abbreviations for amino acids: A alanine, R arginine, N asparagine, D aspartic acid, C cysteine, Q glutamine, E glutamic acid, G glycine, H histidine, I isoleucine, L leucine, K lysine, M methionine, F phenylalanine, P proline, S serine, T threonine, W tryptophan, Y tyrosine, V valine.

	Peptide	Function	Reference
α 1 I	CSRARKQAAS <u>IKVAV</u> SADR	Cell adhesion, neurite outgrowth Stimulation of metastasis, collagenase production Bone cell differentiation Binding of 110 kDa cell surface protein Stimulation of plasminogen activation Signal transduction, cell growth Promotion of angiogenesis and tumor growth Tumor growth, colony formation Binding of APP T-lymphocyte adhesion Disruption of gastrulation in sea urchin	Tashiro <i>et al.</i> , 1989 Kanemoto <i>et al.</i> , 1990 Vukicevic <i>et al.</i> , 1990 Kleinman <i>et al.</i> , 1991 Stack <i>et al.</i> , 1991 Kubota <i>et al.</i> , 1992 Kibbey <i>et al.</i> , 1992 Yamamura <i>et al.</i> , 1993 Kibbey <i>et al.</i> , 1993 Weeks <i>et al.</i> , 1994 Hawkins <i>et al.</i> , 1995
α 1 G α 1 III	RGD	Endothelial differentiation Cell adhesion Cell adhesion, spreading.	Grant <i>et al.</i> , 1989 Aumailley <i>et al.</i> , 1990 Tashiro <i>et al.</i> , 1991
α 1 IVa	LSNIDYILIKAS	Metastasis promotion	Kuratomi <i>et al.</i> , 1999
α 1 IVb	RDQLMTVLAN VT	Metastasis promotion	Kuratomi <i>et al.</i> , 1999
α 1 IVb	SINNTAVMQRL T	Metastasis promotion	Kuratomi <i>et al.</i> , 1999
α 1 VI	RQVFQVAYIIK A	Metastasis promotion	Kuratomi <i>et al.</i> , 1999

$\alpha 1$ G	KQNCLSSRASFRGCVRNLRLSR	$\alpha 3\beta 1$ integrin binding	Gehlsen <i>et al.</i> , 1992
$\alpha 1$ G	SINNNR	Alveolar formation, cell adhesion Cell adhesion	Matter and Laurie, 1994 Chen <i>et al.</i> , 1997b
$\alpha 1$ G	KATPMLKMRTSFHGCIK,	Cell adhesion, heparin binding, neurite outgrowth	Skubitz <i>et al.</i> , 1991
$\alpha 1$ G	KEGYKVRLNI TLEFRTTSK	Cell adhesion, heparin binding, neurite outgrowth Binding of $\alpha 3\beta 1$ integrin	Skubitz <i>et al.</i> , 1991 Pattaramalai <i>et al.</i> , 1996
$\alpha 1$ G	KNLEISRSTFDL LRNSYGRK	Cell adhesion, neurite outgrowth	Skubitz <i>et al.</i> , 1991
$\alpha 1$ G	DGKWHTVKTE YIKRKAF	Cell adhesion, neurite outgrowth	Skubitz <i>et al.</i> , 1991
$\alpha 1$ G	RKRLQVQLSI RT	Neurite outgrowth Metastasis promotion Inhibition of branching epithelial morphogenesis Stimulation of matrix metalloproteinase secretion	Richard <i>et al.</i> , 1996 Kim <i>et al.</i> , 1998 Kadoya <i>et al.</i> , 1998 Weeks <i>et al.</i> , 1998
$\alpha 2$ G	KNRLTIELEVR T	Neurite outgrowth	Richard <i>et al.</i> , 1996
$\beta 1$ III	YIGSR	Cell adhesion, chemotaxis, binding to 67 kDa receptor, neuronal attachment Inhibition of metastasis Bone cell differentiation Endothelial differentiation Seroli cell morphogenesis Inhibition of angiogenesis and tumor growth Disruption of gastrulation in sea urchin	Graf <i>et al.</i> , 1987 Iwamoto <i>et al.</i> , 1987 Nakai <i>et al.</i> , 1992 Yamamoto <i>et al.</i> , 1994 Vukicevic <i>et al.</i> , 1990 Grant <i>et al.</i> , 1989 Hadley <i>et al.</i> , 1990 Sakamoto <i>et al.</i> , 1991 Hawkins <i>et al.</i> , 1995
$\beta 1$ III	PDSRG	Cell adhesion, inhibition of metastasis	Kleinman <i>et al.</i> , 1989
$\beta 1$ III	YGYG DALR	$\alpha 2\beta 1$ integrin binding	Underwood <i>et al.</i> , 1995

β 1 IV	RYVVLPRPVCF EKGMNYTVR	Heparin binding, cell adhesion	Charonis <i>et al.</i> , 1988
β 1 VI	RIQNLLKITNLR IKFVK	Heparin binding	Kouzi-Koliakos <i>et al.</i> , 1989
β 1 V	LGTPG	Binding to 67 kDa elastin receptor	Mecham <i>et al.</i> , 1989
β 2 I	LRE	Motoneuron stop signal, inhibition of neurite outgrowth Promotion of motor axon growth	Hunter <i>et al.</i> , 1989b,1991 Brandenberger <i>et al.</i> , 1996
γ 1 I	RNIAEIIKDI	Neurite outgrowth, neurotrophic effect, neurotoxic effect Neuronal migration Nuclear translocation Required for di/trimerization of laminin Axonal differentiation Axon guidance Modulation of electrical activity of neurons	Liesi <i>et al.</i> , 1989 Liesi <i>et al.</i> , 1995, 1996 Liesi <i>et al.</i> , 1995 Utani <i>et al.</i> , 1994 Matsuzawa <i>et al.</i> , 1996 Matsuzawa <i>et al.</i> , 1998 Hager, <i>et al.</i> , 1998

1.6. Molecular interactions of laminin-1

Laminin-1 has the ability to self-associate into polymers by aggregation of the amino-terminal globular domains VI (Yurchenco *et al.*, 1992; Yurchenco and Cheng, 1993). This self-association requires calcium to induce the required conformational change (Yurchenco and Cheng, 1993).

Various molecules interact with laminin-1 either directly or indirectly and can therefore modulate its biological effects. Nidogen-1 (entactin) binds to the fourth EGF-like repeat of the domain III in the γ 1-chain (Mayer *et al.*, 1993) increasing hepatocyte adhesion and spreading (Levavasseur *et al.*, 1994). Nidogen-2 binds with lower affinity to the same binding site as nidogen-1 (Kohfeldt *et al.*, 1998). A basement membrane-associated glycoprotein, fibulin, binds close or directly to the G-domain of laminin-1 (Pan *et al.*, 1993). Thrombospondin binds laminin-1 directly (Frazier, 1987), as does heparansulfate proteoglycan that binds to the long arm of laminin-1 (Laurie *et al.*, 1986). Agrin, a large

multidomain heparansulfate proteoglycan, attaches to the central region of the three-stranded, coiled-coil oligomerization domain in the long arm of laminin-1 (Denzer *et al.*, 1998). Heparin has several binding sites in the laminin-1 molecule, with the most important one being in the G-domain of laminin-1 (Ott *et al.*, 1982; Laurie *et al.*, 1986). In addition, the α -chain of laminin-1 has a heparin-binding site in domain VI (Colognato-Pyke *et al.*, 1995). Electron microscopy has revealed two additional heparin-binding sites in the globular domains IV and VI of β 1-chain (Charonis *et al.*, 1988 for domain IV; Kouzi-Koliakos *et al.*, 1989 for domain VI). Also in the long arm of laminin-1 an additional heparin binding site is present in domain II close to the region where the α 1, β 1, and γ 1 chains intersect (Skubitz *et al.*, 1988). Type IV collagen has been found to bind laminin-1 either directly to the long arm (Charonis *et al.*, 1986) or indirectly via nidogen-1 (Fox *et al.*, 1991) to domain III of the γ 1-chain (Mayer *et al.*, 1993). Type VII collagen binds directly to the β 3 chain of laminin-5 using its non-collagenous, globular domain (Chen *et al.*, 1999). Although fibronectin does not bind directly to laminin-1 it can interact with laminin-1, because both laminin-1 and fibronectin bind heparin (Hayashi *et al.*, 1980; Ott *et al.*, 1982; Timpl, 1989). An extracellular matrix molecule tenascin-C does not bind directly to laminin-1 (Lightner and Erickson, 1990). However, indirect binding of tenascin-C to laminin-1 or laminin-2 may occur via agrin. Agrin binds directly tenascin-C, laminin-1, and laminin-2 (Cotman *et al.*, 1999).

Apolipoprotein E binds laminin-1 directly (Huang *et al.* 1995). It increases neuronal adhesion to laminin-1 and alters neurite morphology (Huang *et al.* 1995). The amyloid precursor protein (APP) has also been shown to bind laminin (Narindrasorasak *et al.*, 1992). APP has been suggested to be a laminin receptor, and it has been found to bind to the IKVAV-sequence in the carboxyl terminus of the α 1-chain of laminin-1 (Kibbey *et al.*, 1993). Serum amyloid P (SAP), a component of all amyloid plaques (Pepys *et al.*, 1994) and a normal component of a number of basement membranes (Dyck *et al.*, 1980; Al Mutlag *et al.*, 1993), binds to laminin-1 (Zahedi, 1997). Chicken neuronal-glial cell adhesion molecule (Ng-CAM) also binds to the short arms of laminin-1 (Grumet *et al.*, 1993). The mammalian homologue of Ng-CAM, the L1-antigen, also binds to the G2 domain of laminin-1 (Hall *et al.*, 1997a). However, laminin binding of the L1-molecule take place via sulfated HNK-1 carbohydrate epitopes (Hall *et al.*, 1997a). Some sulfated glycolipids also bind laminin-1 (Roberts *et al.*, 1985; Kobayashi *et al.*, 1994; Hillery *et al.*, 1996; Hall *et al.*, 1997b). Lectins, plant proteins that have high affinity for specific sugar residues of laminin-1 (Woo *et al.*, 1990; Ozeki, *et al.*, 1995; Saarela *et al.*, 1996), and calreticulin, a molecular chaperone with lectin like properties (McDonnell *et al.*, 1996), are also known to bind to laminin-1 carbohydrates.

Several receptors that mediate interactions of various domains of laminins with cells have been identified. Generally the laminin receptors are divided into two categories, the non-integrin-type and the integrin-type cell surface receptors (Hemler, 1990; Mecham, 1991; Liesi, 1991; Timpl and Brown, 1994). The first non-integrin type laminin receptor identified is the 67 kDa laminin receptor isolated from melanoma cells (Rao *et al.*, 1983). Later this receptor was identified in several other cell types, including primary neuronal cells (Douville *et al.*, 1988). Another non-integrin type receptor, the 110 kDa receptor (Kleinman *et al.*, 1991), is localized in brains of embryonic and postnatal mice (Luckenbill-Edds *et al.*, 1995). This receptor has been proposed to be the amyloid precursor protein (APP) also binding to the IKVAV sequence of the $\alpha 1$ -chain of laminin-1 (Kibbey *et al.*, 1993). Dystroglycan- α is a non-integrin laminin receptor that binds laminin-1, laminin-2, and laminin-4 (Hemler, 1999). Binding of the laminin-dystroglycan- α complex to the cell surface is mediated by attachment of dystroglycan- α to dystroglycan- β that is a transmembrane protein (Hemler, 1999). In laminin-1, the binding site of dystroglycan- α is in the major heparin binding domain in the E3 fragment (Gee *et al.*, 1993). Cranin, a brain isoform of dystroglycan- α , (Smalheiser and Kim, 1995) binds to the laminin E3 fragment in the long arm of the $\alpha 1$ -chain (Smalheiser, 1993) and is involved in neurite outgrowth (Smalheiser and Schwartz, 1987). At least seven integrins ($\alpha_1\beta_1$, $\alpha_2\beta_1$, $\alpha_3\beta_1$, $\alpha_6\beta_1$, $\alpha_6\beta_4$, $\alpha_7\beta_1$, and $\alpha_9\beta_1$) are known to bind laminins, and the integrins $\alpha_1\beta_1$, $\alpha_2\beta_1$, $\alpha_6\beta_1$, $\alpha_6\beta_4$, $\alpha_7\beta_1$, and $\alpha_9\beta_1$ bind specifically laminin-1 (Hemler, 1991; Timpl and Brown, 1994; Forsberg *et al.*, 1994; McKerracher *et al.*, 1996).

1.7. Laminins in the nervous system

One of the best characterized functions of laminin-1 is its involvement in neuronal development (Sanes, 1989; Liesi 1990; Luckenbill-Edds, 1997). Laminin-1 promotes neurite outgrowth, neuronal migration, and nerve regeneration (Sanes, 1987; Liesi 1990; Luckenbill-Edds, 1997). Laminin-1 also plays a role in neuronal differentiation (Heaton and Swanson, 1988; Frade *et al.*, 1996). The roles of laminins in the nervous system are discussed in sections 2.1.1., 2.2.2.a, 2.3.1., 2.4.1.3. and 2.5.4.1. Knockout studies further demonstrate the significance of laminins in the nervous system development (see section 1.4.).

In adult mammalian brain, antibodies against native laminin-1 demonstrate this protein mostly in basement membranes, whereas in those CNS areas and systems that support axon growth and neuronal regeneration laminin-1 is detected in astrocytes (Liesi, 1985a). In the developing brain, laminin-1 is expressed along the routes of the migratory neurons and pioneer axons (Liesi, 1985a,b; Cohen *et al.*, 1987; Liesi and Silver, 1988; Letourneau *et al.*, 1988). Transient expression of laminin-1 is induced in reactive astrocytes of the adult injured rat brain (Liesi *et al.*, 1984). The hypothesis that induction of laminin-1 in the injured adult CNS supports regeneration comes from the fact that astrocytes of the adult rat olfactory bulb, the only regenerative area in adult mammalian brain, express laminin continuously (Liesi, 1985b). Furthermore Schwann cells of the adult peripheral nervous system (PNS) (Cornbrooks *et al.*, 1983) and astrocytes of the adult CNS of the lower vertebrates that can regenerate their injured nerves (Liesi, 1985b; Hopkins *et al.*, 1985) produce laminin-1.

Laminin-2 has been localized in retinal ganglion cells and along the optic pathway in chicks (Morissette and Carbonetto, 1995). Laminin-2 is also expressed in other parts of the CNS, e.g. in the dendritic spines of the adult mammalian hippocampus (Tian *et al.*, 1997), in neuronal fibers and structures of the limbic brain region (Hagg *et al.*, 1997) as well as in the developing mammalian olfactory system (Raabe *et al.*, 1997). In the developing mouse cerebellum laminin-2 immunoreactivity is restricted to the Purkinje cells (Powell *et al.*, 1998). In the PNS laminin-2 the predominant laminin isoform present in the endoneurial basement membranes of the peripheral nerves (Sanes *et al.*, 1990), is produced by Schwann cells (Leivo and Engvall, 1988).

Laminin-3 is concentrated in the basement membranes of the neuromuscular junctions (Hunter *et al.*, 1989a,b) but is also transiently expressed during development of the rat neocortex (Hunter *et al.*, 1992a). During CNS development laminin-3 is present in the pial surface that overlies the floor plate of the spinal cord (Hunter *et al.*, 1992a). Prenatally, laminin-3 appears transiently in the capillary basement membranes throughout the CNS (Hunter *et al.*, 1992a).

In the developing human nervous system, mRNA for all five laminin α -chains has been detected. The α -1 chain mRNA is present in the 17 wk human embryo brain in the neural retina, olfactory bulb, cerebellum, and in the meninges whereas the α -2-mRNA is detected

in the choroid plexus and the meninges (Vuolteenaho *et al.*, 1994). The mRNAs for the $\alpha 3$ -, $\alpha 4$ -, and $\alpha 5$ -chains are present in the 6-12 wk human embryo brain (Liesi, unpublished result). In 20 wk embryo brain, the $\alpha 4$ -chain mRNA is not detected (Iivanainen *et al.*, 1995). The $\beta 1$ -chain mRNA has been detected in 6-12 wk human embryo brain (Liesi, unpublished result) and the $\beta 2$ -chain in the 17 wk human embryo brain (Iivanainen *et al.*, 1994). In 18-19 wk embryos the mRNA for the $\gamma 1$ -chain and in 17 wk embryos the mRNA for the $\gamma 2$ have been demonstrated (Kallunki *et al.*, 1992). The $\gamma 4$ -chain mRNA was found already in the 6-12 wk embryo brain (Liesi, unpublished result). In the adult human CNS, the expression of laminin chains is more restricted. Expression of the mRNA for $\gamma 1$ -, $\gamma 3$ - and $\gamma 5$ -chains of laminins (I; Koch *et al.*, 1999; Van *et al.*, 1997) has been detected. In rodents the qualitative expression of laminin mRNAs is largely similar to that found in the human CNS. However, unlike the adult human CNS the mRNAs for $\alpha 4$, $\alpha 5$, and $\beta 2$ -chains are expressed in the adult mouse (for reference see Table 5b). Listing of expression of mRNAs for laminins in the human and rodent CNS is shown in Table 5.

Table 5a. Expression of mRNAs for different chains of laminins in the human CNS.

Laminin chains	mRNA expression		
$\alpha 1$	+	embryo 17 wk	Vuolteenaho <i>et al.</i> , 1994
	-	adult	I
$\alpha 2$	+	embryo 17 wk	Vuolteenaho <i>et al.</i> , 1994
$\alpha 3$	+	embryo 6-12 wk	Liesi, unpublished
$\alpha 4$	+	embryo 6-12 wk	Liesi, unpublished
	-	embryo 20 wk	Iivanainen <i>et al.</i> , 1995;
	-	adult	Iivanainen <i>et al.</i> , 1995
$\alpha 5$	+	embryo 6-12 wk	Liesi, unpublished
	-	adult	Durkin <i>et al.</i> , 1997;
$\beta 1$	+	embryo 6-12 wk	Liesi, unpublished
	+	adult	I
$\beta 2$	+	embryo 17 wk	Iivanainen <i>et al.</i> , 1994
$\beta 3$	-	embryo 6-12 wk	Liesi, unpublished
$\gamma 1$	+	embryo 18-19 wk	Kallunki <i>et al.</i> , 1992
	+	adult	I
$\gamma 2$	+	embryo 17 wk	Kallunki <i>et al.</i> , 1992
$\gamma 3$	-	embryo	Koch <i>et al.</i> , 1999
$\gamma 3$	+/-	adult	Koch <i>et al.</i> , 1999
$\gamma 4$ (netrin-1)	+	embryo 6-12 wk	Liesi, unpublished
$\gamma 5$ (netrin-2)	+	adult	Van Raay <i>et al.</i> , 1997

Table 5b. Expression of mRNAs for different chains of laminins in the rodent CNS.

Laminin chains	mRNA expression		
$\alpha 1$	+	embryo 15.5 d	Miner <i>et al.</i> , 1997
	-	adult	Miner <i>et al.</i> , 1997
$\alpha 2$	-	nb*	Bernier <i>et al.</i> , 1994
	-	adult	Miner <i>et al.</i> , 1997
$\alpha 3$	+	embryo 13-17 d	Galliano <i>et al.</i> , 1995
	-	adult	Miner <i>et al.</i> , 1997
$\alpha 4$	+	embryo 17.5d, adult	Miner <i>et al.</i> , 1997
$\alpha 5$	+	embryo 17,5d adult	Miner <i>et al.</i> , 1995
$\beta 1$	+	embryo, nb, adult	Sarthy and Fy, 1996
$\beta 2$	+	nb*, adult	Libby <i>et al.</i> , 1997
$\beta 3$	-	embryo 14 d, nb*	Utani <i>et al.</i> , 1995a
$\gamma 1$	+	embryo 14 d	Sugiyama <i>et al.</i> , 1995
$\gamma 2$	+	embryo 14d	Sugiyama <i>et al.</i> , 1995
$\gamma 3$	+	adult	Iivanainen <i>et al.</i> , 1999
$\gamma 4$ (netrin-1)	+	embryo 10,5-11,5	Serafini <i>et al.</i> , 1996
$\gamma 6$ (netrin-3)	+	embryo 11,5-14,5d	Wang <i>et al.</i> , 1999

*nb, newborn

Schwann cells of the PNS (Davis *et al.*, 1985) and astrocytes of the CNS (Liesi and Risteli, 1989) produce a variant form of laminin with the conventional $\alpha 1$ -chain missing. Recently, human glial laminin has been shown to contain a novel β -chain together with a regular $\gamma 1$ -chain and a shorter α -chain, possibly related to $\alpha 3B$ -, $\alpha 4$ -, and $\alpha 5$ -chains (LeMosy *et al.*, 1996). Glial expression of laminin-1 is shown to correlate with neuronal migration in all areas of the central nervous system investigated (Liesi 1985a). In the developing optic nerve axons are in contact with punctate extracellular deposits of laminin (Cohen *et al.*, 1987; Liesi and Silver, 1988; Letourneau *et al.*, 1988; Liesi and Risteli, 1989). The fine punctate deposits of laminin-1 localized in association with growing axons are composed of $\beta 1$ - and $\gamma 1$ -chains of laminin, whereas the $\alpha 1$ immunoreactivity is present in the basement membranes (Liesi and Risteli, 1989).

2. Nervous system development, degeneration and regeneration

In recent years a large number of molecules, involved in nervous system development, degeneration, and regeneration have been identified. The functions of these molecules, including laminins, are only partially understood and many questions remain to be answered concerning the mechanisms that allow these molecules to orchestrate the behavior of neuronal cells.

2.1. Neuronal differentiation

The differentiated cell types of the vertebrate nervous system arise from the neurogenic epithelium (neuroectoderm) of the neural plate. In recent years major advances have been made in understanding the inductive signals of neurogenesis during early embryonic development. Studies on dissociated ectodermal cells during the gastrula stage showed that these cells reaggregate after long dissociation and express neuronal markers (Godsave and Slack, 1989; Sato and Sargent, 1989). This suggested that neuronal induction may occur without any inductive signals from the dorsal mesoderm. The results indeed imply that embryonic cells would become nerve cells if they do not get any other inductive signals. Thus, the early neuronal differentiation is under a negative control. One of the molecules that inhibit neurogenesis is the bone morphogenetic protein (BMP) 4 (Wilson and Hemmati-Brivanlou, 1995). Several results indicate that the ectoderm responds to BMP-activation in a dose-dependent manner: The neural fate associates with the lowest BMP activity, the epidermal fate to the highest, and the neural crest to the intermediate level (Chitnis, 1999). The inhibitory effect of BMP is inactivated by neural inducers such as noggin, follistatin, and chordin (Sasai *et al.*, 1995). All these molecules are secreted by the Spemann organizer (Smith and Harland, 1992; Hemmati-Brivanlou *et al.*, 1995; Sasai *et al.*, 1994). These molecules may bind to BMP and block the activation of BMP receptors (Piccolo *et al.*, 1996; Zimmerman *et al.*, 1996). This suppresses BMP signaling and leads to activation of a number of genes that promote neuronal fate in the dorsal ectoderm (Sasai, 1998).

The specificity of neuroectodermal cells depends on the position they initially occupy in the neural plate. The dorsoventral development of the neural tube is mediated by two different signaling systems (Tanabe and Jessell, 1996). The dorsal specification is

mediated by BMPs that are initially expressed by epidermal ectoderm and later by the roof plate (Tanabe and Jessell, 1996). BMP signaling induces the fates of several sensory interneurons in the dorsal spinal cord. Recently genetic ablation studies have shown that the roof plate is essential for specification of certain dorsal interneurons (Lee *et al.*, 2000). The ventral specification of the neural tube is mediated by Sonic hedgehog (Shh; Tanabe and Jessell, 1996). Initially, Shh is expressed by the notochord, inducing the formation of the floor plate, where it also will be expressed. The expression pattern of Shh is closely linked to the development and differentiation of the entire ventral neural tube (Martí *et al.*, 1995). Shh expressed by the notochord defines the differentiation of motor neurons and ventral interneurons (Ericson *et al.*, 1996). Shh-deficient mice show perturbed ventral patterning of the CNS (Chiang *et al.*, 1996). Shh prevents the dorsalization of the ventral neuronal tube by antagonizing the effect of BMPs (Liem *et al.*, 1995).

During closure of the neural tube the cells in the anterior part of the neural tube begin to form the primitive forebrain, midbrain, and hindbrain. Posteriorly, the neural tube forms spinal cord. Shortly after closure of the neural tube the neuroepithelial cells proliferate rapidly. During this proliferative phase the cells can either remain as undifferentiated neuronal stem cells or they may differentiate into neurons and glial cells. The mechanisms controlling the undifferentiated and differentiated state of neurons is not fully understood. The cell surface protein Notch and its ligand Delta-1 have been shown to participate in the regulation of the choice between the undifferentiated and differentiated states of neurons (Henrique *et al.*, 1997). In different parts of the CNS different sets of homeotic genes have been found to determine the direction of cellular development. E.g. the development of the midbrain and cerebellum depends on an organizing center at the midbrain-hindbrain junction known as isthmus (Marin and Puelles, 1994). In the isthmus region several important secreted and regulatory genes are expressed, such as *Wnt-1* (McMahon and Bradley, 1990) and *En1* (Wurst *et al.*, 1994; Danielian and McMahon, 1996). Knockout experiments on these genes eliminate cerebellum either totally or to a large extent (McMahon and Bradley, 1990; Wurst *et al.*, 1994; Danielian and McMahon, 1996). *En1* may further be regulated by a secreted glycoprotein WNT (McMahon *et al.*, 1992). Expression of WNT protein is regulated by the *Pax*-gene family (Song *et al.*, 1996). Inactivation of *Pax2* leads into a massive loss of cerebellum and posterior midbrain (Favor *et al.*, 1996). Recent studies indicate that *Pax6* is expressed in the rhombic lip and in the presumptive early granule cell moving away from the rhombic lip (Engelkamp *et al.*,

1999). *Math1*, the mouse homologue of the *Drosophila* gene *atonal* encoding for a basic helix-loop-helix transcription factor, has been shown to be essential for the genesis of cerebellar granule neurons *in vivo* (Ben-Arie *et al.*, 1997). Segmentation plays a prominent role in the hindbrain (Lumsden and Krumlauf, 1996). The hindbrain neural tube is divided in segmental units called rhombomeres. Expression of Hox genes patterns an ordered set of domains along the neuraxis (Lumsden and Krumlauf, 1996). Thus, it is possible that the identity of individual rhombomeres could be defined by a cooperative action of the Hox proteins. Retinoic acid (RA) regulates the expression of Hox genes and overexpression of RA has been shown to result in an anterior shift of the Hox gene expression (Lumsden and Krumlauf, 1996).

2.1.1. Laminin in neuronal differentiation

The involvement of laminin-1 in neuronal differentiation became evident in studies on frog retinal pigment epithelial (RPE) cells (Reh 1987). RPE cells, grown on laminin-1, transdifferentiate into neurons instead of the lens cells (Reh 1987). Laminin-1 is also known to influence neuronal differentiation of the early chicken neural tube (Heston and Swanson 1988). The E8 fragment of laminin-1 stimulates proliferation and differentiation of neuroepithelial cells into neurons *in vitro* (Drago *et al.*, 1991; Frade *et al.*, 1996). Integrin $\alpha1\beta6$ receptors have been found to be down-regulated when retinal ganglion cells progressively differentiate (de Curtis *et al.*, 1991; Cohen and Johnson, 1991). Transformation of chromaffin cells into sympathetic neurons in response to basic fibroblast growth factor occurs if laminin-1 is used as culture substratum (Chu and Tolkovsky, 1994). The IKVAV domain of the $\alpha1$ -chain of laminin-1 interacts with the plasmalemmal protein LBP110 (Chalazonitis *et al.*, 1997). This protein is acquired by the neural crest-derived precursors of the enteric neurons after they colonize the gut and, therefore, promote the development of the enteric neurons (Chalazonitis *et al.*, 1997). Laminin $\beta2$ -chain may also be involved in the differentiation of rod photoreceptors (Hunter *et al.*, 1992b) and may control their developmental choices between the rod photoreceptor and bipolar cell fates (Hunter and Brunken, 1997).

2.1.2. Neuronal cell lines

Different kinds of neuronal cell lines have been developed to study the molecular mechanisms of neuronal differentiation. These cell lines may be divided in different groups depending of the origin. Teratocarcinoma cells, derived from tumors of the fetal germ cells, e.g. teratomas, provide cell lines in which the entire differentiation process from pluripotent cells to terminally differentiated cells can be followed (Martin, 1980). See also section 2.1.3

Human neuroblastoma cell lines are derived from childhood solid tumors containing of primitive cells of precursors of the autonomic nervous system. These neuroblastoma cell lines have been shown to undergo spontaneous and chemically induced differentiation into cells resembling those of the mature nervous system. Neuroblastoma cell lines have been used to study the mechanism of neuronal differentiation (Abemayor and Sidell, 1989; Pålman *et al.*, 1995; Maggi *et al.*, 1998) and degeneration (Omar and Pappola 1993; Neill *et al.*, 1994). In addition to human neuroblastomas mouse neuroblastoma cell lines have been chemically induced from both central and peripheral neurons (Liesi, 1984).

Immortalized neural cell lines can be produced using somatic cell fusion to neuroblastoma cell lines, isoforms of transforming agents such as *myc* and *neu* or by using SV40 large tumor antigen (Martinez-Serrano and Björklund, 1997). These gene transfer techniques have allowed introduction of growth factors, neuropeptides, neurotransmitters as well as biosynthetic and metabolic enzymes into the neural progenitor cells (Gökhan *et al.*, 1998). Using this technique several conditionally immortalized neural stem/progenitor cell lines have been established from embryonic hippocampus, cerebellum (Gökhan *et al.*, 1998), and spinal cord (Li *et al.*, 2000). The conditionally immortalized neural cell lines can be transplanted back into the mammalian brain and may represent an experimental resource for characterization of molecular mechanism involved in CNS development, plasticity, and regeneration (Gökhan *et al.*, 1998). Undifferentiated stem cells of the adult brain have been shown to differentiate into neurons, astrocytes, and oligodendrocytes *in vitro* (Reynolds and Weiss, 1992, 1996; Palmer *et al.*, 1997; Johansson *et al.*, 1999; Chiasson *et al.*, 1999). The neural stem cells are most abundant in the walls of the lateral ventricles (Weiss *et al.*, 1996) and in the hippocampus (Palmer *et al.*, 1997). Stem cells that associate with the ventricular systems have been shown to originate either from the ependyma (Johansson *et al.*, 1999) or the subependyma (Chiasson *et al.*, 1999).

2.1.3. Teratocarcinoma cells as a model system for neuronal differentiation

Teratocarcinoma cells have been used as a model system for early embryonic development (Strickland and Mahdavi 1978; Martin 1980,). These pluripotent cells are capable of differentiating into various organs and cell types. The inductive signals of this differentiation have been studied as putative morphogens, e.g. factors that may control cell determination during early embryonic development (Martin and Evans 1975; Sell and Pierce, 1994; Andrews, 1998). Depending on cell density laminin-1 induces differentiation of teratocarcinoma cells either into neurons or multinucleated striated muscle cells (Darmon, 1982; Sweeney *et al.*, 1990; Kim *et al.*, 1995).

Retinoic acid (RA) alone or in combination with dibutyryl cyclic AMP (dbcAMP) has been used to differentiate teratocarcinoma cells into visceral or parietal endoderm (Strickland and Mahdavi 1978, Sporn and Roberts 1983). DbcAMP alone does not provoke differentiation of the F9 cells but the cells must first be exposed to RA in order to be permissive to the effects of dbcAMP (Darrow *et al.*, 1990). Serum deprivation together with RA and dbcAMP has been shown to induce neuronal differentiation of various teratocarcinoma cell lines (Pleiffer *et al.*, 1981, McBurney *et al.*, 1982, Levine and Flynn 1986, Kubo 1989), including the F9-cells (Kuff and Fewell, 1980, Liesi *et al.*, 1983).

2.2. Neuronal migration

2.2.1. Current models of neuronal migration

The classical viewpoint on neuronal migration is that migration of embryonic and early postnatal neurons occurs via physical guidance of the radially oriented glial cell processes (Jacobson, 1991). According to this point of view all radial migration of neurons proceeds along the radial glial cells while all horizontal neuronal migration proceeds along other neuronal fibers (Rakic, 1990). Thus, the glial cells are considered to have all the molecular and physical guidance cues that the migratory neurons need (Sidman and Rakic 1973; Rakic, 1990) and to provide all the directionality required for neuronal movement.

In the past years an alternative opinion, the nuclear translocation model of neuronal migration, has emerged. It implies that neurons move without the glial guidance by nuclear translocation inside their preformed processes (Morest, 1970; Domesick and Morest, 1977; Nakatsuji and Nagata, 1989; Book and Morest, 1990; Liesi, 1992). Nuclear translocation of neuroblasts has been described as early as in 1935 in the young cerebral cortex (Sauer, 1935), but it was then thought to relate to the division of neuroblasts rather than to their migration. The fact that CNS neurons migrate via nuclear translocation inside a preformed processes was first predicted by Morest based on his careful studies on the Golgi-stained sections of different brain regions (Morest, 1970). However, the lack of direct visual evidence prevented the acceptance of the nuclear translocation model, although experiments using intraperitoneal injections of specific gliotoxins indicated that cerebellar granule neurons can migrate in the absence of the glial cells (Sotelo and Rio, 1980).

In recent time lapse video microscopy studies, the cerebellar granular neurons were seen to migrate without glial cells in association with other neurites (Nakatsuji and Nagata, 1989). When cerebellar granule neurons were cultured on a laminin substratum, the granule neurons migrated via neurite extension and contact formation followed by nuclear translocation inside preformed neuronal processes without the presence of the glia (Liesi, 1992). Novel developments of infrared video microscopy allowed the visualization of neuronal migration in slices of living postnatal cerebellum (Hager *et al.*, 1995). Using this modern methodology, the nuclear-translocation-type neuronal migration was found to be the primary form of neuronal movement in postnatal cerebellum (Hager *et al.*, 1995; Liesi *et al.*, 1995). The nuclear translocation mode of neuronal migration was found to involve two distinct phases. During the first, the neurite extension phase, granule neurons extend neurites towards the presumptive internal granule cell layer. Interestingly, the stop signal for granule neuronal migration is suggested to be their contact with mossy fibers (Baird *et al.*, 1992), their final synaptic counterparts. Thus, it is possible that the neurite-extension phase continues until the granule neurons make contact with their appropriate targets. After the contact is formed the second phase, the nuclear translocation phase, follows. The nuclei then translocate inside their preformed neurites into the internal granule cell layer. In the cerebral cortex a glia-independent form of neuronal migration has also been detected using DiI-labeling of living migratory neurons (O'Rourke *et al.*, 1992, 1995).

Confocal studies indicate that migratory cortical neurons first proceeded vertically (as if they followed the radial glial fibers), but then turned 90 degrees and moved horizontally (O'Rourke *et al.*, 1992, 1995). The horizontal mode of migration cannot be guided by the radial glia.

2.2.2. Molecules involved in neuronal migration

Molecules involved in neuronal migration belong to two primary categories: The ECM proteins and the cell adhesion molecules (CAMs). Many molecules affecting axon guidance also seem to have an effect on neuronal migration. According to the neuronal translocation model, neuronal migration and axon guidance are closely related events. Therefore it is feasible that the same molecules may be involved in both neuronal migration and axon guidance.

2.2.2.a ECM molecules

A number of ECM proteins have been shown to play a role in neuronal migration (Liesi, 1990; Sanes, 1990). In developing brain, laminin-1 is localized along the routes of migratory neurons (Liesi, 1985a). In chick CNS β 1-integrin antisense RNA inhibited neuronal migration *in vivo* (Galileo *et al.*, 1992). These results provide functional evidence for the role of laminin in neuronal migration, because β 1-integrin-type laminin receptors have been shown to mediate the effects of laminin on both central and peripheral neurons in culture (Tomaselli *et al.*, 1986; Hall *et al.*, 1987; Cohen *et al.*, 1986, 1987). Antibodies against the native EHS-tumor laminin-1 fail to inhibit neuronal migration on a laminin-1 substratum (Linder *et al.*, 1986), although recent experiments indicate that antibodies against the neurite outgrowth promoting domain of the γ 1 chain of laminin-1 inhibit neuronal migration both *in vitro* (Liesi *et al.*, 1992) and *in vivo* (Liesi *et al.*, 1995). The non-functional antibodies against native laminin-1 (used by Linder *et al.*, 1986) recognize the P1- fragment in the N-terminus of laminin-1 (Ott *et al.*, 1982), while the neurite outgrowth domains of laminin-1 localize in the C-terminal parts of this molecule (Martin and Timpl, 1987; Liesi *et al.*, 1989; Paulsson, 1992). Thus, antibodies (I) against the C-terminal neurite outgrowth promoting domain inhibit neuronal migration in cerebellar microcultures (Liesi *et al.*, 1992) and in living slices of postnatal rat cerebellum microcultures (Liesi *et al.*, 1995).

Laminin-2 promotes olfactory neuronal migration (Calof and Lander, 1991), whereas laminin-3 inhibits neuronal migration by a gated mechanism, e.g. neurons can migrate on laminin-3 but do not migrate from laminin-3-free substratum towards laminin-3 substratum (Porter and Sanes, 1995). Laminin-3 is transiently expressed along the routes of the migratory cortical neurons (Hunter *et al.*, 1992a).

Netrins, the vertebrate homologues of the UNC-6 gene and homologues of the $\gamma 1$ -chain of laminin-1, are also involved in neuronal migration (Hedgecock *et al.*, 1990; Kennedy *et al.*, 1994). In the reeler mutant mice the orderly inside-out deposition of neocortical cells is disturbed (Frotscher, 1997). The affected gene in the reeler mutation has been found to code for reelin, a large extracellular matrix protein that is secreted by the Cajal-Retzius cells (D'Arcangelo *et al.*, 1995; Frotscher, 1997). Antibodies against thrombospondin-1 inhibit granule neuronal migration by a dose-dependent manner (O'Shea, 1990). Tenascin-C either promotes or inhibits neuronal migration (Faissner and Kruse, 1990; Husmann *et al.*, 1992, 1995; Faissner, 1997). Antibodies against tenascin-C inhibit cerebellar neuronal migration (Chuong, 1990). A chondroitin sulfate proteoglycan (CSPG), astrochondrin, has been shown to be involved in cerebellar neuronal migration (Streit *et al.*, 1993). Apart from stimulation of neuronal migration (Streit *et al.*, 1993; Garwood *et al.*, 1999) CSPGs have also been shown to act as molecular barriers of neuronal migration (Grumet *et al.*, 1996; Landolt *et al.*, 1995) and their inhibitory effect can be modulated by cell adhesion molecules (Grumet *et al.*, 1996; Dou and Levine, 1995). The involvement of fibronectin in CNS neuronal migration has been proposed by studies in which fibronectin has been localized along the migratory cell pathway of embryonal cerebellar neurons (Schachner *et al.*, 1978; Hatten *et al.*, 1982). These studies have been confronted by opposite results indicating that fibronectin is not involved in cerebellum neuronal migration (Hynes *et al.*, 1986). Neuronal migration of neural crest cells has been blocked by antibodies against fibronectin (Rovasio *et al.*, 1983). The CSAT antibodies which recognize a cell surface receptor for both fibronectin and laminin (Bronner-Fraser, 1986) or peptides containing the cell-binding site of fibronectin (Boucaut *et al.*, 1984) also interfere with neural crest cells migration.

2.2.2.b Cell Adhesion molecules

Several cell adhesion molecules are involved in neuronal migration. Neural cell adhesion molecule (N-CAM), Ng-CAM, and L1 antigen (Chuong, 1990; Matsushita *et al.*, 1997; Rønn *et al.*, 1998) as well as astrotactin (Hatten and Mason, 1990; Zheng *et al.*, 1996) are all present in the cerebellum during the time of neural migration. Antibodies against all these adhesion molecules inhibit neuronal migration (Chuong, 1990; Hatten and Mason, 1990). Inactivation of N-CAM by genetic deletion have been shown to disturb the migration of several types of neurons (Cremer *et al.*, 1994; Ono *et al.*, 1994). The adhesion molecule on glia (AMOG), the β subunit of the Na/K-ATPase, is expressed in the cerebellum during the time of neuronal migration and neuronal migration is inhibited by antibodies against AMOG (Antonicek *et al.*, 1987; Gloor *et al.*, 1990).

2.2.2.c Other mediators of neuronal migration

Ion channels also play a role in neuronal migration. Blockers of both voltage-gated N-type calcium channels (Komuro and Rakic, 1992) and N-methyl-D-aspartate (NMDA) receptors (Komuro and Rakic, 1993) inhibit neuronal migration of the cerebellar granule neurons. The rate of granule neuronal migration depends on both the extracellular concentrations of Ca^{2+} and the Ca^{2+} influx through the calcium channels (Komuro and Rakic, 1992, 1993, 1996). In these experiments inhibitors of potassium and sodium channels or L- and T-type calcium channels had no effect on neuronal migration (Komuro and Rakic, 1992).

Proteolytic enzymes play a role in neuronal migration (Kalderon, 1982; Moonen *et al.*, 1982). Cultured cerebellar granule neurons secrete tissue plasminogen activator (tPA), a serine protease (Krystosek and Seeds, 1981), and bind tPA with high affinity onto their surfaces (Verrall and Seed, 1988). Both tPA activity and tPa mRNA levels of the postnatal cerebellum are maximal at P7 (Friedman and Seeds, 1995), which coincides with the most active period of granule cell migration. Inhibitors of serine proteases inhibit granule cell migration both *in vitro* and *in vivo* (Seeds *et al.*, 1990; Seeds *et al.*, 1997).

In addition to the above mentioned molecules additional gene defects and mutations may disturb the neuronal migration process (Goldowitz and Hamre, 1998). Staggerer, weaver, and reelin mice are examples of naturally occurring mutations that affect neuronal migration in the cerebellum (Sidman and Rakic, 1973; Goldowitz and Hamre, 1998). In

humans lissencephaly is a neuronal migration disorder resulting in brain malformation, epilepsy, and mental retardation (Pilz *et al.*, 1998). Two genes associated with lissencephaly have been identified. The gene for X-linked lissencephaly is called doublecortin and its function is to stabilize microtubules during neuronal migration (Allan and Walsh, 1999; Francis *et al.*, 1999; Gleeson, 1999). Another lissencephaly associated gene, LIS1, the gene for the β subunit of platelet activating factor acetylhydrolase, also influences microtubule dynamics (Sapir *et al.*, 1997). LIS1 knockout mice exhibit abnormal neuronal migration (Hirotsume *et al.*, 1998). Neuronal migration is also influenced by a number of external physical (e.g. ionizing radiation, heat), chemical (e.g. toxins, various drugs, alcohol) or biological factors (e.g. some viruses) (Jacobson, 1991).

2.2.3. Neuronal migration during cerebellar development

In the adult cerebellum five types of neurons are present: Purkinje cells, granule cells, basket cells, Golgi II cells, and stellate cells. The Purkinje cells are the only efferents of the cerebellum whereas all other neurons are interneurons. The adult cerebellum is composed of three cortical layers in which the basket and stellate cells are in the outmost layer called the molecular layer. Purkinje cells occupy the central layer, the Purkinje cell layer, and the inner cell layer, the granule cell layer, is largely formed by the granule neurons (Fig. 3). Impulses are conducted into the cerebellar cortex by climbing fibers and mossy fibers. Climbing fibers synapse with the dendrites of the Purkinje cells. Mossy fibers synapse with both dendrites of the granule cells and axons of the Golgi II cells. The axons of the granule cells, the parallel fibers, make synaptic connections with dendrites of the Purkinje, basket, Golgi II, and stellate cells. Stellate cells synapse with dendrites of the Purkinje cells, and basket cells with the somas of the Purkinje cells. The Golgi II cells synapse with the granule cell dendrites.

Cerebellar neurons originate from two separate germinal zones. The Purkinje and Golgi II cells originate from the neuroepithelial ventricular zone of the fourth ventricle. Recently the basket and stellate cells have also been shown to originate from the same germinal zone as the Purkinje cells (Zhang and Goldman, 1996). The granule cells originate from the germinal cells of the rhombic lip, the lateral most area of the presumptive floor of the 4th ventricle. These germinal cells migrate to form the external granule cell layer (EGL) on the surface of the cerebellar plate (Jacobson, 1991; Goldowitz and Hamre, 1998).




Figure 3. Organization of adult mammalian cerebellar neurons into the cortical layers; Mol = molecular layer, Pur = Purkinje cell layer, Gran = granule cell layer, BC = basket cells, CF= climbing fiber, GC = Golgi II cells, gr = granule cells, Pur= Purkinje cells, SC = stellate cells, PF=parallel fibers.

During neuronal migration neuroblasts move from their origin of birth to their final destinations. Migration of cerebellar neurons occurs in two phases. During the first phase neuroblasts migrate from the ventricular germinal zone of the fourth ventricle along the basal lamina outward to create the mantle layer of the cerebellar plate. This first migratory phase that occurs in the mouse around E11-E13 (Miale and Sidmam, 1961) gives rise to the Purkinje cells. The Golgi II cells originate from the same ventricular zone around E12-E15 and migrate directly into the presumptive molecular layer (Miale and Sidmam, 1961; Zhan and Goldman, 1996; Goldowitz and Hamre, 1998). The neuroblasts that generate granule neurons divide in the rhombic lip at E13 to E15 and migrate over the surface of the cerebellar plate to form the external granular cell layer (EGL). After having formed the

external granule cell layer these precursors of granule neurons enter a proliferative phase (Miale and Sidmam, 1961) before they enter their second phase of neuronal migration. During this second postnatal phase the granule neurons migrate from the EGL passing the Purkinje cells to form the internal granule cell layer of the mature cerebellum (Miale and Sidmam, 1961; Zhan and Goldman, 1996; Goldowitz and Hamre, 1998). The migration of granule neurons in the developing cerebellum is schematically presented in Figure 4. Progenitors of the basket and stellate cells migrate first from the primary germinal zone to the cerebellar white matter and divide there postnatally (Zhang and Goldman, 1996). Finally they enter the cerebellar cortex (Zhang and Goldman, 1996).




Figure 4. Migration of granule neurons from the external granule cell layer to the internal granule layer. Abbreviations; EGL, external granule cell layer; Mol, molecular layer; Pur, Purkinje cell layer; IGL, internal granule cell layer.

2.2.4. The weaver mutant mouse as a model of neuronal migration and death

The molecular and cellular mechanisms governing neuronal migration can be studied using neurological mutants having defects in neuronal migration. E.g, 20 mutations that alter the development of cerebellum have been identified (Sidman *et al.*, 1965). The weaver mouse is a naturally occurring, autosomally recessive mutation that has been genetically mapped to chromosome 16 in the mouse (Reeves *et al.*, 1989). The weaver mice have a neuronal defect that disturbs both cerebellar and nigro-neostriatal development (Sidman 1965, Roffler-Tarlov and Graybiel 1986). The early postmitotic granule cells are unable to migrate or form bipolar neurites (Sotelo, 1975, Rakic and Sidman 1973a,b,c). A schematic presentation of cell layers in both the weaver mutant and normal mouse cerebella is shown in Fig. 5. In the weaver mouse cerebellum a large majority of the vermal granule neurons die within the first two postnatal weeks (Rakic and Sidman, 1973c) leaving the adult weaver cerebellum low in granule neurons (Sotelo, 1975; Rakic and Sidman 1973c). Purkinje cells appear normal at the start of their postnatal life, but are later misplaced and a proportion of them die during the third postnatal week of life (Smeyne and Goldowitz, 1990; Maricich *et al.*, 1997). Neuronal degeneration in the substantia nigra (SN) is most abundant during the postnatal days 24-25 (Schmidt, *et al.*, 1982; Oo *et al.*, 1996). In addition to the disturbed cerebellar and SN development the weaver mice have a number of other abnormalities. Most of the homozygous weaver males are sterile due to of a failure in spermatogenesis (Harrison and Roffler-Tarlov, 1994). Germ cells are shown to die during the development of the testes (Harrison and Roffler-Tarlov, 1998). Furthermore, both heterozygous and homozygous weaver mice undergo tonic-clonic seizures (Eisenberg and Messer, 1989), which are thought to be the most common cause of their death. Cell death in the weaver cerebellum has been suggested to occur via apoptosis (Migheli *et al.*, 1995; Wullner *et al.*, 1995), whereas apoptosis is not the mechanisms of neuronal death in the weaver SN (Oo *et al.*, 1996). The weaver mutant mouse is used as an animal model for studies on neuronal migration, neuronal death, and neurodegenerative disorders, such as Parkinson's disease.




Figure 5. Schematic presentation of cell layers in early postnatal cerebellum of the a) normal and b) weaver mutant mouse.

The weaver gene defect impairs neuronal migration both *in vivo* and *in vitro* (Rakic and Sidman, 1973a-c; Trenkner *et al.* 1978; Willinger and Margolis, 1985). Several studies have shown that the weaver granule neurons fail to migrate and die under normal cell culture conditions (Trenkner *et al.* 1978; Willinger and Margolis, 1985). Studies using chimeric mice indicate that the weaver defect is intrinsic to the granule neurons (Goldowitz, 1989). This is further verified by *in vitro* and *in vivo* transplantation studies in which the weaver granule cells can be rescued by the wild-type granule neurons (Gao and Hatten, 1993) or their membrane extracts (Gao *et al.*, 1992). Although the weaver gene defect is intrinsic to the granule neurons the weaver Bergmann glial cells are also abnormal. They project normally, but appear immature compared to the normal Bergmann glial cells (Bignami and Dahl, 1974a,b; Sotelo and Changeux, 1974) and resemble “reactive” astrocytes by their increased expression of both GFAP and laminin (II).

Recent reports suggest that a point mutation in a G-protein coupled inwardly rectifying K^+ channel gene, GIRK2 (Lesage *et al.*, 1994), could be the weaver gene (Patil *et al.* 1995; Slesinger *et al.*, 1996). This mutation maps close to the weaver locus in chromosome 16 and is a single amino acid substitution (glycine $\rightarrow$ serine) in the pore-forming H5 region of the GIRK2 channel protein (Patil *et al.* 1995; Fig. 6.).




Figure 6. Proposed weaver-mutation in the GIRK2 channel gene. The position of the mutated amino acid (nucleotide change G→A) in H5 domain is marked with a black dot.

A point mutation in the GIRK2 channel pore domain creates a G protein-insensitive, non-selective cation channel that gates not only K^+ but also Na^+ and Ca^{2+} (Navarro *et al.*, 1996). Both GIRK2 and the related protein GIRK 1 have been localized in the granule neurons of the external granule cell layer of the normal postnatal cerebellum during the period of neuronal differentiation and neuronal migration (Slesinger *et al.*, 1996). Interestingly, the GIRK2 channel protein was not studied in the postnatal weaver cerebellum (Slesinger *et al.*, 1996).

In spite of these results it is becoming increasingly clear that the GIRK2 point mutation can not explain the weaver phenotype. The resting membrane potentials of weaver neurons can be rescued by aprotinin (II), a L-type Ca^{2+} -channel blocker verapamil (Liesi and Wright, 1995), antibodies against the γ 1-chain of laminin-1 (Liesi and Wright, 1996), and ethanol (Liesi, *et al.*, 1997). Several recent experiments have shown that the GIRK2 channel is inactive in weaver neurons during the developmental stages when the neurons are dying (Mjaatved *et al.*, 1995; Surmeier *et al.*, 1996). Furthermore, mice lacking the

GIRK2 channel gene have normal cerebella (Signorini *et al.*, 1997). As weaver granule neurons can be rescued by inhibiting the calcium entry into the neurons or chelating calcium by BABTA-AM (Liesi and Wright, 1996; Liesi *et al.*, 1997), it is possible that any measure that reduces intracytoplasmic levels of Ca^{2+} will be sufficient to prevent death of the weaver neurons. In line with this view point the weaver granule cells fail to express functional NMDA receptors (Liesi *et al.*, 1999), which helps to reduce influx of Ca^{2+} that would occur via activated NMDA receptor channels.

2.3. Neurite outgrowth and axonal targeting

During the development of the nervous system pioneer axons have to find their right targets to establish specific neuronal connections. Growth cones of growing axons responds to different guidance cues, such as soluble chemoattractants and/or chemorepellents as well as contact repulsion and/or attraction. Some of these guidance cues have been identified and found to be extracellular matrix molecules and cell adhesion molecules (Tessier-Lavigne and Goodman, 1996).

2.3.1. Laminin and other extracellular matrix molecules in neurite outgrowth

Laminin-1 is shown to promote neurite outgrowth in several *in vitro* (Baron-Van Evercooren *et al.*, 1982; Manthorpe *et al.*, 1983; Liesi *et al.*, 1984a, 1989; Edgar *et al.*, 1984; Gundersen, 1987; Lein *et al.*, 1992; Matsuzawa *et al.*, 1996; 1998) and *in vivo* studies (Liesi *et al.*, 1985b; Hopkins *et al.*, 1985; Cohen *et al.*, 1987; Letourneau, 1988; Liesi and Silver, 1988).

Five different neurite outgrowth promoting peptides, one in the $\gamma 1$ -chain (Liesi *et al.*, 1989b), and four in the $\alpha 1$ -chain (Tashiro *et al.*, 1989; Skubitz *et al.*, 1991), have been identified in the laminin-1 molecule. Axonal differentiation and directional axon growth of rat hippocampal neurons have been shown to be promoted by patterned substrates of a decapeptide derived from the $\gamma 1$ -chain neurite outgrowth domain of laminin-1 (Matsuzawa *et al.*, 1996; 1998). The neurite outgrowth promoting peptide of the $\gamma 1$ -chain of laminin-1 modulates the electrical properties of neocortical neurons (Hager *et al.*, 1998).

The first genetic evidence for the involvement of laminin in neuronal migration and axon growth came from studies on neuronal development in *C. elegans* (Hedgecock *et al.*, 1990; Ishii *et al.*, 1992). In this species a mutation of the *Unc-6* gene affects neuronal migration and pioneer axon growth indicating that *Unc-6* takes part in these processes (Hedgecock *et al.*, 1990; Ishii *et al.*, 1992). The *Unc-6* gene product is homologous to the IV and V domains of the γ -1 chain of laminin-1 and to netrins, recently identified homologues of the γ 1-chain of laminin-1 (Serafini *et al.*, 1994).



In addition to the role of laminin-1 in development of the optic pathway (Liesi and Silver, 1988) laminin-2 has also been proposed to play a role in the development of the visual system. Retinal ganglion cells (RGC) lose their ability to respond to laminin-1 during the developmental period when axons grow into the optic tectum (Cohen *et al.*, 1986), whereas laminin-2 supports neurite outgrowth of chick RGC throughout their embryonic development until the postnatal day 15 (Cohen and Johnson, 1991). Neurite outgrowth of retinal ganglion cells (RGC) on both laminin-1 and laminin-2 is blocked by anti- β 1 integrin antibodies (Cohen and Johnson, 1991). Laminin-2 promotes regeneration of sympathetic nerves *in vivo* (Anton *et al.*, 1994).

Laminin-3 is a potent inhibitor of neurite outgrowth of motor neurons (Porter *et al.*, 1995; Porter and Sanes, 1995). The β 2-chain of rat laminin-3 bears a site, LRE, that has been shown to inhibit neurite outgrowth promoted by other extracellular matrix molecules (Porter *et al.*, 1995). Interestingly, the human β 2-chain of laminin-3 lacks the LRE-site (Wewer *et al.*, 1994), which questions the importance of LRE as a general stop-signal for growing axons. Whether the LRE domain inhibits or promotes neurite outgrowth may also depend on the assembly of this peptide domain with other chains of laminin (Brandenberger *et al.*, 1996). E.g. native chick laminin-4, which contains the β 2 chain, does not inhibit but rather promotes motor axon growth (Brandenberger *et al.*, 1996).

Several other ECM molecules may either promote or suppress neurite outgrowth (Table 6). Neurite outgrowth promoting ECM molecules include fibronectin (Akers *et al.*, 1981) and trombospondin (Neugebauer *et al.*, 1991; Arber and Caroni, 1995). Collagens I, IV and VII have high neurite outgrowth promoting activities on cultured neurons from embryonic mouse brain, but they are less effective on neurons from the postnatal brain (Hirose *et al.*, 1993). Tenascin-C (Bourdon *et al.*, 1983) exhibits both axon growth-

promoting and axon growth-inhibiting activities (Grierson *et al.*, 1990; Lochter *et al.*, 1991; Faissner, 1997). Proteoglycans are also found to have a dual role in axon growth. The growth promoting effects of chondroitin sulfate (CS), heparan sulfate (HS), dermatan sulfate (DS) and keratan sulfate (KS) are dependent on the neuronal growth substratum (Campagna *et al.*, 1995; Dou and Levine, 1995; Snow *et al.*, 1996). The inhibitory effects of proteoglycans are neutralized if the relative concentration of laminin-1 is increased in relation to the inhibitory proteoglycans (McKeon *et al.*, 1995; Snow *et al.*, 1996).

Table 6. Behaviour of dissociated rat central neurons on ECM-proteins. The effect of each substrate is illustrated as a schematic drawing.

Protein	Neurons	Effect	Stimul.	Inhibition
Laminin-1	Cerebellum HC, cortex SC, retina		+++	N.D.
Laminin-2 (Merosin)	Retina Cerebellum (P1-P3)		+	N.D.
Laminin-3 (s-laminin)	SC CG		+	-
γ 1 chain neurite outgrowth domain	Cerebellum HC, Retina SC		++	N.D.
Netrin-1	SC, Retina		+++	-
Thrombospondin	Cerebellum Retina		+++	N.D
Fibronectin	Cerebellum Retina		+	N.D
Tenascin-C/R	HC; Retina Cerebellum		++	-
Type IV Collagen	Cerebellum Cortex		N.D	-
HSPG	Cerebellum Retina SC		++	-
CSPG	Cerebellum SC		++	--
KSPG	Cerebellum SC		++	--

2.3.2. Netrins, semaphorins and ephrins in axon guidance

Netrins are soluble axon outgrowth promoting molecules, homologous to the $\gamma 1$ chain of laminin-1 and *Unc-6* (Hedgecock *et al.*, 1990; Ishii *et al.*, 1992). Netrins are found to either attract or repel different subsets of axons towards or away from the ventral midline (Hamelin *et al.*, 1993; Serafini *et al.*, 1994, 1996; Colamarino and Tessier-Lavigne, 1995). The receptors that mediate the attractive effect of netrins are DCC and neogenin (for ref. see Chisholm and Tessier-Lavigne, 1999; Livesey, 1999). The repellent effect of netrins is mediated by receptors of the UNC-5 family (for ref. see Chisholm and Tessier-Lavigne, 1999; Livesey, 1999).

Semaphorins are a large family of secreted and membrane bound molecules (Semaphorin Nomenclature Committee, 1999) that were first identified as repellents for axonal growth (Mark *et al.*, 1997; Mueller, 1999). Receptors for secreted semaphorins, neuropilins, are necessary for repulsive guidance cues (Kitsukawa *et al.*, 1997; Giger *et al.*, 2000). Recently semaphorins (Sema-1a) were also shown to have attractive cues for developing axons (Wong *et al.*, 1999).

Ephrins and their receptors, the Eph receptor tyrosine kinases, mediate axonal patterning and pathway selection through contact dependent repulsion (Flanagan and Vanderhaeghen, 1998). Eph receptors are named ephA or ephB depending on the class of ligands they bind (Eph Nomenclature Committee, 1997). Ephrins are divided into two classes: EphrinA binds to EphA and is GPI linked to membranes, while ephrinB is a transmembrane protein that binds EphB. Further evidence for the role Eph receptors in axon guidance has been obtained from targeted mutation studies. Embryos lacking the EphB2 receptor function do not have a part of the anterior commissure (Henkemeyer *et al.*, 1996). Ephrins and Eph receptors are required for establishing appropriate connections in the retinotectal systems (Cheng *et al.*, 1995; Drescher *et al.*, 1995; Monschau *et al.*, 1997; Friesen *et al.*, 1998).

2.3.3. Cell adhesion molecules

Pioneer axons may serve as guidance cues for the axons following them. When a large number of axons follow each other they become grouped together, which is called fasciculation. Cell adhesion molecules function in the fasciculation process. In *Drosophila* Fasculin II (Harrelson and Goodman, 1988), a homologue of the vertebrate NCAM, is involved in fasciculation (Grenningloh *et al.*, 1991; Lin *et al.*, 1994). L1, NCAM-180, and

N-cadherin promote neurite outgrowth *in vitro* (Doherty *et al.*, 1989; Lemmon *et al.*, 1989; Bixby and Zhang, 1990) and *in vivo* (Tomasiewicz *et al.*, 1993; Dahme *et al.*, 1997) but they act on different steps of neurite outgrowth (Takei *et al.*, 1999). L1 functions in neurite extension and NCAM-180 in growth cone protrusion (Takei *et al.*, 1999). NrCAM and axonin-1 mediate commissural axons to cross the midline. Reagents that disturb the interactions of growth cone associated axonin-1 with NrCAM in floor plate result in pathfinding errors *in vivo* (Stoeckli and Landmesser, 1995).

2.4. Neuronal degeneration

Neuronal degeneration is a normal phenomenon during nervous system development (Hutchins and Barger, 1998). Virtually all cell populations in the vertebrate nervous system undergo massive cell death during the early stages of their development. Approximately half of all the neurons produced during neurogenesis die (Raff *et al.*, 1993). Several reasons may cause neuronal degeneration during development. E.g. the failure of neurons to obtain support from some critical neurotrophic factors (O'Leary and Cowan, 1984; Pilar *et al.*, 1980) or their inability to make appropriate axonal contacts (Oppenheim, 1990) might lead to their death. Sex hormones also play a role in neuronal degeneration during development (Dibner and Black, 1976; Wright and Smolen, 1985). Developmentally occurring cell death is suggested to occur via apoptosis, characterized by shrinkage of the cell and its nucleus, condensation of chromatin, and production of membrane-enclosed particles containing intracellular material known as "apoptotic bodies" (Kerr *et al.*, 1972; Wyllie *et al.*, 1980; Arends and Wyllie, 1991).

In the adult brain, ischemic brain injury that leads to intracellular overdose of Ca^{2+} may lead to neuronal death (Choi, 1988a). Glutamate excitotoxicity is thought to participate in pathogenesis of the ischemic brain injury (Choi, 1988b; Choi and Rothman, 1990). Glutamate-induced neuronal death may occur either via apoptosis or necrosis depending on the mitochondrial function (Ankarcrona *et al.*, 1995). Necrosis is associated with an extreme energy failure of the mitochondria (Ankarcrona *et al.*, 1995) whereas relatively intact mitochondrial activity appears to be necessary for the apoptosis to proceed (Ankarcrona *et al.*, 1995). Thus, necrosis is a passive process characterized by cell and organelle swelling and lysis of the intracellular contents of the cell into the extracellular environment. Neurodegenerative diseases, such as Alzheimer's disease, Huntington's

disease, and amyotrophic lateral sclerosis (ALS) all cause neuronal death in the adult nervous system. In these diseases neuronal death is shown to occur via apoptosis (Smale *et al.*, 1995; Anderson *et al.*, 1996; Zhang *et al.*, 1997 for AD; Dragunow *et al.*, 1995; Herdreen *et al.*, 1994 for Huntington's disease; Alexianu *et al.*, 1994; Rothstein *et al.*, 1994 for ALS).

2.4.1. Alzheimer's disease

Alzheimer's disease (AD) is a neurodegenerative disorder that leads to early impairment of memory and other intellectual functions in humans. The most important neuropathological signs of Alzheimer's disease include the accumulation of senile plaques and neurofibrillary tangles in the brain tissue combined with extensive neuronal loss in hippocampus (HC), neocortex, and other areas of the brain. Senile plaques are extracellular deposits with an amyloid core surrounded by degenerating neurons, sprouting neurites, reactive astrocytes, and activated microglia. The major constituent of the plaques is the amyloid β protein peptide ($A\beta$ 1-42; Masters *et al.*, 1985; Selkoe *et al.*, 1986). Additional proteins found in the plaques include α -antichymotrypsin (ACT; Abraham *et al.*, 1988), apolipoprotein E and J (Namba *et al.*, 1991; Wisniewski and Frangione, 1992; Choi-Miura *et al.*, 1992), IgG (Eikelenboom and Stam, 1982), several complement proteins (Kalaria and Perry, 1993), glycosaminoglycans (Snow and Wight, 1989), fibronectin (Howard and Pilkington, 1990), and laminin-1 (I).

2.4.1.1. Pathophysiology of Alzheimer's disease

Alzheimer's disease appears in sporadic and familial forms (familial AD = FAD). Certain familial forms of AD have been linked to chromosome 21 (St George-Hyslop, *et al.*, 1987). The amyloid β protein gene maps to this chromosome (Kang *et al.*, 1987, Tanzi *et al.*, 1987). In all cases of AD there is excessive deposition of $A\beta$ (Masters *et al.*, 1985; Selkoe *et al.*, 1986), an insoluble, 42 residue proteolytic processing product of the amyloid precursor protein (APP). In Alzheimer's disease APP, normally located on cell membranes (Kang *et al.*, 1987), is proteolytically cleaved and released into the extracellular space. All mutants of β APP that have been linked to AD lead to altered proteolytic processing of this protein. This altered proteolytic processing favors the production of amyloidogenic and neurotoxic $A\beta$ fragment of β APP (Selkoe, 1995). An eleven amino-acid-long peptide (amino acids 25-35) derived from the 42-amino-acid amyloid β protein peptide has been shown to promote neurite outgrowth at low concentrations and to be neurotoxic at high

concentrations (Whitson *et al.*, 1989; Yankner *et al.*, 1989, 1990a,b; Rober *et al.*, 1991). This suggests a role for this peptide both in neuronal degeneration and sprouting events detected in the plaques. Neuronal death induced by amyloid β -peptide *in vitro* occurs by an apoptotic way (Loo *et al.*, 1993; Watt *et al.*, 1994).

In addition to its linkage to chromosome 21 AD is also genetically linked to chromosomes 1, 14, and 19. Mutations of the Presenilin-1 (PS-1) gene in chromosome 14 (Alzheimer's Disease Collaborative Group, 1995; Pereztur *et al.*, 1995; Sherrington *et al.*, 1995) and of the Presenilin-2 (PS-2) gene in chromosome 1 (Levy-Lahad *et al.*, 1995; Rogaev *et al.*, 1995) are found to cause the early onset familial form of Alzheimer's disease (FAD). These PS-1 and PS-2 genes encode for two highly homologous proteins (Levy-Lahad *et al.*, 1995; Rogaev *et al.*, 1995; Sherrington *et al.*, 1995) that have been localized either intracellularly within the endoplasmic reticulum and the Golgi complex (Kovacs *et al.*, 1996) or alternatively in the nuclear membrane and in association with interphase kinetochores and centrosomes (Li *et al.*, 1997). Recently, PS-1 has also been localized to the cell surface (Schwarzman, *et al.*, 1999). In addition to the PS-1 gene a gene for α -antichymotrypsin (ACT) also localizes in chromosome 14 (Billingsley *et al.*, 1993). ACT is a protease inhibitor found in the amyloid deposits of AD and in preplaques (Abraham *et al.*, 1988). ACT has been proposed to accelerate the fibril-formation of A β (Ma *et al.*, 1994). In some families with the late onset form of AD a linkage to chromosome 19 has been demonstrated (Pericak-Vance *et al.* 1991). This is relevant for the role of apolipoprotein E in AD, because apolipoprotein E gene localizes in chromosome 19 (Lin-Lee *et al.*, 1985). Apolipoprotein E4 is a known risk factor for Alzheimer's disease (Corder *et al.*, 1993) although its mechanism of action is presently unknown (Mahley and Huang, 1999).

In addition to linkage studies several results indicate that defects in energy metabolism may contribute to the pathogenesis of AD (Blass and Gibson, 1991). In the late-onset AD mitochondrial cytochrome c oxidase (CO) activity in brains and platelets of the affected individuals is decreased (Parker *et al.*, 1990; Mutisya *et al.*, 1994; Chagnon *et al.*, 1995). Recently, mutations of the mitochondrial CO gene that associate with the late-onset AD have been found to result in overproduction of reactive oxygen species *in vitro* (Davis *et al.*, 1997). The free radicals formed may participate in the cleavage of β -amyloid protein and result in the aggregation of A β peptide in the AD brain tissue.

2.4.1.2. Extracellular matrix molecules in Alzheimer's disease

Several extracellular matrix molecules have been localized in the AD brain. Fibronectin is found in plaques and in crystal-like formations in the gray matter (Howard and Pilkington, 1990). Laminin-1 (I) is localized to the plaques of AD. In cerebral microvessels collagen IV content is increased in AD (Kalaria and Pax, 1995). At least four different classes of proteoglycans (PGs) have been detected in the AD brain. These include heparan sulfate PGs (Snow *et al.*, 1988; 1990; Snow and Wight, 1989), dermatan sulfate PGs (Snow *et al.*, 1992), chondroitin sulfate PGs (DeWitt *et al.*, 1993), and keratan sulfate PGs (Snow *et al.*, 1996).

APP binds to several ECM proteins, including HSPGs (Narindrasorasak *et al.* 1991; Snow *et al.*, 1994) and laminin-1 (Narindrasorasak *et al.* 1992; Kibbey *et al.* 1993). In fact, APP may be a laminin receptor that binds specifically to the IKVAV-sequence of the $\alpha 1$ -chain of laminin-1 (Kibbey *et al.* 1993). Biological functions of A β -peptide could be regulated by its binding to ECM molecules (Koo *et al.*, 1993). E.g, low concentrations of substrate-bound A β was found to promote neurite outgrowth when combined with low amounts of the ECM proteins laminin or fibronectin (Koo *et al.*, 1993).

Recently, ECM molecules, including laminin-1, have been shown to influence the biogenesis of APP and the generation of amyloidogenic fragments containing amyloid β -peptide (Bronfman *et al.* 1996ab, Mönning *et al.* 1995). Perlecan, a HSPG, accelerates A β fibril formation and stabilizes these fibrils (Castillo *et al.*, 1997). Importantly, laminin-1 has been shown to reduce the fibril formation of A β -peptide *in vitro* (Bronfman *et al.*, 1996a, 1996b, 1998; Monji *et al.*, 1998a, 1998b; Drouet *et al.*, 1999). In the hippocampus and the frontal cortex of Alzheimer's disease patients the expression of laminin-1 is increased compared to normal control brains (I). In AD laminin-1 was identified as large extracellular punctate deposits in all senile plaques (I).

2.5. Nervous system regeneration

Regeneration of the adult mammalian peripheral and central nervous system is limited. Injuries to the central nervous system of adult mammals regenerate mainly in the olfactory bulb (Graziadei and Monti Graziadei, 1978). The peripheral nervous system is capable of regeneration, although regeneration of peripheral nerve injuries in adult mammals occurs slowly and does not necessarily lead into functional recovery.

2.5.1. Regeneration in the peripheral and central nervous systems

Axotomy or crushing of a peripheral nerve leads to Wallerian degeneration, where the portion of the axon distal to the injury site degenerates. During Wallerian degeneration the environment in the trauma site is created for regeneration. This includes chromatolysis of the cell body and removal of the distal end of the axon with myelin-derived debris by macrophages. If the regeneration is successful the changes in the cell body are reversible and the proximal stump of the axon grows within the connective tissue remaining in the distal stump and regenerate. Schwann cells in the distal stump secrete chemotrophic factors that attract growing axons by mechanisms related to axon growth during development. (Sunderland, 1991).

Axons of adult CNS do not regenerate, whereas axons of adult PNS regenerate. Lack of regeneration in the mammalian CNS is thought to be due to the yet-unidentified factors that prevent regeneration in the CNS but not in the PNS. This has been shown by studies in which the CNS axons were shown to regenerate if they grow into grafts of the peripheral nerves (Benfey and Aguayo 1982, Villegas-Perez *et al.*, 1988).

In recent years some of the CNS factors inhibitory for CNS regeneration have been identified. Cultured CNS oligodendrocytes and CNS myelin were found to inhibit neurite outgrowth of cultured neurons (Schwab and Caroni, 1988). Some of this inhibition was mediated by two CNS myelin proteins (absent from PNS myelin), NI-35 and NI-250 (Caroni and Schwab, 1988a). The use of antibodies IN-1 and IN-2 against the NI-35 and NI-250 molecules did neutralize the inhibitory effects of these antigens (Caroni and Schwab, 1988b). Transplantation of myeloma cells producing the IN-1 antibody into the dorsal fronto-parietal cortex of rats 7-10 days before a spinal cord lesion allowed CNS axons to regenerate (Schnell and Schwab, 1990). Recently identifications of *Nogo* gene in

humans and rats, that encodes of an inhibitory myelin protein have been reported (Chen *et al.*, 2000; Prinjha *et al.*, 2000; GrandePré *et al.*, 2000). The Nogo-A isoform may be the NI-250 protein recognized by the IN-1 antibody (Chen *et al.*, 2000; Prinjha *et al.*, 2000; GrandePré *et al.*, 2000). Myelin-associated glycoprotein (MAG; Trapp, 1990), another myelin protein that inhibits neurite outgrowth (Mukhopadhyay *et al.*, 1994; McKerracher *et al.*, 1994), is present in myelin sheaths of both CNS and PNS axons (Figuelewicz *et al.*, 1981; Quarles and Trapp, 1984). The inhibitory effect of MAG in the PNS is neutralized by laminin-1 (David *et al.*, 1995). Recent results indicate that adult CNS myelin does not always inhibit regeneration but the reactive glial extracellular matrix at the lesion site may be a major factor associated with the failure of axon regrowth (Davies *et al.*, 1997). Furthermore, several recent studies indicate that the failure of the axonal regrowth in the CNS but not in the PNS may be due to the inability of damaged CNS tissue to activate an appropriate inflammatory response after damage (Lazarov-Spiegler *et al.*, 1996, 1998; Rabchevsky and Streit, 1997; Rapalino *et al.*, 1998; Zeev-Brann *et al.*, 1998).

Embryonic CNS tissue regenerates (Kromer *et al.*, 1981; Björklund and Stenevi, 1984), although the mammalian brain has a critical age of development after which regeneration of the CNS does not occur (Smith *et al.*, 1986). This critical period coincides with myelination and oligodendrocyte development (Smith *et al.*, 1986; Varga *et al.*, 1995). Although CNS neurons are generally capable of regeneration (Benfey and Aguayo, 1982) and environmental factors are the major reason for their failure to regenerate, it has also been proposed that the intrinsic properties of CNS neurons may suppress regeneration. Axonal regeneration in the postnatal rat *in vitro* has been shown to depend on the maturation of axons (Li *et al.*, 1995) and injured axons of the adult rat have failed to regenerate, even in a permissive glial pathway (Davies, *et al.*, 1996). The regenerative failure of most older axons may be controlled by genetic programming absent in the developing neurons (Chen *et al.*, 1995, 1997a; Aigner *et al.*, 1995). A candidate gene that may regulate axon growth is the proto-oncogene *bcl-2* (Holm and Isacson, 1999), one of the primary factors responsible for keeping cells from choosing the apoptotic pathway. At the onset of a regenerative failure *in vitro* the expression of *bcl-2* is decreased (Chen *et al.*, 1997a). Retinal ganglion cells derived from transgenic mice over-expressing the *bcl-2* gene retain their ability to grow axons throughout their lifespan (Chen *et al.*, 1997a).

2.5.2. Attempts to regenerate PNS injuries

A standard method to rejoin transected peripheral nerves is neurorrhaphy using microscopic instruments and thin microfilament threads (Sunderland, 1991; Terzis and Smith, 1991). This methodology requires a specialist and is time consuming. Both gluing (Narakas, 1988) and laser technology (Almqvist, 1988) have been applied, but these methods are not easier and do not improve regeneration (Vastamäki, 1990). Thus, novel methods to regenerate injuries of the PNS are needed.

Tubulation of both the proximal and distal nerve ends is believed to aid regeneration. Tubulation prevents the displacement of nerve stumps, orient axon growth and vascularization, concentrates growth and trophic factors and protects regenerating tissue from scar invasion (Madison *et al.*, 1992). Various kinds of tubular prostheses have been used to bridge nerve gaps. Silicone tubes alone were found to increase regeneration compared to the untreated controls (Lundborg *et al.*, 1982). The silicone tube facilitates elongation of the axons across 10-mm gaps in rats (Lundborg *et al.*, 1982). Extended gaps could also be successfully bridged with silicone tubes filled with collagen-glycosaminoglycan (GAG; Chamberlain *et al.*, 1998) or artificial nerve grafts composed of polyamide filaments (Lundborg *et al.*, 1997). These grafts facilitated regeneration across a 15-mm gap in the rat sciatic nerve (Lundborg *et al.*, 1997; Chamberlain *et al.*, 1998). However, the use of silicone tubes necessitates surgery to suture the nerve ends into the silicone tubes. Another more serious disadvantage in the use of silicone tube grafts is the fact that axons regenerated through the silicone tubes degenerate after one year of implantation unless the silicone tube is removed. Axonal degeneration may be caused by the tube material itself or by the constriction of the proximal and distal nerve stumps due to long-term tubulization (Le Beau *et al.*, 1988). Thus, usage of silicone tubes necessitates a second surgery to remove the silicone tubes after regeneration has taken place. To avoid the second surgery biodegradable tubes have been developed. The biodegradable tubes tested include polylactate tubes (Da Silva *et al.*, 1985; Madison *et al.*, 1987), and poly(organo)phosphazene tubes (Langone *et al.*, 1995). Collagen or collagen with a GAG substrate (Collin and Donoff, 1984; Ellis and Yannas, 1996; Chamberlain *et al.*, 1998) have also been tried in addition to collagen tubes filled with Schwann cells (Kim *et al.*, 1994). Thin cellulose sheets have shown to be as effective as microsurgical neurorrhaphy in reconnecting the transected nerve (III).

2.5.2.1. Laminin treatment of the PNS injuries

Laminin-1 is a potent neurite outgrowth promoting molecule that is also involved in axon guidance (Baron van Evercooren *et al.*, 1982; Liesi *et al.*, 1984, 1989; Cohen *et al.*, 1987; Letourneau *et al.*, 1988). Astrocytes in the CNS of the lower vertebrate (Liesi 1985; Hopkins *et al.*, 1985) and Schwann cells of the mammalian PNS constitutively express laminin-1 (Cornbrooks *et al.*, 1983). Astrocytes of the olfactory bulb, the only regenerating area of the adult mammalian CNS, continuously express laminin-1 (Liesi, 1985b). These observations indicate that expression of laminin-1 correlates with nerve regeneration. This has suggested many researchers to use laminin-1 grafts in the treatment of PNS injuries. Laminin has been applied in both silicon and biodegradable tubes as well as in gels with additional ECM components or in collagen tubes also containing fibronectin (Madison *et al.*, 1985; Madison *et al.*, 1987; Bailey *et al.*, 1993; Tong *et al.*, 1994; Labrador *et al.*, 1998). The functional role of laminin-1 in neuronal regeneration is further emphasized by the fact that the inhibitory effect of a myelin-associated glycoprotein (MAG) on neurite outgrowth is neutralized by laminin-1 (David *et al.*, 1995). Microglial grafts that supposedly counteract the mechanisms that inhibit CNS regeneration induce laminin expression at the injury site (Zak, 1987; Rabchevsky and Streit, 1997).

The Schwann cell laminin differs from the basement membrane form of the protein in that it lacks the conventional $\alpha 1$ -chain (Davis 1985; Edgar *et al.*, 1988). It is possible that the α -chain of Schwann cell laminin is either missing or that it is shorter than the $\alpha 1$ -chain of laminin-1 (e.g. $\alpha 4$ -chain). It appears that no α -chain is necessary for the regeneration of peripheral nerves. This is evident from a study in which a decapeptide from the neurite outgrowth domain of the $\gamma 1$ -chain was coupled with a type I collagen matrix to form a filter. This peptide graft supported nerve regeneration as well as suturation of the rat sciatic nerve (Kauppila *et al.* 1993). Animals that received laminin grafts to repair their sciatic nerves showed significantly lower autotomy scores (numbers of self-mutilated phalanx) compared to the animals with conventional neurorrhaphy. The thickness of the graft, however, formed a limitation for its use. If a large number of adjacent nerve fascicles were to be separately reconstructed, the graft resulted in a compression damage of the repaired nerves (Kauppila *et al.* 1993). A thick graft could also impair regeneration by preventing the diffusion of factors critical for regeneration.

2.5.3. Attempts to regenerate CNS injuries

The first transplantation studies of nervous tissue were performed as early as in 1863 by using a portion of the optic nerve to regenerate a dog hypoglossal nerve (For references see Dellon and Dellon, 1993). Various types of mature CNS neurons have expressed regenerative capacity of their injured axons when the axons are in contact with peripheral nerve grafts (David and Aguayo, 1981; Munz *et al.*, 1985; Morrow *et al.*, 1993; Decherchi and Gauthier, 1996) and a 20% recovery of the dorsal root fibers into the spinal cord was obtained using living glial cell implants (Kliot, *et al.*, 1990). Cultured Schwann cells were found to promote regeneration of CNS neurons when transplanted into the lesioned adult rat spinal cord (Paino and Bunge, 1991; Li and Raisman, 1994). Recently macrophages, prestimulated with spontaneously regenerating peripheral nerve tissue, have been shown to stimulate optic nerve regeneration in the adult rat (Lazarov-Spiegler *et al.*, 1996). These activated macrophages were also found to enhance the clearance of damaged myelin from the lesion site (Lazarov-Spiegler *et al.*, 1998). Grafts of cultured microglial cells were also shown to induce prominent neurite outgrowth of the lesioned spinal cord of adult rats (Rabchevsky and Streit, 1997).

No therapy for the repair of spinal-cord injuries is presently available (Olson, 1997). Recently, promising results regarding both the histological and functional recovery of CNS injuries in rats has been achieved by chronic infusions of IN-1 antibody, neurotrophin-3 or both (Diener and Bregman, 1994; Schnell *et al.*, 1994; Bregman *et al.*, 1995). The first evidence for true functional regeneration in the adult mammalian spinal cord was reported when the cut rat spinal cord was bridged by peripheral nerves in-between the individual spinal axon tracts and areas of neuronal cell bodies (Cheng *et al.*, 1996). The bridge was further stabilized by FGF-impregnated fibrin glue (Cheng *et al.*, 1996).

Neural transplantation of dopaminergic neurons of human embryos into the brains of Parkinson's disease patients offers a novel therapeutic tool to treat this neurodegenerative disease (Lindvall *et al.*, 1994; Kopyov *et al.*, 1996). Human CNS progenitor cells, maintained in a proliferative state in culture, have been shown to migrate and differentiate into both neurons and astrocytes following intracerebral grafting in rats (Svendsen *et al.*, 1997). Although a low number of tyrosinase hydroxylase-positive neurons (approximately 200) were identified *in vitro*, they could allow functional recovery, because allograft

studies have shown that only 120 dopamine cells are needed to produce functional recovery effects (Brundin *et al.*, 1985). Transplanted pig fetal neural cells have also been grafted in brains of Parkinson's disease patients (Deacon *et al.*, 1997). The implanted pig dopaminergic neurons matured and extended neurites into the human host brain (Deacon *et al.*, 1997). Continuous neurogenesis is shown to occur in the adult mammalian brain, which may allow future development of methods that favour regeneration in the CNS.

AIM OF THE STUDY

Aim of this study was

- to investigate the possible involvement of the neurite outgrowth domain of the $\gamma 1$ -chain of laminin-1 in pathophysiology of human neurodegenerative disorders (I)
- to use the weaver mutant mouse as a model system to investigate the possible role of laminin-1 $\gamma 1$ -chain in neuronal migration defects and in neuronal death (II)
- to develop novel tools for neurography (III)
- to use the F9 cells to develop a novel model system for future studies on neuronal differentiation (IV)

MATERIALS AND METHODS

I

Production of Antisera to Laminin Peptides

Synthetic peptides derived from the C-terminal part of the mouse γ -1-chain (Sasaki and Yamada, 1987) or α 1-chain (Tashiro *et al.*, 1989) of laminin-1 were from Multiple Peptide Systems (La Jolla, CA). Rabbits for anti-1533 a and anti-1533 b were immunized with a 30 amino acid peptide (p32; I Fig. 1) and rabbits for anti-1543 were immunized with a 10 amino acid peptide (p20; I Fig. 1) conjugated to itself to form a larger polymer (Linder and Robey, 1987). Rabbits for anti-2091 were immunized with a 22 amino acid peptide from the α -chain of laminin-1 (PA-22; I Fig. 1.). The peptide columns used to absorb each antibody were done by coupling the peptides to Sepharose 4B (Pharmacia, Uppsala, Sweden) according to the instructions of the manufacturers.

Characterization of Anti-peptide Antisera

The peptides p20 and p32 were labeled with ^{125}I using the Bolton-Hunter reagent and the chloramide T method (Risteli and Risteli, 1987), respectively. The radioimmunoassay (Timpl and Risteli, 1982) and inhibition (Risteli and Risteli, 1987) assays were carried out as described previously. Mouse laminin-1 was purified as described earlier (Timpl *et al.*, 1979). Ten micrograms of purified laminin-1 was run in 5% sodium dodecyl sulfate (SDS) gel electrophoresis under reducing conditions and blotted onto nitrocellulose filter as described before (Towbin *et al.*, 1979). The transferred proteins were visualized with 0.2% Ponceau (Sigma, St Louis, MO) in 4% trichloroacetic acid and processed for immunostaining using anti-peptide antibodies or antibodies to native laminin-1 as described earlier (Liesi and Risteli, 1989). 100 ng of native laminin-1 or 10 ng of each synthetic peptide were dot blotted onto a nitrocellulose filter. The filter was dried and processed as described earlier (Liesi and Risteli, 1989).

Immunostaining of Brain Tissue Sections

Brain tissue samples from Alzheimer's disease and Down's syndrome patients and from normal, age-matched controls were obtained from the National Neurological Research Bank (VAMC Wadsworth Division, Los Angeles, CA). All samples were neuropathologically verified and were from either the hippocampus or the frontal cortex of the diseased brains. Brain samples from a total of four different Alzheimer's disease patients and five normal, age-matched controls were analyzed. In addition autopsy material from the hippocampus of a 44-year-old Finnish Down's syndrome patient was

obtained. The deep frozen brain tissue was cut into 10 μm cryostat sections and dried for 2 hours at room temperature. The sections were fixed with either 2% paraformaldehyde or 0.4% p-benzoquinone (Koch-Lights Inc.) in phosphate-buffered saline (PBS) for 15 min at 4°C and processed for immunocytochemistry for laminin or other molecules as described elsewhere (Liesi and Silver 1988). In short, the sections were dehydrated in graded series of alcohols (50%-70%-96%-100%-xylene) for 5 min and rehydrated through the same series to PBS. After rinse in PBS for about 30 min the sections were incubated overnight at 4°C in normal sheep serum (Sigma, St. Louis, MO) and incubated with the primary antisera for 24 hours at 4°C.

The antiserum against mouse laminin-1 (Liesi and Silver 1988) was used at a dilution of 1:2000 and affinity-purified antibodies against the P1 fragment of human laminin-1 (Risteli and Timpl, 1981) at a concentration of 10 $\mu\text{g}/\text{ml}$. These two antibodies were applied as an additional control for the specificity of the staining patterns observed. Antibodies against the synthetic peptides derived from laminin-1 (I) were applied at 1:1000- 1:2000 dilutions. The sections were incubated with the primary antibodies for 24h, briefly washed in PBS, and exposed to sheep antirabbit immunoglobulins coupled to fluorescein isothiocyanate (FITC; Wellcome, Beckenham, UK). In double staining for α -antichymotrypsin or amyloid β protein the sections were further incubated with normal rabbit serum followed by incubation with rabbit antibodies to α -antichymotrypsin (Dakopatts, Copenhagen, Denmark) or rabbit antibodies to the amyloid β protein peptide (Masters *et al.* 1985). The antibodies were diluted 1:500 and 1:1000, respectively. The specificity of this sequential immunostaining procedure for two rabbit antibodies has been confirmed (Liesi *et al.* 1983b). After a brief wash in PBS, the sections were mounted in PBS:glycerol (1:1) and viewed with an Olympus BH2 microscope using epifluorescence and appropriate filter combinations.

Demonstration of mRNA in Brain Tissue

The brain tissue to be analyzed was frozen and ground in liquid nitrogen, solubilized in guanidine isothiocyanate, and layered on a CsCl cushion (Chirgwin *et al.*, 1979). The separated RNA was electrophoresed in formaldehyde agarose gels (Goldberg, 1980) and visualized by ethidium bromide shadowing. After transfer, the RNA was hybridized with oligolabelled cDNA probes to the $\alpha 1$ (Sasaki *et al.*, 1988), $\beta 1$ (Sasaki *et al.*, 1987) and $\gamma 1$ (Sasaki and Yamada, 1987) chains of laminin-1. Shortly, the cDNA probes were

oligolabelled using Random Primed DNA Labelling Kit (Boehringer Mannheim) and purified in G50 column. The filters were incubated in prehybridization solution (50% deionized formamide, 50mM Hepes, 3×SSC, 5×Denharts, 0.2 mg/ml SS-DNA, 1 mM EDTA, 0.2% SDS, 0.12%NaCl) for one hour at +37°C. After prehybridization SS-DNA (6.4 mg) and oligolabelled cDNA probe was added to the prehybridization solution and filters were further incubated overnight at +37°C. After hybridization filters were washed in 2×SSC with 0.1% SDS at+37°C 15 min and exposed to X-ray film. A mouse α -actin cDNA probe (Minty *et al.*, 1982) served as a control for the quantity of mRNA loaded. The same Northern blot lane was hybridized with each of the four probes in a sequential order. To ensure that washing off of the probe at 60°C in 50% deionized formamide/0.1×SSC/0.1% SDS for 30 min did not wash off the mRNA, the α -chain message was detected first, the γ -chain message second followed by the β -chain and α -actin transcripts.

Immunoblotting of Laminin-1 in Brain Tissue

The frozen brain tissue was transferred into a sterile 50 ml Falcon tube containing 10 ml of lysis buffer (2% Triton-X-100, 0.5 M NaCl, 10 mM Tris HCL, pH 7.4, 1mM PMSF). After homogenization the tissue was dissolved in the lysis buffer and incubated for 30 min at 4°C. The undissolved material was centrifuged at 14 000g for 20 min at 4°C. The supernatant was transferred to a new tube and dissolved in 5 ml Laemmli sample buffer with 0.1 M β -mercaptoethanol. The proteins were analyzed via 5% SDS-gel electrophoresis. After transfer to nitrocellulose (Towbin *et al.*, 1979) the filter was stained with 0.2% Ponceau S in 4% trichloroacetic acid and processed for immunoblotting using antibodies to native laminin-1 as described elsewhere (Liesi and Risteli, 1989).

Polymerase Chain Reaction (PCR) Analysis of the Laminin β Chain mRNA in Brain Tissue

A specific DNA copy of the laminin β 1 chain RNA sequence was synthesized using two oligonucleotide primers (B1 and B2) derived from the published sequence of human laminin β 1 chain gene (Vuolteenaho *et al.*, 1990). The primer set spanned a region of 287 bp, extending from exon 9 to exon 10. First-strand cDNA synthesis was accomplished by extension with the downstream primer B2 (5'-GGATCTCGGATGTCCCTCTCTGG-3') in a 20 μ l reaction volume containing 1 μ g

total RNA, 30 pmol PCR primer, and reverse transcriptase from AMV (Promega) under conditions suggested by the supplier. Fifty percent of the heat-treated first-strand cDNA mixture was then included in 50 μ l volume with 1 μ M of each of the primers B1 (5'-CATGTGCAGGCATAACACCAAGG-3') and B2, 0.2mM each of dNTPs, 10 mM Tris-HCL (pH 8.4), 50 mM KCL, 1.5 mM MgCl₂, 0.1mg/ml gelatin, and 1.5 U AmpliTag DNA polymerase (Perkin-Elmer Cetus, Norwalk, CT). Forty-five cycles of PCR amplification were performed at 95°C, 55°C and 72°C for 1 min each in a DNA thermal cycler (Techne PHC-2) according to the method of Saiki *et al.* (1985). The double-stranded PCR products resulting from this amplification were fractionated by electrophoresis on a 2% agarose gel followed by elution, extraction with equal volumes of phenol and chloroform, and precipitation with ethanol. One-half of this template was subjected to DNA sequence analysis using the dideoxy chain termination method of Sanger (1977) with Sequenase (U.S. Biochemicals). The sequence was determined in both directions using the PCR primers as sequencing primers. The sequencing reactions were run on 5% polyacrylamide gels containing 7 M urea and were visualized by autoradiography (Konica X-ray film).

II

Weaver Mice

Heterozygous (+/wv) mice carrying the weaver mutation were obtained from the Jackson Laboratories (Bar Harbor, ME) and bred at the colony of the Institute for Basic Research, New York or at the Department of Anatomy, University of Helsinki. The mice were free of mycoplasma, MHV, Sendai, and other common mouse pathogens. They were bred on a B6CBA-A^{w-J}/A wv genetic background and homozygous weaver (wv/wv) or control (+/+) mice were used for experiments 7 to 13 days after birth.

Neuronal Cultures

Cultures of the granule neurons were initiated from cerebella of 7- to 10-day-old homozygous (wv/wv) weaver mice and their normal (+/+) litter mates. The cerebella were aseptically removed and the cells dissociated using a trypsin/DNase-treatment as described in detail by Trenkner (1991). One hundred thousand cells were plated on laminin-coated, 22-mm glass coverslips and cultured for 24 hr in serum-free RPMI 1640 culture medium (Gibco, BRL) supplemented with antibiotics and 2 mM glutamine. The cultures were fixed in 2% paraformaldehyde in PBS for 15 min for quantification and

immunostaining experiments. In a subset of experiments cultures were grown in the presence of 100 U/ml of aprotinin (ICN Biomedicals, Costa Mesa, CA) added into the culture medium 1hr after plating. For biochemical studies on proteolytic enzyme activities the cells were cultured overnight, lysed, collected in Laemmli sample buffer without β -mercaptoethanol (LSB-), and run in 3-12 % gradient SDS gels. Quantification of neurite outgrowth was done in six normal cultures, six weaver cultures, and six weaver cultures containing aprotin: The numbers of neurons in six representative fields on each laminin -coated glass coverslip were evaluated using phase-contrast microscopy. The mean numbers of neurons, and of neurons bearing long neurites (>10 times cell soma), were evaluated in each case. Approximately 60 neurons were counted per coverslip. Oneway variant analysis (ANOVA) on the Instat (1.11a) program was used for statistical analysis. The statistical comparisons between each individual group of neurons were performed using the Bonferroni's modified t-test.

Immunocytochemistry

Cerebellar tissues of the normal (+/+) and homozygous (wv/wv) weaver mutant mice were frozen in powdered dry ice and cut to 10- μ m cryostat sections. The sections were dried for 2 hr at room temperature, lightly fixed in freshly prepared 0.4% *p*-benzoquinone (Sigma, St. Louis, MO) in PBS, and processed for immunocytochemistry as described in detail by Liesi and Silver (1988). Antibodies against laminin-I (Liesi and Silver, 1988) and its neurite outgrowth domain (anti-1533a and anti-1534 in I) were those used previously, and their specificities were confirmed (I). The anti-1533a was used for *in vivo* studies and the anti-1543 was used for *in vitro* studies. Rabbit antibodies against tissue plasminogen activator (tPA) were those used in earlier studies (Tienari *et al.*, 1991). Antibodies against laminin-I and its neurite outgrowth domain were applied at 1:2000 and those against tPA at 6 μ g/ml for overnight incubations at +4 C. Binding of primary antibodies was detected using sheep anti-rabbit immunoglobulins coupled to FITC (Wellcome, Beckenham, UK). In some experiments sections were immunostained for the glial fibrillary acid protein (GFAP; Bignami and Dahl, 1974a, b) or the L1 antigen using rabbit polyclonal antibodies (Rathjen and Schachner, 1984). The immunostained tissues were viewed and photographed using an Olympus BH2 fluorescence microscope with appropriate filter combinations.

Zymographic Assays of the Normal and Weaver Mouse Cerebellar Tissues in Vivo and in Vitro

Cerebellar tissue or granule neurons on a laminin substratum were collected, lysed in LSB-, and run in 3-12% gradient gels. The gels were washed free of SDS and overlaid with agarose containing casein and plasminogen as described in detail (Tienari *et al.*, 1991). Proteolytic enzyme activities of the cell extracts were monitored by incubating gels with the overlays for 12-24 hr followed by staining of the overlays with amino black to detect the sites for enzyme activity. Human recombinant urokinase (1.0 U/ml uPA; Calbiochem, San Diego, CA) and tissue plasminogen activator (5 ng/ml tPA, American Diagnostica, Greenwich, CT) were used as controls. The specificity of the enzyme reactions was monitored by control experiments in which 100 µg/ml of the neutralizing antibodies against either uPA or tPA (Tienari *et al.* 1991) was added into the agarose overlays to inhibit the functions of the respective enzymes present in the cell or tissue extracts.

Electrophysiology

Resting membrane potentials of the normal and weaver neurons on a laminin substratum were determined using a List EPC-7 patch clamp amplifier. The resting membrane potentials of the cultured cells were measured immediately after entering the whole cell patch configuration as described (Hamill *et al.* 1981). Pipettes were pulled from borosilicate glass and lightly fire polished. Patch pipettes contained (in mM): 140 CsCl, 10 BAPTA, 2 MgCl₂, and 10 HEPES (pH 7.2). Experiments were performed at room temperature in the RPMI 1640 culture medium.

III

Animals

Male Wistar rats (Hannover strain; Harlan, Netherlands) were used in all experiments. The rats were 8 months old (380-470 g) and had water and food available *ad libitum*. They were housed in groups of six animals with a light cycle 6.00-18.00 h and a relative humidity of 35-55%. All experiments were approved by the Institutional Ethics Committee of the Institute of Biomedicine, University of Helsinki and by the Provincial Government of Uusimaa, Finland.

Behavioral Testing of Neuropathic Symptoms

The mechanical withdrawal thresholds of the sciatic nerve areas of the hindpaws of twelve rats were first determined as described earlier (Mansikka and Pertovaara, 1995). Briefly, the rat was standing or walking on a metal grid and the right hindpaws were stimulated with a series of calibrated von Frey monofilaments (Stoelting, USA). The central pads of the paw served as the stimulus site. The monofilaments were applied to the foot pad in series of increasing force until the rat withdrew the limb. The lowest force producing a withdrawal response was considered the threshold. The threshold for each hindpaw was based on three separate measurements. The tests were performed double-blind 1 and 2 months after trauma. The statistical comparisons were performed using Friedman repeated measures of variance on ranks followed by two tailed Mann-Whitney U-test. $P < 0.05$ was considered significant in all statistical comparisons of this series of experiments.

Traumatization and Reconstructive Surgery

The rats (6 in each experimental group) were anesthetized with pentobarbital (50 mg/kg, Orion, Finland) and the right sciatic nerve was exposed under aseptic conditions. The nerve was transected at mid-thigh and either resutured with two perineural 10-0 monofilament sutures or reconnected by using a moistened lens cleaning paper (Illford, U.K.) which was wrapped around the stumps of the transected nerves (Kauppila *et al.* 1993). The anatomical orientation of the nerve stumps was restored by observing the fascicular and vascular anatomy of the transected nerves. After reconstruction the wounds were closed in layers with 3-0 silk sutures.

Scoring of Autotomy

Autotomy of the digits was observed once a week during a period of six months. Autotomy was scored as described earlier (Kauppila and Pertovaara, 1991): Each self-mutilated phalanx represented one score point. Two-tailed Mann-Whitney U-test was used for statistical comparisons. The tests for mechanical withdrawal thresholds were performed one month and two months after nerve resuturation because the regenerated axons reach the mid-thigh and establish their connections to the periphery during the next month (Kauppila *et al.* 1993).

Electrophysiologic Testing of Recovery

Six months after surgery the rats were anesthetized with pentobarbital (50 mg/kg) and reinnervation of the soleus and gastrocnemius muscles was studied as described earlier (Kauppila *et al.* 1993). The sciatic nerve (both operated and unoperated side) was exposed, transected proximal to the lesion, mounted on a bipolar platinum stimulating electrodes, and covered with liquid paraffin. The hindlimb was immobilized with needles and the achilles tendon was cut and connected to a strain gauge (Grass, U.S.A). The initial load was adjusted to 5 g. The sciatic nerve was stimulated proximal to the trauma site with square wave pulses of 0-01 or 0.1 ms duration and constant voltage of 15 V (stimulator Nihon-Kohden, Japan). Five consecutive stimuli were used for testing at both stimulus durations. The tension resulting in a muscle twitch was recorded after each stimulus and the differences in tension between the control and operated side were compared with the one-sided Students t-test. This test was chosen, because we wanted to detect even the smallest putative differences between the forces of the control (left) side and the trauma (right) side.

Testing of Anatomic Recovery

To evaluate the degree of anatomical recovery the rats were killed with an overdose of pentobarbital, and the muscles dissected out and weighed at the end of the electrophysiological experiments (Kauppila *et al.* 1993). The statistical analysis was carried out using the Student two-tailed t-test.

Immunocytochemistry

The nerves were dissected free, frozen on dry ice and cut in 10 µm cryostat sections. The cryostat sections were fixed in 0.4% p-benzoquinone in PBS for 15 min, washed in PBS, and dehydrated and rehydrated as described earlier (Liesi and Silver, 1998). Mouse monoclonal antibodies against the 200 kDa neurofilament protein (RT97, Boehringer, Germany) were applied at 1:1000 dilutions. After an overnight incubation with the first antibodies the sections were washed in PBS and exposed to goat anti-mouse immunoglobulins coupled to TRITC (Cappel, PA, U.S.A) for 1 h. The immunocytochemistry was viewed and photographed with an Olympus BH-2 microscope with appropriate filter combinations.

IV

Cell Cultures

The cells were those used previously (Wartiovaara *et al.*, 1978; Liesi *et al.*, 1983). The uninduced cells were maintained in plastic Petri dishes (Nunc, Denmark) in 10% fetal calf serum supplemented with RPMI 1640, antibiotics, and glutamine. For induction studies the cells were trypsinized and plated on 13 mm glass coverslips sterilized by flaming in alcohol. The cells were cultured at a density of 10^5 /coverslip in 3% fetal calf serum/RPMI overnight after which RA (Sigma, St.Louis, MO; 10^{-7}) and dibutyryl cAMP (Sigma; 10^{-3}) were included into the culture medium. The induction was completed by culturing the cells for 10 days with a media change every three days. The control cultures were plated at the same density but either maintained in 10% fetal calf serum or in 3% fetal calf serum without RA/dbcAMP and fixed after 10 days of cultivation. The cells were fixed in 2% paraformaldehyde in PBS for 15 min and immunostained immediately afterwards. For the RNA isolation studies the cells were plated at an initial cell density of 100 000 on a 9 cm plastic Petri dish either in the presence of 10% fetal calf serum or in 3% fetal calf serum with or without the inducing agents.

Cloning of a F9 Teratocarcinoma Cells

Trypsinized F9 cells were plated at low density (1×10^3) on 10cm bacteriological dishes and picked one by one to the microwells of the tissue culture plastic or to ELISA plates precoated with laminin (100 $\mu\text{g}/\mu\text{l}$). The cells were grown in 10 % fetal calf serum supplemented with RPMI 1640, antibiotics, and glutamine. Cells from the confluent microwells were transferred into the 3 cm plastic Petri dishes and further expanded onto the 10 cm plastic Petri dishes. To ensure that each clone originated from a single F9 cell the cells from 10 cm dishes were sub-cloned again in single cells as described above, regrown, and stored in 10% DMSO/FCS in liquid nitrogen. The clones were initially tested for their neuronal properties by plating them on 25 mm glass coverslips in 10% FCS supplemented with RPMI 1640, antibiotics, and glutamine. After 48 h, the cloned cells were fixed in methanol and immunostained for the neurofilament triplet proteins and other neuronal marker proteins. Only the clones derived from the laminin-coated microwells were chosen for future studies, because they expressed neuronal properties in a serum free medium (not shown), e.g. the cells had both a neuronal phenotype with neurite like extensions and expression of neurofilaments and other neuronal markers. One of such clones, the clone D9L2, was selected for further studies. The D9L2 clone was plated on 25 mm glass coverslips in a serum free RPMI supplemented with

antibiotics and glutamine. After 48 hrs the cells were fixed and processed for immunocytochemistry.

Northern Analysis

Total RNA of the uninduced (grown in 10% serum), control (grown in 3% serum), and induced F9 cells (grown in 3% serum with RA or with RA and dbcAMP) were isolated as described (Liesi and Risteli, 1989). 10 µg of each RNA was run in formaldehyde agarose gels, blotted onto nitrocellulose and hybridized with oligolabeled cDNA probes for the rat 68 kDa neurofilament protein (Julien *et al.*, 1985), and β -actin (Minty *et al.*, 1982) as described (Liesi and Risteli, 1989).

Immunocytochemistry for Neurofilament Proteins

Uninduced F9 teratocarcinoma cells grown in 10 % serum, the control cells in 3% serum, the RA/dbcAMP induced cells in 3% serum, and the D9L2 clone were fixed in 2% paraformaldehyde in PBS for 15 min, washed in PBS, and permeabilized with cold (−20°C) methanol for 5 min. The cells were immunostained using either polyclonal antibodies against neurofilament triplet proteins (Dahl, 1981) or a control antibody absorbed with neurofilaments (Dahl, 1981) as described earlier (Liesi and Risteli, 1989). Monoclonal antibodies against the 68 kDa neurofilament protein (Boehringer, Germany) and monoclonal antibodies against the 200 kDa neurofilament protein (RT97; Boehringer, Germany) were further applied at 5µg/ml concentration. Rabbit anti-NCAM antibodies (Gegelashvili *et al.*, 1993) were diluted 1:500 and mouse monoclonal antibodies against the neuron specific β -tubulin isoform (TUJI; Lee *et al.*, 1990) and were applied at 1 µg/ml concentration. The immunostained cultures were viewed with an Olympus BH2 fluorescence microscope with appropriate filter combinations.

RT-PCR Analysis

170 ng of each isolated RNA was used for one tube RT-PCR (Titan™ One Tube RT-PCR System, Boehringer Mannheim) to detect the 68 kDa neurofilament and the 200 kDa neurofilament gene transcripts. The RT-PCR was carried out for 40 min at 56 °C for RT-reaction and denaturation at 94 °C for 4 min followed by 35 cycles of denaturation at 93 °C for 1 min, annealing at 58 °C for 2 min, and elongation at 72 °C for 3 min. The primers used in RT-PCR are listed in IV, Table 1. PCR products of NF 68 kDa were isolated using a Sephaglas Bandprep kit (Pharmacia Biotech) and sequenced by the Sanger method.

RESULTS

I. LAMININ AND ITS NEURITE OUTGROWTH PROMOTING DOMAIN IN THE BRAIN IN ALZHEIMER'S DISEASE AND DOWN'S SYNDROME PATIENTS

1.1. Immunochemical characterization of antibodies against the γ 1-chain neurite outgrowth promoting peptides of laminin-1

Three different antibodies against the neurite outgrowth promoting peptide derived from the γ 1-chain (Liesi *et al.*, 1989) were produced and their specificities confirmed using dot-blots and radioimmunoassays. Anti-1543 was specific for a 10 amino acid long peptide p20 (1543-1553). Anti-1533a and b recognized a 30 amino acid long peptide p32 (1533-1563) and native and denatured form of laminin-1. Anti-1533b recognize also the p20 peptide. Specificity of anti-1533a for p32 peptide was confirmed by inhibition assays (I, Fig 2c). Inhibition assays for anti-1533b indicated specificity for both peptides.

1.2. Immunocytochemical studies on laminin and its neurite outgrowth peptide in AD brain

Antibodies against mouse EHS-tumor laminin, mainly recognizing the P1 fragment (Ott *et al.*, 1982) of laminin-1, detected laminin-1 as punctate extracellular deposits in all senile plaques in brain samples of aged male Down's syndrome patients and in brains of Alzheimer's disease patients. This was verified by double immunocytochemistry for laminin-1 and A β or α -chymotrypsin (I, Fig 4A,B). Some glial cells and their processes were also weakly immunoreactive for laminin-1. In normal control brains capillaries were the only structures immunoreactive for laminin-1. Antibodies against the human P1 fragment of laminin-1 (Risteli and Timpl, 1981) showed a similar distribution, but gave a stronger positive signal of the glial elements (I, Fig, 4C,D). Absorption of the laminin-1 antibodies by passing them through a laminin-1 column abolished immunoreactivity of the plaques and glial fibers (I, Fig. 4E,F). Using antibodies against the human laminin-1 P1 fragment laminin-1 immunoreactivity was detected around some capillaries in control brain tissue as well as in Alzheimer's disease and Down's syndrome brains.

Antibodies against synthetic peptides, derived from the C-terminal domain of the γ 1-chain (Sasaki and Yamada, 1987) of mouse laminin-1 (anti-1543 against the decapeptide; anti-1533a and anti-1533b against the 30-amino acid peptide, I), showed no punctate immunoreactivity in the plaque regions (I, Fig. 4A,B). Instead the glial cells and their fibers were immunoreactive for these peptide antibodies in the diseased (I, Fig. 4G), but not in normal control brains. This immunoreactivity was abolished by preabsorption of the antibodies through a peptide column. (I, Fig. 4H). An antibody against a neurite outgrowth domain of the α 1-chain of laminin-1 stained only the capillary basement membranes. Peptide antibodies that recognized the 10-amino-acid peptide p20 (anti-1543 and anti-1533b) showed binding of this peptide antigen as fine extracellular punctate deposits in the affected Down's syndrome brain tissue (I, Fig. 5D). The deposits of this peptide antigen were also found in the plaque areas, but there was no specific correlation with the plaques. Similar fine punctate deposits of the peptide antigen are relevant by these antisera in all tAlzheimer's brain tissue investigated. This immunoreactivity was absent in normal control brains and was abolished in tAlzheimer's disease brains, if the antibodies were preabsorbed with the corresponding peptide conjugated to Sepharose.

1.3. Demonstration of laminin-1 mRNA in Alzheimer's disease brain

Northern analysis showed a differential expression of the mRNAs for the three laminin-1 polypeptide chains in Alzheimer's disease and normal control brains (I, Fig.6). The mRNA for the α 1-chain was expressed in neither case (I, Fig.6). The 8.2 kb γ 1-chain transcript of laminin-1 was present both in control and Alzheimer's disease brains, but the expression was increased 10-fold in Alzheimer's disease brain tissue compared to the normal control brain tissue (I, Fig.6). The 5.6 kb β 1-chain transcript was detectable only in Alzheimer's disease brain tissue (I, Fig.6). As Northern blots showed no β 1-chain transcripts in control brains, a more sensitive RT-PCR was performed on normal and Alzheimer's disease brain tissues. Direct sequencing of the RT-PCR products showed that the β 1-chain transcripts were present in both Alzheimer's disease and control brain tissues.

1.4. Immunoblotting experiments on Alzheimer's disease brain tissue

Immunoblots of tissue extract of Alzheimer's disease brain tissue revealed no expression of the laminin α 1-chain, although both β 1- and γ 1-chains were seen using antibodies against the native laminin-1 molecule (I, Fig.7). In normal control brains the expression of laminin α 1-chain or β 1- and γ 1-chains was not detectable (I, Fig.6). The protein staining showed equal amounts of samples being loaded in the gels. These experiments together with Northern analysis indicated that in the Alzheimer's disease brain tissue the expression of laminin-1 is elevated compared to the normal brain tissue.

II. INCREASED PROTEOLYTIC ACTIVITY OF THE GRANULE NEURONS MAY CONTRIBUTE TO NEURONAL DEATH IN THE WEAVER MOUSE CEREBELLUM

2.1. *In vitro* studies on neuronal migration and death in the weaver cerebellum

Normal wild-type granule neurons (+/+) extended long neurites on a laminin-1 substratum (II, Table 1, Fig.1A) and deposited laminin around themselves (II, Fig.1B). The homozygous weaver (wv/wv) granule cells showed impaired neurite outgrowth on a laminin-1 substratum (II, Table 1, Fig.1B). The weaver neurons degraded laminin-1 from their substratum (II, Fig.1D). Polyclonal antibodies against the neurite outgrowth domain of the γ 1-chain of laminin-1 showed binding of this antigen along the surfaces of the weaver granule neurons (II, Fig.2A), whereas wild type granule cells did not bind this antigen (II, Fig.2B). A serine protease inhibitor, aprotinin, promoted the survival of weaver granule neurons on a laminin substratum and restored their neurite outgrowth to the level of normal granule neurons (II, Fig.4). After 12 hours in culture patch clamp studies indicated that normal neurons had resting membrane potentials (RMPs) of -61.2mV $\pm$ 2.8, whereas weaver neurons had RMPs of only -37.7 mV $\pm$ 3.4 (II, Table 2.) Aprotinin restored the RMPs of the weaver granule neurons to normal levels (-58.6mV $\pm$ 3.7). In zymographic assays normal granule neurons did not secrete detectable amounts of urokinase or tissue plasminogen activator *in vitro* (II, Fig.3A), whereas the weaver granule cell secreted tissue plasminogen activator but no urokinase (II, Fig.3B).

2.2. *In vivo* studies on neuronal migration and death in the weaver cerebellum

In the normal wild-type mouse the external granule cell layer of the cerebellum was L1-antigen positive, and the white matter of the cerebellum showed some L1-immunoreactivity (II, Fig.5A). In the weaver mouse the external granule cell layer was completely devoid of L1-antigen and only moderate L1-immunoreactivity was detected in the white matter (II, Fig.5B). Immunocytochemistry for GFAP showed that Bergmann glial fibers were highly immunoreactive for GFAP in the weaver cerebellum (II, Fig.5D), whereas the glial cells of the normal cerebellum were weakly immunoreactive for this antigen (II, Fig.5C).

Using a laminin-1 antibody that recognizes native laminin we detected an overall increase in the production of laminin-1 in the weaver cerebellum. In the normal and weaver cerebella the Purkinje cell layer showed the highest levels of laminin expression (II, Fig.6A), but in the weaver cerebellum the Purkinje cell layer and the external granule cell layer expressed higher amounts of laminin than in the normal cerebellum (II, Fig.6B). Similarly, an antibody against the γ 1-chain of laminin-1, anti-1533a, that recognizes the γ 1-chain of the native laminin-1 molecule showed an overall increase in the laminin-1 γ 1-chain expression in the weaver cerebellum (II, Fig.7B,D) compared to the normal cerebellum (II, Fig.7A,C). In the normal cerebellum the γ 1-chain antigen was detected in the Purkinje cells. In the external granule cells there was only weak immunoreactivity for this antigen. In the weaver cerebellum the immunoreactivity for the γ 1-chain was present in Purkinje cells, external granule cells, and in the molecular layer, as well as glial fibers. The weaver granule cells in the external granule cell and molecular layers showed intense immunoreactivity for the γ 1-chain antigen (II, Fig.7D).

Immunocytochemical localization of tissue plasminogen activator demonstrated a coexpression with the γ 1-chain immunoreactivity (compare II Figs. 7D and H). In the weaver cerebellum tPA showed an overall increase, but the distribution of tPA in the weaver cerebellum was similar to that in the normal cerebellum (compare II Figs. 7E,G and 7F,H). In the normal cerebellum tPA was expressed in fibers extending through the external granule cell layer, in the molecular layer, and in the Purkinje cell layer. In the weaver cerebellum the external granule cell, molecular, and Purkinje cell layers showed the strongest tPA-immunoreactivity. The upper parts of the external granule cell layer contained tPA immunoreactive glia-like fibers, and diffusely immunostained the immature

granule cells. An increase in the weaver tPA activity, consistent with the immunocytochemistry, was also verified by zymographic assays on cerebellar tissues of normal and weaver mutant mice. At P7 there was no difference in the expression of tPA between normal and weaver cerebellar tissues. However, at P13 the weaver cerebellum showed a 10-fold increase in tPA activity compared to the normal cerebellum.

III USE OF PAPER FOR TREATMENT OF A PERIPHERAL NERVE TRAUMA IN THE RAT

3.1. Immunocytochemical analysis of nerve regeneration

The 200 kDa NF expression was studied in sagittal sections of the regenerating rat sciatic nerve 6 months after reconstruction. When cellulose was used for regeneration antibodies against the 200 kDa NF protein showed neurofilament expression in nerve fibres at the distal tip of the injured nerve (III, Fig. 2C). This 200 kDa NF expression was comparable to that seen if the sciatic nerve was reconstructed using suturation (III, 2A). In the middle of the cellulose reconstructed nerve the 200 kDa NF protein was localized as well-organized nerve bundles (III, Fig. 2D) that appeared similar to the bundles of the sutured nerve (III, Fig. 2B). Thus, the 200 kDa NF positive nerve fibres had regenerated through the entire graft. Compared to suturation the cellulose treatment induced a fibrous scar around the site of injury, whereas no scar developed in-between the transected nerve stumps. These results indicate that cellulose grafts promoted neurite outgrowth through the injury zone.

3.2. Twitch tensions of the muscles after treatment

Twitch tensions of the muscles produced by electrical stimulation using either 0.01 ms or 0.1 ms single stimuli were not statistically different between the paper-grafted side and the uninjured control side (III, Fig. 1). If the nerve repair was performed using ordinary neurorrhaphy, the twitch tensions were significantly reduced on the injury side compared to the uninjured control side (III, Fig.1). Furthermore, the proportional muscle mass, the muscle mass of the trauma side compared to the control side, was significantly greater when the reconstruction was performed using paper compared to the ordinary neurorrhaphy.

3.3. The scores and rate of autotomy

The autotomy scores and the rates of autotomy were compared between paper reconstruction and neurorraphy. The incidence of autotomy was 50% in both groups. The latency of onset of autotomy was 2.3 ± 0.8 weeks (mean $\pm$ S.E.M.) in the neurorraphy group and 2.7 ± 0.4 weeks in the paper-treated group. The sutured and cellulose-treated groups did not differ significantly from each other regarding the time within which the maximal autotomy scores were reached. The mean times required were 2.7 ± 1.1 weeks in the sutured group and 3.6 ± 1.1 weeks in the cellulose-treated group. The final autotomy scores were 4.2 ± 2.1 weeks for the suturation group, and 4.0 ± 2.0 weeks for the paper-treated group.

IV NEUROFILAMENT PROTEINS ARE CONSTITUTIVELY EXPRESSED IN F9 TERATOCARCINOMA CELLS

4.1. Immunocytochemical analysis of induced F9 cells

Polyclonal antibodies against the neurofilament triplet protein were used to demonstrate that neurofilament proteins were expressed in the uninduced F9 cells grown in 10% fetal calf serum. Under these culture conditions the cells grew as embryonal bodies in which the neurofilaments were seen as short filamentous accumulations within the cell explants (IV, Fig. 2A). When antibodies against the neurofilament triplet proteins were absorbed with purified neurofilaments the filamentous immunoreactivity of the embryonal bodies was abolished (IV, Fig. 2C). Monoclonal antibodies against the 68 kDa neurofilament protein revealed similar filamentous accumulations in the uninduced F9 cell cultures (IV, Fig. 2B). A prolonged (10 days *in vitro*) cultivation of the F9 cells in 3% serum in the presence of RA and cAMP induced a neuronal phenotype with extensive neurite outgrowth and spreading of the F9 cells from the aggregates. Importantly, the F9 cells with a neuronal phenotype expressed the neurofilament triplet proteins (IV, Fig. 2D). Double immunocytochemistry for the 200 kDa neurofilament protein confirmed that the neuronal phenotype in these cultures was accompanied by expression of this mature type neurofilament protein (IV, Fig. 2E). These NF-positive neuron-like cells grew either as small clusters on top of the undifferentiated F9 cells, or were attached to the glass surface and sent out long neurites. (IV, Fig. 2E). In addition to neurofilament proteins (Figs. 2-3) the uninduced F9 cells also expressed other proteins involved in neuronal maturation, such

as N-CAM and TUJI (IV, Fig.3). The control cells (in 3% serum without RA and cAMP) failed to express immunocytochemically detectable neurofilament proteins (Fig. 3), but expressed both N-CAM and TUJI (Fig. 3).

4.2. Northern blot analysis of the induced F9 cells

Northern blot analysis showed that the 3.5 and 2.3 kb transcripts of the 68 kDa neurofilament protein gene (IV, Fig. I.) were expressed at the same level in the uninduced F9 cells cultured in 10% fetal calf serum and in the RA/cAMP induced F9 cells cultured in 3% serum (IV, Fig. I.). The uninduced F9 cells in 3% fetal calf serum did not express the 68 kDa neurofilament protein gene transcripts at detectable levels (IV, Fig. I.). Equal quantities of mRNA were loaded, as shown by ethidium bromide shadowing of each loaded mRNA. The mRNA levels were further evaluated by demonstration of β -actin mRNA levels in each sample using a cDNA probe for human β -actin. Each lane showed one sharp undegraded actin transcript at approximately 1.7 kb level (IV, Fig. I.). Densitometric scanning of each lane indicated that the levels of actin transcription were roughly comparable -between all samples.

4.3. RT-PCR analysis

RT-PCR analysis of both the 68 and 200 kDa NF gene transcripts further confirmed that the NF genes were constitutively expressed in the F9 cells. The 640 bp PCR product of the 200 kDa NF was expressed in uninduced (grown in 10% serum), induced (grown in 3% serum with RA or RA/dbcAMP) and control (grown in 3% serum) cultures of the F9 cells (IV, Fig 4). The 419 bp transcript of the 68 NF was also expressed in all conditions mentioned above. Sequencing of the 419 bp PCR products from all culture conditions verified that the 68 kDa NF gene products were identical and 98% similar to the cloned mouse NF 68 kDa gene.

4.4. Cloning of a neuronal F9 cell line

Single cells were picked from heterogeneous populations of F9 cells and grown in 10 % serum. The clones obtained were tested for their neuronal properties in a 24 hr assay in a serum free medium. One clone, D9L2, expressed both neurofilament triplet proteins, the 200 kDa neurofilament protein and TUJI (IV, Fig. 3 i-l). Thus, we conclude having successfully cloned a F9 cell line with neuronal properties.

DISCUSSION

The results of this thesis indicate that laminin-1 and its neurite outgrowth promoting γ 1-chain peptide (Liesi *et al.*, 1989) may be involved in neurodegenerative processes in Alzheimer's disease (I) and in the weaver mutant mouse (II). The results also show that thin grafts of paper may serve as potential carriers of laminin-1 and its neurite outgrowth promoting γ 1-chain peptide in attempts to regenerate peripheral nerves (III). Lastly, the present results indicate that laminin can be successfully used to subclone a novel neuronal cell line of the F9-teratocarcinoma cells (IV). This cell line may be used as a simplified model system to study the molecular mechanisms of neuronal differentiation.

I. Laminin and its neurite outgrowth-promoting domain in the brain in Alzheimer's disease and Down's syndrome patients

Even though laminin-1 (i.e. the isoform of laminins that is best characterized for its function in the nervous tissue) is generally known to promote neurite outgrowth and regeneration in both the CNS and the PNS (Liesi, 1990), results from this laboratory and other groups have merged to suggest that laminin-1 and its other isoforms may act in soluble form (Liesi *et al.*, 1989; Colamarino and Tessier-Lavigne, 1995) and may have a dual neurotrophic-neurotoxic function (Liesi *et al.*, 1989). An 11 amino acid long peptide derived from A β , the major constituent of the Alzheimer's plaques (Masters *et al.*, 1985; Selkoe *et al.*, 1986), was also shown to have a dual neurotrophic/neurotoxic effect on primary neurons (Yankner *et al.*, 1990b). This similarity led us to investigate whether the γ 1-chain of laminin-1 is present in AD and Down's syndrome brains and could therefore participate in the neuronal death mechanism in these disorders.

We found that the immunocytochemical distribution of laminin-1 served as a reliable marker for both senile plaques and pre-plaques in Alzheimer's disease (I); such a highly specific expression of laminin-1 in Alzheimer's disease and Down's syndrome brains was a novel finding. Previously, laminin-1 was demonstrated in brains of Alzheimer's disease patients in areas surrounding capillaries (Snow *et al.*, 1990) and it was thought to leak through the capillaries (Perlmutter and Chiu, 1990). However, accumulation of laminin-1 nearby the capillaries in AD brains may not be essential for pathophysiology of the disease, because similar accumulations were also detected in normal brain tissue (I). In contrast, normal control brains showed no expression of the punctate deposits of

laminin-1 (I), which further supported the view point that deposition of laminin-1 in the plaque areas could play a role in pathophysiology of Alzheimer's disease. E.g. the punctate deposits of laminin-1 in AD plaques could attempt to enhance neurite outgrowth and be responsible for sprouting events shown to take place in Alzheimer's brains (Scheibel and Tomiyasu, 1978; Geddes *et al.*, 1986). This point of view is feasibly based on the localization of similar punctate deposits of laminin-1 along growing axon tracts during embryonic brain development (Cohen *et al.*, 1987; Liesi and Silver, 1988; Letourneau *et al.*, 1988).

Antibodies against the γ 1-chain neurite outgrowth promoting domain did not localize in as large punctate deposits in the plaques (I), which might indicate that laminin-1 was deposited in the plaque regions in such a conformation that the antigenic epitopes recognized by the peptide antibodies could be hidden or blocked. E.g. HSPGs, also present in the plaque regions (Snow *et al.*, 1990), could bind to the heparin binding domains of laminin-1 (Engel, 1991) and prevent the binding of peptide antibodies to the neurite outgrowth domain of the γ 1-chain close to the heparin binding site. Alternatively, the neurite outgrowth domain of the γ 1-chain might have been degraded in the plaque areas and could not be recognized by the antibodies. The C-terminal parts of laminin are known to be more sensitive to proteolysis than other parts of the molecule (Ott *et al.*, 1982). Both the plaques and the reactive astrocytes of the Alzheimer's brain are rich in lysosomal proteinases, such as cathepsins B and D (Cataldo *et al.*, 1990) and these enzymes could degrade laminin (Heck *et al.*, 1990; Steadman *et al.*, 1993; Buck *et al.*, 1992; Guinec *et al.*, 1993). If the neurite outgrowth domain of the γ 1-chain has been degraded, however, sprouting in the plaques could also be due to the growth factor-like properties of laminin-1 (Panayotou *et al.*, 1989) or other growth factors in the plaque regions (Birecree *et al.*, 1988; Stopa *et al.*, 1990; Fenton *et al.*, 1998).

Instead of a localization as large punctate deposits in the plaques, the γ 1-chain peptide antibodies recognized fine punctate deposits of the neurite outgrowth domain in AD brain tissue (I). These fine punctate deposits were detected by antibodies that recognized a 10 amino acid neurite outgrowth domain of the γ 1-chain but not native laminin (I). This result indicates that proteolytic degradation of laminin-1 into smaller peptides, antigenically similar to the peptides used to produce the γ 1-chain specific antibodies, might occur in the Alzheimer's brain. The fact that glial cells in Alzheimer's disease and

in Down's syndrome brains were strongly immunoreactive for the γ 1-chain peptide while native laminin-1 antibodies showed only weak staining of glial cells suggests the glial laminin was rich in the γ 1-chain or had antigenic epitopes needed for recognition of the γ 1-chain better exposed.

A role for laminin-1 in AD is further suggested by the fact that the IKVAV-peptide from the α 1-chain of laminin-1 has been shown to bind APP (Kibbey *et al.*, 1993). This binding may facilitate the deposition of APP and amyloid in plaques (Kibbey *et al.*, 1993). We failed to detect mRNA for the α 1-chain of laminin-1 in AD or control brains (I). Therefore, if the IKVAV sequence exists in plaques, there must be an uncharacterized α -chain or some other neurite outgrowth promoting protein carrying the IKVAV sequence. Recent studies have shown that laminin reduces the fibril formation of amyloid- β -peptide (A β 1-42) *in vitro* (Bronfman *et al.*, 1996a, 1996b, 1998; Monji *et al.*, 1998a, 1998b; Drouet *et al.*, 1999) and modulates the biogenesis of APP (Monning *et al.*, 1995; Coulson *et al.*, 1997). These results indicate that laminin-1 and its γ 1-chain may have a direct interaction with APP and that this interaction may be have importance for the pathophysiology of Alzheimer's disease.

The induction of both laminin-1 and APP occurs in reactive astrocytes of the injured adult rodent brain (Liesi *et al.*, 1984; Siman *et al.*, 1989). Here we report an approximately 10-fold increase in laminin-1 γ 1-chain expression in AD brains compared to control brains (I). The increased expression of laminin-1 and its γ 1-chain may be due to the response of the diseased brain to the tissue injury occurring in AD brains. Factors involved in the gene regulation for the expression of laminin-1 are largely unknown. However, expression of the γ 1-chain of laminin-1 is known to be induced by interleukin-1 beta (Richardson *et al.*, 1995) or Sp-1 transcription factors (Lietard *et al.*, 1997). Both interleukin-1 beta and Sp-1 are induced by trauma (Giulian and Lachman, 1985; Pearson *et al.*, 1999; Feng *et al.*, 1999). The trauma-induced over-expression of both APP and laminin-1 together with increased proteolysis could release A β and laminin-1 γ 1-chain peptides in the AD brain tissue. As the γ 1-chain peptide of laminin-1 is neurotoxic at high concentrations *in vitro* (Liesi *et al.*, 1989), its binding and accumulation in the AD brain tissue may produce a synergistic neurotoxic effect together with A β peptide. This neurotoxicity can be further enhanced by the release of excitatory neurotransmitters (Koh *et al.*, 1990) and growth factors (Yankner *et al.*, 1990a,b; Kowall *et al.*, 1991). Thus, we

hypothesize that laminin-1 synthesis is initially induced as tissue response to trauma with no direct link to the disease. However, laminin-1 may be involved in neuronal death in both AD and Down's syndrome via its interaction with APP (Narindrasorasak *et al.*, 1992) and its toxic peptides as well as via accumulation of the neurotoxic γ 1-chain peptides in plaques and the brain tissue.

II Increased proteolytic activity of the granule neurons may contribute to neuronal death in the weaver mouse cerebellum

Increased expression of the γ 1-chain neurite outgrowth peptide (Liesi *et al.*, 1989) in AD brain tissue (I) suggested that this peptide may be involved in neuronal death. We used the weaver mutant mouse, an animal model of neuronal degeneration, to study further the possible involvement of the γ 1-chain peptide in neuronal death. We found that expression of laminin-1, and its γ 1-chain neurite outgrowth domain were elevated in the cerebellum of the weaver mutant mouse (II). Expression of tissue plasminogen activator (tPA) in the weaver cerebellum was also high compared to the normal cerebellum (II). As tPA co-localized with laminin-1 in the weaver cerebellum (II), increased expression of both tPA and laminin-1 and its γ 1-chain may result in increased proteolytic cleavage of laminin-1 and its γ 1-chain. Thus, toxic amounts of peptides derived from the γ 1-chain neurite outgrowth domain might accumulate in the weaver cerebellum and result in massive neuronal death. This point of view is supported by the fact that weaver granule neurons degrade their laminin-1 substratum (II) and bind increased amounts of the neurite outgrowth domain of the γ 1-chain (II). Importantly, weaver neurons can be rescued by a serine protease inhibitor aprotinin (II) or by antibodies against the γ 1-chain peptide (Liesi and Wright, 1996). As both the RMPs and the neurite outgrowth potential of weaver granule neurons can be rescued by aprotinin (II), increased proteolysis may be one of the primary defects in the weaver mouse cerebellum.

The role of tPA in neuronal degeneration and degradation of laminin has been demonstrated by the fact that mice deficient for tPA or plasminogen are resistant to neuronal degeneration (Tsirka *et al.*, 1997), and that degradation of laminin by tPA proteolysis is shown to precede neuronal death after an injection of excitotoxin (Chen and Strickland, 1997; Nagai, *et al.*, 1999). Recent results by Mecnas *et al.* (1997)

contradict these data and show that neurons in tPA $-/-$ homozygous weaver mice are not rescued. It is not currently known why Mecnas *et al.* failed to rescue the weaver neurons, but it is possible that the complete lack of tPA in the *wv/wv-tPA-tPA-* mice opposed to a reduction of tPA-proteolysis by aprotinin (II) is generally harmful to neurons, especially since tPA induced proteolysis is known to be essential for neuronal migration (Kalderon, 1982; Seeds *et al.*, 1990).

In addition to the abnormal laminin-1 expression and proteolytic activity of the weaver neurons (II), additional molecular mechanisms may impair neuronal migration in the weaver mouse cerebellum. A point mutation of the GIRK2 potassium channel gene has been proposed to be responsible for the weaver phenotype (Patil *et al.*, 1995; Slesinger *et al.*, 1996). However, the importance of the GIRK2 gene as a weaver gene is fading because electrophysiological experiments have failed to detect functional GIRK2 channels in the cultured weaver granule neurons (Mjaatved *et al.*, 1995; Surmeier *et al.*, 1996). Furthermore, the GIRK2 knockout mice have normal cerebella (Signorini *et al.*, 1997), which indicates that the GIRK2 channel is not essential for early postnatal development of the cerebellum and cannot therefore be the weaver gene. Rescue of the weaver granule neurons from death by verapamil a L-type calcium channel blocker (Liesi and Wright, 1996), or other means that reduce the levels of intracytoplasmic calcium (Liesi *et al.*, 1997) strongly suggest that the weaver gene action is mediated by calcium-dependent mechanisms. The Weaver granule neurons fail to express functional N-methyl-D-aspartate (NMDA) receptors (Liesi and Wright, 1996), which may be due to the fact that the $\epsilon 2$ subunit is absent in the weaver cerebellum (Liesi *et al.*, 1999). The $\epsilon 2$ subunit is induced in the granule neurons after rescue with verapamil (Liesi and Wright, 1996; Liesi *et al.*, 1999), which suggests that the down regulation of NMDA receptors may be a protective measure to reduce calcium entry into weaver granule neurons via functional NMDA-receptors. As NMDA receptors have been shown to play a role in neuronal migration (Komuro and Rakic, 1992, 1993), the lack of NMDA-receptor function may result in the expression of the weaver phenotype (Liesi and Wright, 1996; Liesi *et al.*, 1999). The role of NMDA receptors in the expression of the weaver phenotype is further emphasized by results that blocking of the NMDA receptor function in the homozygous weaver mice by eliminating the $\zeta 1$ -subunit rescues the weaver granule neurons from death (Jensen *et al.*, 1999).

III Use of cellulose for treatment of a peripheral nerve trauma in the rat

Laminin-1 and its $\gamma 1$ -chain domain support axon growth of CNS neurons (Matsuzawa *et al.*, 1996; 1997). Laminin-1 is known to associate with the regenerating CNS (Liesi 1985b) and peripheral nerves have been shown to grow along the laminin-rich basement membranes (Ide *et al.*, 1983). Thus, laminin-1 grafts have been used by several laboratories in attempts to repair peripheral nerve injuries (Madison *et al.*, 1985; 1987; Bailey *et al.*, 1993; Tong *et al.*, 1994; Labrador *et al.*, 1998 Kauppila *et al.*, 1993). Laminin-1 or its $\gamma 1$ -chain neurite outgrowth promoting peptide (Liesi *et al.*, 1989) coupled with a type-I-collagen have been used to support neuronal regeneration *in vivo* (Kauppila *et al.*, 1993). These grafts supported regeneration comparable to that achieved by suturation (Kauppila *et al.*, 1993). The main limitation for the use of such laminin-1 grafts was the thickness, which made these grafts difficult to handle and caused compression of damaged nerves. As laminin-1 and its $\gamma 1$ -chain neurite outgrowth peptide were found effective in peripheral nerve regeneration (Kauppila *et al.*, 1993), the development of more suitable graft materials is needed.

We therefore tested thin cellulose grafts for their ability to regenerate peripheral nerves. We applied cellulose because it has been successfully used for the treatment of burns as well as in otolaryngology in humans (Palva, 1982; Hazarika, 1985) and could be immediately applied to human nerve surgery. The use of cellulose grafts was more feasible than using collagen-I grafts, since proteins could be covalently coupled to cellulose. Thin sheets of cellulose were used to reconstruct severed peripheral nerves in rats (III). Cellulose was sticky and allowed good positioning of the severed nerves, but provoked a stronger foreign body reaction inflammation than resuturation (III). When cellulose grafts were used in restorative surgery the twitch-induced forces of the muscles between the operated and control sides were identical (III), which was an unexpected result. In sutured animals the twitch-induced forces of muscles decreased as compared to control side. The latter result is consistent with earlier studies by Brunetti *et al.* (1985) and Kauppila *et al.* (1993). Also macroanatomical measurements of the muscle mass favor the hypothesis that cellulose around the trauma site may favor neuroregeneration. In fact, the proportional muscle mass had a tendency to increase when cellulose grafting was used for reconstruction. Increase in the proportional muscle mass is considered as a measure of nerve regeneration (Kauppila *et al.*, 1993; Kauppila, 1994; Greensmith *et al.*,

1995). Thus, our results indicate that nerve regeneration improved using cellulose grafts compared to conventional suture.

Thin cellulose grafts were found to induce scar formation around the trauma site (III), whereas scar formation was minimal if reconstruction was made by suture. Inflammation near the trauma site is known to promote regeneration of both rat dorsal root and sciatic nerve injuries (Lu and Ricahardson, 1991; Dahlin, 1992), which suggests that scar-formation may be one of the factors in cellulose grafting that supports regeneration.

Cellulose grafting was as efficient as suture in preventing dyesthesias induced by self-mutilation. The low autotomy scores of both cellulose grafting and suture indicated that cellulose grafting effectively supported sensory regeneration of the denervated paws. This could be concluded, since the high incidence of autotomy correlates with poor recovery (Kauppila, 1994) and autotomy is known to disappear when regenerating axons form their connections (Wall *et al.*, 1979; Kauppila *et al.*, 1993; DeLeo *et al.*, 1994; Kauppila, 1994).

The present results suggest that cellulose grafts may be used to repair peripheral nerves. However, additional research will be required before this technique will be clinically applicable. This is due to the fact that rats have only few fascicles in their sciatic nerves and therefore the repair of rat nerves differs from a normal clinical situation in which several adjacent fascicles are damaged and need to be rejoined. Scar formation that occurs in cellulose grafting may result in intraneural fibrosis in humans and hamper regeneration, although regeneration of rat neurons occurred successfully.

IV Neurofilament proteins are constitutively expressed in F9 teratocarcinoma cells

Neuronal differentiation has been shown to be under a negative control, i.e. cells will become neurons if they do not receive inductive signals to become other cell types (Hemmati-Brivanlou and Melton, 1997). Bone morphogenetic protein (BMP) is one of the recently identified factors involved in initial neuronal induction (Wilson and Hemmati-Brivanlou, 1995). Binding of neuronal inducers (such as noggin, follistatin, and chordin) to BMP inactivates BMP and results in expression of several transcription

factors that promote the neuronal lineage (Sasai, 1998). Until now, neurogenesis of mammals has been thought to occur during the embryonic and early postnatal period. However, recent research has shown that the adult mammalian brain has neural stem cells that can give rise to both neurons and glial cells (Reynolds and Weiss, 1992, 1996; Lois and Alvarez-Buylla, 1993; Morshead *et al.*, 1994; Palmer *et al.*, 1997; Johansson *et al.*, 1999; Chiasson *et al.*, 1999). These adult stem cells are very similar to those present in the embryonic brain and similar mechanisms are therefore thought to induce their differentiation into various cell types (Johe *et al.*, 1996; Palmer *et al.*, 1997; Johansson *et al.*, 1999; Chiasson *et al.*, 1999).

Apart from normal stem cells neuronal differentiation has been studied extensively using teratocarcinoma cell lines (Pleiffer *et al.*, 1981, McBurney *et al.*, 1982, Levine and Flynn 1986, Kubo 1989; Kuff and Fewell, 1980, Liesi *et al.*, 1983). As an approach to the molecular mechanisms of neuronal differentiation we used F9 teratocarcinoma cells as a model system (IV). We chose this cell line, because Liesi *et al.* (1983) have previously shown that the F9 cells can choose a neuronal lineage when exposed to RA/dbcAMP under serum deprivation. Our studies confirmed these results and also showed that the uninduced F9 cells, grown in 10 % serum, expressed both the 68 kDa neurofilament gene transcripts and protein without having a neuronal phenotype (IV). Even without the neuronal phenotype these cells expressed additional neuronal markers, such as the neuron specific tubulin III isoform (TUJI) and N-CAM. RT-PCR studies showed that both the 68 kDa and the 200 kDa neurofilament (NF) gene transcripts were constitutively expressed by F9 cells (IV). The present results confirm and expand the earlier results by Liesi *et al.* (1983). We show here that RA/dbcAMP treatment of the F9 cells is not required for their expression of neurofilament genes and proteins (IV). However, the expression of a neuronal phenotype by the F9 cells appeared to depend on serum deprivation and RA/dbcAMP stimulation (IV).

Even though the neuronal differential potential of F9 cells has been published by several laboratories, contradictory results have also appeared. The F9 cells have also been found not to express NF-proteins (Tienari *et al.*, 1987). This controversy is not presently understood, but it is possible that laboratories that failed to induce/demonstrate neurofilament expression of the F9 cells may have used a cell line in which the non-

neuronal subpopulation of F9 cells might have been selected over the neuronally differentiating one.

This view is feasible, because we verified that only a subpopulation of F9 cells developed into neuron-like cells under serum deprivation and exposure to RA/dbcAMP (IV). Thus, the F9 cells could not be used in biochemical or molecular studies on neuronal differentiation. In order to solve this problem we subcloned the F9 cells on a laminin-1 substratum by using a single parent cell isolation technique. Using this technique we obtained a homogenous, neuronally differentiating D9L2 cell line that 1) expressed the 200 kDa NF protein and 2) had a neuronal phenotype on the regular tissue culture plastic in a serum free medium without RA and dbcAMP. Laminin-1 was chosen as an initial growth substratum of the isolated cells, because earlier studies indicate that laminin-1 promotes neuronal of several types of neurons, such as early neuroepithelial cells (Heaton and Swanson, 1988; Frade *et al.*, 1996), embryonic hippocampal neurons (Lein *et al.*, 1992), sympathetic neurons (Chu and Tolkovsky, 1994), and enteric neurons (Chalazonitis *et al.*, 1997). Our results indicate that laminin-1 indeed is a favored substratum for the cloning of a neuronal D9L2 cell line. This cell line is currently being used in our laboratory to study the molecular mechanisms of neuronal differentiation.

SUMMARY AND CONCLUSIONS

In this Thesis, I have studied the potential role of laminin-1 and its γ 1-chain neurite outgrowth domain in Alzheimer's disease and in the weaver mutant mouse model. Based on previous results on the role of laminin-1 I have tested cellulose as a suitable graft material in the repair of neuronal injuries and used laminin-1 to develop a neuronal F9 cell line for future studies on neuronal differentiation.

We wanted to study the expression of laminin-1 and its γ 1-chain peptide in Alzheimer's disease and Down's syndrome brains, because both the neurite outgrowth domain of the γ 1-chain of laminin-1 and the A β -peptide were shown to have a dual concentration-dependent neurotrophic/neurotoxic effect. We found that laminin-1 and the neurite outgrowth promoting domain of its γ 1-chain accumulate in Alzheimer's disease and Down's syndrome brains, but not in normal control brains. The punctate deposits of laminin localize in the Alzheimer's plaques and antibodies against the neurite outgrowth promoting domain of the γ 1-chain detect both the extracellular γ 1-chain deposits and glial cells in the diseased brains, but not in healthy control brains. These results suggest that deposition of laminin-1 in plaques, and its γ 1-chain in the astrocytes and Alzheimer brain tissue may either promote sprouting of the affected neurons or contribute to the neurotoxic mechanisms that cause neuronal death in Alzheimer's disease.

Our results on the increased expression of laminin-1 and its γ 1-chain neurite outgrowth domain in Alzheimer's disease and Down syndrome brains implied that laminin and its γ 1-chain may be involved in neuronal death mechanisms. To study this in detail we used weaver mutant mice as an experimental model system for neuronal migration defects and neuronal degeneration. We found that the weaver mouse cerebellum shows an increased expression of laminin-1, and its γ 1-chain. Increased proteolytic activity in the weaver cerebellum may lead to degradation of laminin-1, and accumulation of the neurite outgrowth domain on the surfaces of the weaver granule cells. This may result in accumulation of neurotoxic peptides that provoke the death of weaver neurons.

The role of laminin-1 in peripheral nerve regeneration has been verified in an earlier study, but suitable graft materials have not yet been found for the coupling of laminin-1 and its biologically active peptides. Therefore, we tested cellulose as a graft material to

be coupled with laminin or synthetic peptides with a neurite outgrowth promoting activity. We found that cellulose grafts induced more fibrous scarring around the transection site compared to the microsurgical neurotaphy. However, this scarring did not impair functional recovery or cause signs of neuropathic pain. Thus, our results indicate that cellulose may be a potentially useful material in the repair of peripheral nerves.

Laminin-1 is known to promote neuronal differentiation. Therefore, we used laminin-1 to subclone a novel neuronal cell line of the F9 teratocarcinoma cells. This was necessary, because we found that only a subpopulation of F9 cells expressed neuronal markers and could be differentiated into cells with a neuronal phenotype. In the course of our study, we found that the undifferentiated F9 cells constitutively expressed both the 68 kDa and 200 kDa neurofilament transcripts and proteins as well as other neuronal marker proteins. These results indicate that RA and dbcAMP are not necessary for neurofilament gene expression in F9 cells. Instead, they may be required for the expression of a neuronal phenotype by the F9 cells.

LIST OF REFERENCES

- Abemayor**, E., and Sidell, N. Human neuroblastoma cell lines as models for the in vitro study of neoplastic and neuronal cell differentiation. *Environ. Health. Perspect.*, 80, 3-15, 1989
- Abraham**, C.R., Selkoe, D.J. and Potter, H. Immunochemical identification of the serine protease inhibitor α_1 -antichymotrypsin in the brain amyloid deposits of Alzheimer's disease. *Cell*, 52, 487-501, 1988
- Aigner**, L., Arber, S., Kapfhammer, J.P., Laux, T., Schneider, C., Botteri, F., Brenner, F., and Caroni, P. Overexpression of the neural growth-associated protein GAP-43 induces nerve sprouting in the adult nervous system of transgenic mice. *Cell*, 83, 269-278, 1995
- Akers**, R.M., Mosher, D.F., and Lilien, J.E. Promotion of retinal neurite outgrowth by substratum-bound fibronectin. *Dev. Biol.*, 86, 179-188, 1981
- Alexianu**, M.E., Mohamed, A.H., Smith, R.G., Colom, L.V., and Appel, S.H. Apoptotic cell death of a hybrid motoneuron cell line induced by immunoglobulins from patients with amyotrophic lateral sclerosis. *J. Neurochem.*, 63, 2365-2368, 1994
- Allan**, K.M., and Walsh, C.A. Genes that regulate neuronal migration in the cerebral cortex. *Epilepsy Res.*, 36, 143-154, 1999
- Almqvist**, E. Nerve repair by laser. *Orthopedic Clinics of North America*, 19, 201-208, 1988
- Al Mutlag**, H., Wheeler, J., Robertson, H., Watchorn, C., and Morley, A.R. Tissue distribution of amyloid p component as defined by a monoclonal antibody produced by immunisation with human basement membranes. *Histochem. J.*, 25, 219-227, 1993
- Alzheimer's Disease Collaborative Group**. The structure of the presenilin-1 (S182) gene and identification of six novel mutations in early onset AD families. *Nat. Genet.*, 11, 219-222, 1995
- Anderson**, A.J., Su, J.H., and Cotman, C.W. DNA damage and apoptosis in Alzheimer's disease: colocalization with c-Jun immunoreactivity, relationship to brain area, and effect of postmortem delay. *J. Neurosci.*, 16, 1710-1719, 1996
- Andrews**, P.W. Teratocarcinomas and human embryology: pluripotent human EC cell line. *APMIS*, 106, 158-168, 1998
- Ankarcrona**, M., Dypbukt, J.M., Bonfoco, E., Zhivotovsky, Orrenius, S., Lipton, S.A., and Nicotera, P. Glutamate-induced neuronal death: A succession of necrosis or apoptosis depending on mitochondrial function. *Neuron*, 15, 961-973, 1995
- Anton**, E.S., Sandrock, A.W.Jr., and Matthew, W.D. Merosin promotes neurite growth and Schwann cell migration *in vitro* and nerve regeneration *in vivo*: Evidence using an antibody to merosin, ARM-1. *Dev. Biol.*, 164, 133-146, 1994
- Antonicsek**, H., Persohn, E., and Schachner, M. Biochemical and functional characterisation of a novel neuron-glia adhesion molecule that is involved in neuronal migration. *J. Cell Biol.*, 104, 1587-1595, 1987
- Arber**, S., and Caroni, P. Thrombospondin-4, an extracellular matrix protein expressed in the developing and adult nervous system promotes neurite outgrowth. *J. Cell Biol.*, 131, 1083-1094, 1995
- Arends**, M.J., and Wyllie, A.H. Apoptosis: mechanisms and roles in pathology. *Int. Rev. Exp. Pathol.*, 32, 223-254, 1991

- Armstrong, J.W.**, and Chapes, S.K. Effects of extracellular matrix proteins on macrophage differentiation, growth, and function- comparison of liquid and agar culture systems. *J. Exp. Zool.*, 269, 178-187, 1994
- Arumugham, R.G.**, Hsieh, T.C.Y., Tanzer, M.L., and Laine, R.A. Structure of the asparagine-linked sugar chains of laminin. *Biochim. Biophys. Acta.*, 883, 112-126, 1986
- Aumailley, M.**, Gerl, M., Sonnenberg, A., Deutzmann, R., and Timpl, R. Identification of the Arg-Gly-Asp sequence in laminin A chain as a latent cell-binding site being exposed in fragment P1. *FEBS Lett.*, 262, 82-86, 1990
- Aumailley, M.**, and Grieg, T. Laminins: A family of diverse multifunctional molecules of basement membranes. *J. Invest. Dermatol.*, 106, 209-214, 1996
- Aumailley, M.**, Nurcombe, V., Edgar, D., Paulsson, M., and Timpl, R. The cellular interactions of laminin fragments. Cell adhesion correlates with two fragment-specific high affinity binding sites. *J. Biol. Chem.*, 262, 11532-11538, 1987
- Aumailley, M.**, and Smyth, N. The role of laminins in basement membrane function. *J. Anat.*, 193, 1-21, 1998
- Bailey, S.B.**, Eichler, M-E-, Villadiego, A., and Rich, K.M. The influence of fibronectin and laminin during Schwann cell migration and peripheral nerve regeneration through silicon chambers. *J. Neurocytol.*, 22, 176-184, 1993
- Baird, D.**, Baptista, C., Wang, L-C., and Mason, C. Specificity of a target cell-derived stop signal for afferent axonal growth. *J. Neurobiol.*, 23, 579-591, 1992
- Baron Van Evercooren, A.**, Kleinmann, H., Ohno, S., Marangos, P., Schwartz, J. and Dubois-Dalcq, M. Nerve Growth factor, laminin and fibronectin promote neurite outgrowth in human fetal sensory ganglia cultures. *J. Neurosci Res.* 8, 179-193, 1982
- Beck, K.**, Hunter, I. and Engel, I. Structure and function of laminin: anatomy of multidomain glycoprotein. *FASEB J.* 4, 148-160, 1990
- Ben-Arie, N.**, Bellen, H.J., Armstrong, D.L., McCall, A.E., Gordadze, P.R., Guo, Q., Matzuk, M.M., and Zoghbi, H.Y. *Mat 1* is essential for genesis of cerebellar granule neurons. *Nature*, 390, 169-172, 1997
- Benfey, M.**, and Aguayo, A. Extensive elongation of axons from rat brain into peripheral nerve grafts. *Nature* 269, 150-152, 1982
- Bernier, S.M.**, Utani, A., Sugiyama, S, M Doi, T., Polistina, C., and Yamada, Y. Cloning and expression of laminin $\alpha 2$ chain (M-chain) in the mouse. *Matrix Biol.*, 14, 447-455, 1994
- Bignami, A.** and Dahl, D. Astrocyte specific protein and radial glia in the cerebral cortex of newborn rat. *Nature*, 252, 55-56, 1974a
- Bignami, A.** and Dahl, D. The development of Bergmann glia in mutant mice with cerebellar malformations: Reeler, staggerer and weaver. Immunofluorescence study with antibodies to the glial fibrillary acidic protein. *J. Comp. Neurol.*, 155, 219-230, 1974b
- Billingsley, G.D.**, Walter, M.A., Hammond, G.L., and Cox, D.W. Physical mapping of four serpin genes: alpha 1-antitrypsin, alpha 1-antichymotrypsin, corticosteroid-binding globulin, and protein c inhibitor, within a 280-kb region on chromosome 14g32.1. *Am. J. Hum. Gen.*, 52, 343.353, 1993
- Birecree, E.**, Whetsell, W.O. Jr., Stoscheck, C., King, L.E. Jr., and Nanney, L.B. Immunoreactive epidermal growth factor receptors in neuritic plaques from patients with Alzheimer's disease. *J. Neuropathol. Exp. Neurol.*, 47, 549-560, 1988
- Bixby, J.L.**, and Zhang, R. Purified N-cadherin is a potent substrate for the rapid induction of neurite outgrowth. *J. Cell Biol.*, 110, 1253-1260, 1990

- Björklund, A.**, and Stenevi, U. Intracerebral neural implants: Neuronal replacement and reconstruction of damaged circuitries. *Annu. Rev. Neurosci.*, 7, 279-308, 1984
- Blass, J.P.**, and Gibson, G.E. The role of oxidative abnormalities in the pathophysiology of Alzheimer's disease. *Rev. Neurol.*, 147, 513-525, 1991
- Book, K.J.**, and Morest, D.K. Migration of neuroblasts by perikaryal translocation: Role of cellular elongation and axon outgrowth in the acoustic nuclei of the chick embryo medulla. *J. Comp. Neurol.* 297, 55-76, 1990
- Boucaut, J.C.**, Darribere, T., Poole, T.J., Aoyama, H., Yamada, K.M., and Thiery, J.P. Biologically active synthetic peptides as probes of embryonic development: a competitive peptide inhibitor of fibronectin function inhibits gastrulation in amphibian embryos and neural crest cell migration in avian embryos. *J. Cell Biol.*, 99, 1822-1830, 1984
- Bourdon, M.A.**, Wikstrand, C.J., Furthmayer, H., Matthews, T.J., and Bigner, D.D. Human gliomamesenchymal extracellular matrix antigen defined by monoclonal antibody. *Cancer Res.*, 43, 2796-2805, 1983
- Bouzon, M.**, Dussert, C., Lissitzky, J.C., and Martin, P.M. Spreading of B16 F1 cells on laminin and its proteolytic fragments P1 and E8: Involvement of laminin carbohydrate chains. *Exp. Cell. Res.*, 190, 47-56, 1990
- Brandenberger, R.**, Kammerer, R.A., Engel, J., and Chiquet, M. Native chick laminin-4 containing the $\beta 2$ chain (s-laminin) promotes motor axon growth. *J. Cell Biol.*, 135, 1583-1592, 1996
- Bregman, B.S.**, Kunkel-Bagden, E., Schnell, L., Dai, H.N., Gao, D., and Schwab, M.E. Recovery from spinal cord injury mediated by antibodies to neurite growth inhibitors. *Nature*, 378, 498-501, 1995
- Bronfman, F.C.**, Alvarez, A., Morgan, C., and Inestrosa, N.C. Laminin blocks the assembly of wild-type A beta and the Dutch variant peptide in to Alzheimer's fibrils. *Amyloid*, 5, 16-23, 1998
- Bronfman, F.C.**, Garrido, J., Alvarez, A., Morgan, C., and Inestrosa, N.C. Laminin inhibits amyloid- β -peptide fibrillation. *Neurosci. Lett.*, 218, 201-203, 1996a
- Bronfman, F.C.**, Soto, C., Tapia, L. Tapia, V., and Inestrosa, N.C. Extracellular matrix regulates the amount of the β -amyloid precursor protein and its amyloidogenic fragments *J. Cell. Physiol.* 166, 360-369, 1996b
- Bronner-Fraser, M.** An antibody to a receptor for fibronectin and laminin perturbs cranial neural crest migration *in vivo*. *Dev. Biol.*, 117, 528-536, 1986
- Bruch, M.**, Landwehr, R., and Engel, J. Dissection of laminin by cathepsin G into its long-arm and short-arm structures and localization of regions involved in calcium dependent stabilization and self-association. *Eur. J. Biochem.*, 185, 271-279, 1989
- Brundin, P.**, Isacson, O., and Björklund, A. Monitoring of cell viability in suspension of embryonic CNS tissue and its use as criterion for intra cerebral graft survival. *Brain Res.*, 331, 251-259, 1985
- Brunetti, O.**, Magni, F., and Pazzaglia, U. Neural nature of the time dependent factor(s) involved in motor reinnervation. *Exp. Neurol.*, 90, 667-685, 1985
- Buck, M.R.**, Karustis, D.G., Day, N.A., Honn, K.V., and Sloane, B.F. Degradation of extracellular-matrix proteins by human cathepsin B from normal and tumour tissues. *Biochem. J.*, 282, 273-278, 1992
- Burgeson, R.E.**, Chiquet, M., Deutzmann, R., Ekblom, P., Engel, J., Kleinmann, H., Martin, G.R., Meneguzzi, G., Paulsson, M., Sanes, J., Timpl, R., Tryggvason, K., Yamada, Y. and Yurchenco, P.D. A new nomenclature for the laminins. *Matrix Biol.* 14, 209-211, 1994
- Calof, A.L.**, and Lander, A.D. Relationship between neuronal migration and cell-substratum adhesion: laminin and merosin promote olfactory neuronal migration but are

anti-adhesive. *J. Cell Biol.*, 115, 779-794, 1991

Campagna, J.A., Ruegg, M.A., and Bixby, J.L. Agrin is a differentiation-inducing "stop signal" for motoneurons *in vitro*. *Neuron*, 15, 1365-1374, 1995

Caroni, P., and Schwab, M.E. Two membrane protein fractions from rat central myelin with inhibitory properties for neurite growth and fibroblast spreading. *J. Cell Biol.*, 106, 1281-1288, 1988a

Caroni, P., and Schwab, M.E. Antibody against myelin-associated inhibitor of neurite growth neutralizes nonpermissive properties of CNS white matter. *Neuron*, 1, 85-96, 1988b

Castillo, G.M., Ngo, C., Cummings, J., Wight, T.N., and Snow, A.D. Perlecan binds to the beta-amyloid proteins (A beta) of Alzheimer's disease, accelerates A beta fibril formation, and maintains A beta fibril stability. *J. Neurochem.*, 69, 2452-2465, 1997

Cataldo, A.M., Thayer, C.Y., Bird, E.D., Wheelock, T.R. and Nixon, R.A. Lysosomal proteinase antigens are prominently localized within senile plaques of Alzheimer's disease: Evidence for a neuronal origin. *Brain Res.* 513, 181.192, 1990

Chagnon, P., Betard, C., Robitaille, Y., Cholette, A., and Gauvreau, D. Distribution of brain cytochrome oxidase activity in various neurodegenerative diseases. *Neuroreport*, 6, 711-715, 1995

Chalazonitis, A., Tennyson, V.N., Kibbey, M.C., Rothman, T.P., and Gershon, M.D. The alpha 1 subunit of laminin-1 promotes the development of neurons by interacting with LBP110 expressed by neuronal crest-derived cells immunoselected from the fetal mouse gut. *J. Neurobiol.*, 33, 118-138, 1997

Chamberlain, L.J., Yannas, I.V., Hsu, H.P., Strichartz, G., and Spector, M. Collagen-GAG substrate enhances the quality of nerve regeneration through collagen tubes up to level of autograft. *Exp. Neurol.*, 154, 315-329, 1998

Chammas, R., Veiga, S.S., Line, S., Potocnandk, P., and Brentani, R.R. An-linked oligosaccharide-dependent interaction between laminin and gp120/140. An alpha 6/beta 1 integrin. *J. Biol. Chem.*, 226, 3349-3355, 1991

Champlaud, M.-F., Lunstrum, G.P., Rousselle, P., Nishiyama, T., Keene, D.R., and Burgeson, R.E. Human amnion contains a novel laminin variant, laminin 7, which like laminin 6, covalently associates with laminin 5 to promote stable epithelial-stromal attachment. *J. Cell Biol.*, 132, 1189-1198, 1996

Chandrasekaran, S., Dean, J.W., Griniger, M.S., and Tanzer, M.L. Laminin carbohydrates are implicated in cell signalling. *J. Cell Biochem.* 46, 115-124, 1991

Charonis, A.S., Skubits, A.P., Koloakos, G.G., Reger, L.A., Dege, J., Vogel, A.M., Wohlhueter, R., and Furcht, L.T. A novel synthetic peptide from the B1 chain of laminin with heparin-binding and cell adhesion-promoting activities. *J. Cell. Biol.*, 107, 1253-1260, 1988

Charonis, A.S., Tsilibary, E.C., Saku, T., and Furthmayr, H. Inhibition of laminin self-assembly and interaction with type IV collagen by antibodies to the terminal domain of the long arm. *J. Cell Biol.*, 103, 1689-1697, 1986

Chen, D.F., Jhaveri, S., and Schneider, G.E. Intrinsic changes in developing retinal neurons result in regenerative failure of their axons. *Proc. Natl. Acad. Sci. USA*, 92, 7287-7291, 1995

Chen, D.F., Schneider, G.E., Martinou, J.-C., and Tonegawa, S. Bcl-2 promotes regeneration of severed axons in mammalian CNS. *Nature*, 385, 434-439, 1997a

Chen, F.-L., and Strickland, S. Neuronal death in the hippocampus is promoted by plasmin-catalyzed degradation of laminin. *Cell*, 91, 917-925, 1997

Chen, L., Shick, V., Marret, M.L., Laurie, S.M., Ogle, R.C., and Laurie, G.W. Laminin

E8 alveolarization site: heparin sensitivity, cell surface receptors, and role in cell spreading. *Am. J. Physiol.*, 272, 494-503, 1997b

Chen, M., Marinkovich, M.P., Jones, J.C., O'Toole, E.A., Li, Y.Y., and Woodley, D.T. NC1domain of type VII collagen binds to the beta 3 chain of laminin 5 via a unique subdomain within the fibronectin-like repeats. *J. Invest.Dermatol.*, 112, 117-183, 1999

Chen, M.S., Huber, A.B., van der Haar, Frank, M., Schnell, L., Spillmann, A.A., Christ, F., and Schwab, M. Nogo-A is a myelin-associated neurite outgrowth inhibitor and an antigen for monoclonal antibody IN-1. *Nature*, 403, 434-439, 2000

Cheng, H.J., Nakamoto, M., Bergemann, A.D., and Flanagan, J.G. Complementary gradients in expression and binding of ELF-1 and Mek4 in development of the topographic retinotectal projection map. *Cell*, 82, 371-381, 1995

Cheng, H., Cao, Y., and Olson, L. Spinal cord repair in adult paraplegic rats: partial restoration of hind limb function. *Science*, 273, 510-513, 1996

Chiang, C., Litingtung, Y., Lee, E., Young, K.E., Corden, J.L., Westphal, H., and Beachy, P.A. Cyclopia and defective axial patterning in mice lacking Sonic hedgehog gene function. *Nature*, 383, 407-413, 1996

Chiasson, B.J., Tropepe, V., Morshead, C.M., and van der Kooy, D. Adult mammalian forebrain ependymal and subependymal cells demonstrate proliferative potential, but only subependymal cells have neural stem cell characteristics. *J. Neurosci.*, 19, 4462-4471, 1999

Chirgwin, J.M., Przbyla, A.E., MacDonald, R.J., and Rutter, W.J. Isolation of biologically active ribonucleic acid from sources enriched in ribonuclease. *Biochemistry*, 18, 5294-5299, 1979

Chisholm, A., and Tessier-Lavigne, M. Conservation and divergence of axon guidance mechanism. *Curr. Opin. Neurobiol.*, 9, 603-615, 1999

Chitnis, A.B. Control of neurogenesis-lessons from frogs, fish and flies. *Curr. Opin. Neurobiol.*, 9, 18-25, 1999

Choi, D.W. Calcium-mediated neurotoxicity: relationship to specific channel types and role in ischemic damage. *Trends Neurosci.*, 11, 465-469, 1988a

Choi, D.W. Glutamate neurotoxicity and diseases of the nervous system. *Neuron.*, 1, 623-634, 1988b

Choi, D. W., and Rothman, S.M. The role of glutamate neurotoxicity in hypoxic-ischemic neuronal death. *Annu. Rev. Neurosci.*, 13, 171-182, 1990

Choi-Miura, N.H., Ihara, Y., Fukuchi, K., Takeda, M., Nakano, Y., Tobe, T., and Tomita, M. SP-40, 40 is a constituent of Alzheimer's amyloid. *Acta Neuropathol.* 83, 260-264, 1992

Chu, C.H., and Tolkovsky, A.M. Alternative adrenal chromaffin cell fates induced by basic fibroblast growth factor or cyclic AMP *in vitro* depend on a collaboration with the growth substrate. *Neurosci.*, 59, 43-54, 1994

Chung, A.E., Andfhe, R., Freeman, I.L., Vergnes, J.P., Braginski, J.E., and Carlin, B. Properties of a basement membrane-related glycoprotein synthesized in culture by a mouse embryonal carcinoma-derived cell line. *Cell*, 16, 277-286, 1979

Chuong, C-M. Differential roles of multiple adhesion molecules in cell migration: Granule cell migration in cerebellum. *Experientia* 46, 892-899, 1990

Cohen, J., Burne, J.F., McKinlay, C., and Winter, J. The role of laminin and the laminin/fibronectin receptor complex in the outgrowth of retinal ganglion cell axons. *Dev. Biol.*, 122, 405, 1987

Cohen, J., Burne, J.F., Winter, J., and Bartlett, P.F. Retinal ganglion cells lose response to laminin with maturation. *Nature*, 322, 465-467, 1986

- Cohen, J.**, and Johanson, A.R. Differential effects of laminin and merosin on neurite outgrowth by developing retinal ganglion cells. *J. Cell Sci., Suppl.*, 15, 1-7, 1991
- Collin, W.**, and Donoff, R.B. Nerve regeneration through collagen tubes. *J. Dent. Res.*, 63, 987-993, 1984
- Colognato-Pyke, H.**, O'Rear, J.J., Yamada, Y., Carbonetto, S., Cheng, Y.S., and Yurchenco, P.D. Mapping of network-forming, heparan-binding, and $\alpha 1\beta 1$ integrin-recognition sites within the α -chain short arm of laminin-1. *J. Biol. Chem.*, 270, 9398-9406, 1995
- Colamarino, S.A.**, and Tessier-Lavigne, M. The axonal chemoattractant netrin-1 is also a chemorepellent for trochlear motor axons. *Cell*, 81, 621-629, 1995
- Cooper, A.R.**, and MacQueen, H.A. Subunits of laminin are differentially synthesized in mouse eggs and early embryo. *Dev. Biol.*, 96, 467-471, 1983
- Corder, E.H.**, Saunders, A.M., Strittmatter, W.J., Schechel, D.E., Gaskell, P.C., Small, G.W., Roses, A.D., Haines, J.L., and Pericak-Vance, M.A. Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science*, 261, 921-923, 1993
- Cornbrooks, C.**, Carey, D. McDonald, J., Timpl, R. and Bunge, R. In vivo and in vitro observations on laminin produced by Schwann cells. *Proc. Natl. Acad. Sci., USA* 80, 3850-3854, 1983
- Cotman, S.L.**, Halfter, W., and Cole, G.J. Identification of extracellular matrix ligands for the heparan sulfate proteoglycan agrin. *Exp. Cell Res.*, 25, 54-64, 1999
- Couchman, J.R.**, Höök, M., Rees, D.A., and Timpl, R. Adhesion, growth and matrix production by fibroblasts on laminin substrates. *J. Cell Biol.*, 96, 177-183, 1983
- Coulson, E.J.**, Barrett, G.L., Storey, E., Bartlett, P.F., Beyreuther, K., and Masters, C.L. Down-regulation of the amyloid protein precursor of Alzheimer's disease by antisense oligonucleotides reduces neuronal adhesion to specific substrata. *Brain Res.*, 770, 72-80, 1997
- Cremer, H.**, Lange, R., Christoph, A., Plomann, M., Vopper, G., Roes, J., Brown, R., Baldwin, S., Kraemer, P., Scheff, S., Barthels, D., Rajewsky, K., and Wille W. Inactivation of the N-CAM gene in mice results in size reduction of the olfactory bulb and deficits in spatial learning. *Nature*, 367, 455-459, 1994
- Dahl, D.** Isolation of neurofilament proteins and of immunologically active neurofilament degradation products from extracts of brain, spinal cord and sciatic nerve. *Biochim. Biophys. Acta*, 668, 299-306, 1981
- Dahlin, L.B.** Stimulation of regeneration of the sciatic nerve by experimentally induced inflammation in the rat. *Scan. J. Plast. Reconstr. Surg. Hand Surg.*, 26, 121-125, 1992
- Dahme, M.**, Bartsch, U., Martini, R., Anliker, B., Schachner, M., and Mantei, N. Distribution of the mouse L1 gene leads to malformation of the nervous system. *Nat. Genet.*, 17, 346-349, 1997
- Danielian, P.S.**, and McMahon, A.P. Engrailed-1 as a target of the WNT-1 signalling pathway in vertebrate midbrain development. *Nature*, 383, 332-334, 1996
- D'Arcangelo, G.**, Miao, G.G., Chen, S.C., Soares, H., Morgan, J.I., and Curran, T. A protein related to extracellular matrix proteins deleted in the mouse mutant reeler. *Nature*, 374, 719-723, 1995
- Darrow, A.L.** Rickles, R.J. and Strickland S. Maintenance and use of F9 teratocarcinoma Cells. *Methods in enzymology*, 190, 110-117, 1990
- Darmon, M.Y.** Laminin provides a better substrate than fibronectin for attachment, growth, and differentiation of 1003 embryonal carcinoma cells. *In Vitro*, 18, 997-1003, 1982
- Da Silva, C.F.**, Madison, R., Dikkes, P., Chiu, T.-H. and Sidman, R. An in vivo model

to quantify motor and sensory peripheral nerve regeneration using bioresorbable nerve guide tubes. *Brain Res.*, 342, 307-315, 1985

David, S., and Aguayo, A.J. Axonal elongation into peripheral nervous system "bridges" after central nervous system injury in adult rats. *Science*, 214, 931-933, 1981

David, S., Braun, P.E., Andckson, D.L., Kottis, V., and McKerracher, L. Laminin overrides the inhibitor effects of peripheral nervous system and central nervous system myelin-derived inhibitors of neurite growth. *J. Neurosci. Res.*, 42, 594-602, 1995

Davies, S.J., Field, P.M., and Raisman, G. Regeneration of cut adult axons fails even in the presence of continuous aligned glial pathways. *Exp. Neurol.*, 142, 203-216, 1996

Davies, S.J.A., Fitch, M.T., Memberg, S.P., Hall, A.K., Raisman, G., and Silver, J. Regeneration of adult axons in white matter tracts of the central nervous system. *Nature*, 390, 680-683, 1997

Davis, R.E., Miller, S., Herrnstadt, C., Ghosh, S.S., Fahy, E., Shinobu, L.A., Galasko, D., Thal, L.J., Beal, M.F., Howell, N., and Parker, W.D. jr. Mutations in mitochondrial cytochrome c oxidase genes segregate with late-onset Alzheimer disease. *Proc. Natl. Acad. Sci. USA*, 94, 4526-4531, 1997

Davis, G., Manthorpe, M., Engvall, E., and Varon, S. Isolation and characterization of rat schwannoma neurite-promoting factor: Evidence that the factor contains laminin. *J. Neurosci.*, 5, 2662-2671, 1985

Deacon, T., Schumacher, J., Dinsmore, J., Thomas, C., Palmer, P., Kott, S., Edge, A., Penney, D., Kassiseh, S., Dempsey, P., and Isacson, I. Histological evidence of fetal pig neural cell survival after transplantation into a patient with Parkinson's disease. *Nature Med.*, 3, 350-353, 1997

Dean, J.W., Chandrasekaran, S., and Tanzer, M.L. A biological role of the carbohydrate moieties of laminin. *J. Biol. Chem.*, 265, 12553-12562, 1990

Decherchi, P., and Gauthier, P. In vitro pre-degenerated nerve autografts support CNS axonal regeneration. *Brain Res.*, 726, 181-188, 1996

De Curtis, I., Quaranta, V., Tamura, R.N., and Reichardt, L.F. Laminin receptors in the retina: sequence analysis of the chick integrin alpha 6 subunit. Evidence for transcriptional and posttranscriptional regulation. *J. Cell Biol.*, 113, 405-416, 1991

Deiner, M.S., and Sretavan, D.W. Altered midline axon pathways and ectopic neurons in the developing hypothalamus of netrin-1- and DCC-deficient mice. *J. Neurosci.*, 19, 9900-9912, 1999

DeLeo, J.A., Coombs, D.W., Willenbring, S., Colburn, R.W., Fromm, C., Wagner, R., and Twitchell, B.B. Characterization of a neuropathic pain model: sciatic cryoneurolysis in the rat. *Pain*, 56, 9-16, 1994

Dellon, E.S., and Dellon, A.L. The first nerve graft, Vulpian, and the nineteenth century neural regeneration controversy. *J. Hand Surg.*, 18A, 369-372, 1993

Dennis, J.W., Waller, C.A., and Schirmacher, V. Identification of asparagine-linked oligosaccharides involved in tumor cell adhesion to laminin and type IV collagen. *J. Cell. Biol.*, 99, 1416-1323, 1984

Denzer, A.J., Gesemann, M., Schumacher, B., and Ruegg, M.A. An aminoterminal extension is required for the secretion of chick agrin and its binding to extracellular matrix. *J. Cell Biol.* 131, 1547-1560, 1995

Denzer, A.J., Schulthess, T., Fauser, C., Schumacher, B., Kammerer, R.A., Engel, J., and Ruegg, M.A. Electron microscopic structure of agrin and mapping of its binding site in laminin-1. *EMBO J.*, 17, 335-343, 1998

Dewitt, D.A., Silver, J., Canning, D.R., and Perry, G. Chondroitin sulfate proteoglycans are associated with the lesions of Alzheimer's disease. *Exp. Neurol.*, 121, 149-152, 1993

Dibner, M.D., and Black, I.B. Biochemical and morphological effects of testosterone

treatment on developing sympathetic neurons. *J. Neurochem.*, 30, 1479-1483, 1976

Diener, P.S., and Bregman, B.S. Neurotrophic factors prevent the death of CNS neurons after spinal cord lesions in newborn rats. *Neuroreport*, 5, 1913-1917, 1994

Doherty, P., Barton, C.H., Dickson, G., Seaton, P., Rowett, L.H., Moore, S.E., Gower, H.J., and Walsh, F.S. Neuronal process outgrowth of human sensory neurons on monolayers of cells transfected with cDNAs for five human N-CAM isoforms. *J. Cell Biol.*, 109, 789-798, 1989

Domesick, V., and Morest, D.K. Migration and differentiation of ganglion cells in the optic tectum of the chick embryo. *Neuroscience*, 2, 459-475, 1977

Dou, C-L., and Levine, J.M. Differential effects on glycosaminoglycans on neurite growth on laminin and L1 substrates. *J. Neurosci.*, 15, 8053-8066, 1995

Douville, P.J., Harvey, W.I., and Carbonetto, S. Isolation and partial characterization of high affinity laminin receptors in neural cells. *J. Biol. Chem.*, 263, 14964-14969, 1988

Drago, J., Nurcombe, V., and Bartlett, P.F. Laminin through its long arm E8 fragment promotes the proliferation and differentiation of murine neuroepithelial cells in vitro. *Exp. Cell Res.*, 192, 256-265, 1991

Drescher, U., Kremoser, C., Handwerker, C., Loschinger, J., Noda, M., and Bonhoeffer, F. In vitro guidance of retinal ganglion cell axons by RAGS, a 25 kDa tectal protein related to ligands for Eph receptor tyrosine kinases. *Cell*, 82, 359-370, 1995

Dragunow, M., Faull, R.L.M., Lawlor, P., Beilharz, E., Singleton, K., Walker, E., and Mee, E. In situ evidence for DNA fragmentation in Huntington's disease striatum and Alzheimer's disease temporal lobes. *Neuroreport*, 6, 1053-1057, 1995

Drouet, B., Pincon-Raymond, M., Chambaz, J., and Pilot, T. Laminin 1 attenuates beta-amyloid peptide Abeta(1-40) neurotoxicity of

cultured fetal rat cortical neurons. *J. Neurochem.* 73, 742-749, 1999

Durkin, M., Loechel, F., Mattei, M., Gilpin, B., Albrechtsen, R., and Wever, U. Tissue-specific expression of the human laminin alpha5-chain, and mapping of the gene to human chromosome 20g13.2-13.3 and to distal mouse chromosome 2 near the locus for the ragged (Ra) mutation. *FEBS*, 441, 296-300, 1997

Dyck, R.F., Lockwood, C.M., Kershaw, M., McHugh, N., Duance, V.C., Baltz, M.L., and Pepys, M.B. amyloid P-component is a constituent of normal human glomerular basement membrane. *J. Exp. Med.*, 152, 1162-1174, 1980

Dziadek, M. Role of laminin-nidogen complexes in basement membrane formation during embryonic development. *Experientia*, 51, 901-913, 1995

Dziadek, M., and Timpl, R. Expression of nidogen and laminin in basement membrane during mouse embryogenesis and in teratocarcinoma cells. *Dev. Biol.* 111, 372-382, 1986

Edgar, D., Timpl, R., and Thoenen, H. The heparin-binding domain of laminin is responsible for its effects on neurite outgrowth and neuronal survival. *EMBO J.*, 3, 1463-1468, 1984

Edgar, D., Timpl, R., and Thoenen, H. Structural requirements for the stimulation of neurite outgrowth by two variants of laminin and their inhibition by antibodies. *J. Cell Biol.*, 106, 1299-1306, 1988

Edgar, D. The expression and distribution of laminin in the developing nervous system. *J. Cell Sci.*, 15, 9-12, 1991

Ehrig, K., Leivo, I., Argraves, W.S., Ruoslahti, E., and Engvall. Merosin, a tissue-specific basement membrane protein, is a laminin-like protein. *Proc. Natl. Acad. Sci. USA*, 87, 3264-3268, 1990

Eikelenboom, P. and Stam, F.C. Immunoglobins and Complement factors in

senile plaques. An immunoperoxidase study. *Acta Neuropathol.*, 57, 239-242, 1982

Eisenberg, B., and Messer, A. Tonic/clonic seizures in mouse mutant carrying the weaver gene. *Neurosci. Lett.*, 96, 168-172, 1989

Ellis, D.L., and Yannas, I.V. Recent advances in tissue synthesis in vivo by use of collagen-glycosaminoglycan copolymers. *Biomaterials*, 17, 291-299, 1996

Engel, J., Odermatt, E., Engel, A., Madri, J.A., Furthmayr, Rohde, H., and Timpl, R. Shapes, domain organization and flexibility of laminin and fibronectin, two multifunctional proteins of the extracellular matrix. *J. Mol. Biol.*, 150, 79-120, 1981

Engel, J. EGF-like domains in extracellular matrix proteins: localized signals for growth and differentiation. *FEBS Lett.*, 251, 1-7, 1989

Engel, J. Domains in proteins and proteoglycans of the extracellular matrix with functions in assembly and cellular activities. *Int. J. Biol. Macromol.*, 13, 147-151, 1991

Engel, J. Structure and function of laminin. In: *Molecular aspects of basement membranes*, ed. by Rohrbach, D.H. and Timpl, R., Academic Press, San Diego, Ca, pp.359-383, 1993

Engel, J., Odermatt, E., Engel, A., Madri, J.A., Furthmayr, H., Rohde, H., and Timpl, R. Shapes, domain organizations and flexibility of laminin and fibronectin, two multifunctional proteins of the extracellular matrix. *J. Mol. Biol.*, 150, 97-120, 1981

Engelkamp, D., Rashbass, P., Seawright, A., and von Heyningen, V. Role of *Pax6* in development of cerebellar system. *Development*, 126, 3585-3596, 1999

Engvall, E., Earwicker, D., Haaparanta, T., Ruoslahti, E., and Sanes, J.R. Distribution and isolation of four laminin variants; tissue restricted distribution of heterotrimers assembled from five different subunit. *Cell Regul.*, 1, 731-740, 1990

Engvall, E., and Wewer, U.M. Domains of laminin. *J. Cell Biochem.*, 61, 493-501, 1996

Eph nomenclature committee, Unified nomenclature for Eph family receptors and their ligands. *Cell*, 90, 403-404, 1997

Ericson, J., Morton, A., Kawakami, H., Roelink, T., and Jessel, M. Two critical periods of Sonic Hedgehog signaling required for the specification of motor neuron identity *Cell*, 87, 661-673, 1996

Faissner, A. The tenascin gene family in axon growth and guidance. *Cell Tissue Res.*, 290. 331-341, 1997

Faissner, A., and Kruse, J. J1/Tenascin is a repulsive substrate for central nervous system neurons. *Neuron*, 5, 627-637, 1990

Favor, J., Sandulache, R., Neuhauser-Klaus, A., Pretsch, W., Chatterjee, B., Senft, E., Wurst, W., Blanquet, V., Grimes, P., Sporle, R., and Schughart, K. The mouse Pax2 (1Neu) mutation is identical to a human PAX2 mutation in a family with renal-coloboma syndrome and results in developmental defects of the brain, ear, eye, and kidney. *Proc. Natl. Acad. Sci. U.S.A.*, 93, 13870-13875, 1996

Fazeli, A., Dickinson, S.L., Hermiston, M.L., Tighe, R.V., Steen, R.G., Small, C.G., Stoeckli, E.T., Keino-Masu, K., Masu, M., Rayburn, H., Simons, J., Bronson, R.T., Gordon, J.I., Tessier-Lavigne, M., and Weinberg, R.A. Phenotype of mice lacking functional *Deleted in colorectal cancer* (Dcc) gene. *Nature*, 386, 24, 796-804, 1997

Feng, Z., Chang, R.C.C., Bing, G., Hudson, P., Tiao, N., Jin, L., Hong, J-S. Long-term increase of Sp-1 transcription factors in the hippocampus after kainic acid treatment. *Mol. Brain Res.*, 69, 144-148, 1999

Fenton, H., Finch, P.W., Rubin, J.S., Rosenberg, J.M., Taylor, W.G., Kuo-Leblanc, V., Rodriguez-Wolf, M., Baird, A., Schipper, H.M., and Stoppa, E.G. Hepatocyte growth factor (HGF/SF) in Alzheimer's disease. *Brain Res.*, 779, 262-270, 1998

- Fessler, L. I., Campbell, A.G., Duncan, K.G., Fessler, J.H.** Drosophila laminin: characterization and localization. *J. Cell Biol.*, 105, 2383-2391, 1987
- Figlewicz, D.A., Quarles, R.H., Johnson, D., Barbarash, G.R., Sternberger, N.H.** Biochemical demonstration of the myelin-associated glycoprotein in the peripheral nervous system. *J. Neurochem.*, 37, 749-758, 1981
- Flanagan, J.G., and Vanderhaeghen, P.** The ephrins and Eph receptors in neural development. *Annu Rev. Neurosci.*, 21, 309-345, 1998
- Forsberg, E., Ek, B., Engstrom, A., and Johansson, S.** Purification and characterization of integrin alpha 9 beta 1. *Exp. Cell Res.* 213, 183-190, 1994
- Fox, J.W., Mayer, R., Nischt, M., Aumailley, D., Reinhardt, H., Wiedenmann, K. Mann, K., Timpl, R., Krieg, and Engel, J.** Recombinant nidogen consist of three globular domains and mediates binding of laminin to collagen type IV. *EMBO J.* 10, 3137-3146, 1991
- Frade, J.M., Martinez-Morales, J.R., and Rodriguez-Tebar, A.** Laminin-1 selectively stimulates neuron generation from cultured retinal neuroepithelial cells. *Exp. Cell Res.*, 222, 140-149, 1996
- Francis, F., Koulakoff, A., Boucher, D., Chafey, P., Schaar, B., Vinet, M.-C., Friocourt, G., McDonnell, N., Reiner, O., Kahn, A., McConnell, S.K., Berwald-Netter, Y., Denoulet, P., and Chelly, J.** Doublecortin is a developmentally regulated, microtubule-associated protein expressed in migrating and differentiating neurons. *Neuron*, 23, 247-256, 1999
- Frazier, W.** Trombospondin: a modular adhesive glycoprotein of platelets and nucleated cells. *J. Cell Biol.*, 105, 625-632, 1987
- Friedman, G.C., and Seed, N.W.** Tissue plasminogen activator mRNA expression in granule neurons coincides with their migration in the developing cerebellum. *J. Comp. Neurol.*, 360, 658-670, 1995
- Friesen, J., Yates, P.A., McLaughlin, T., Friedman, G.C., O'Leary, D.D., and Barbacid, M.** Ephrin-A5 (AL-1/RAGS) is essential for proper retinal axon guidance and topographic mapping in the mammalian visual system. *Neuron*, 20, 235-243, 1998
- Frotscher, M.** Dual role of Cajal-Retzius cells and reelin in cortical development. *Cell Tissue Res.*, 290, 315-322, 1997
- Fujii, H., Nishikawa, N., Komazawa, H., Orikasa, A., Ono, M., Itoh, I., Murata, J., Azuma, I., and Saiki, I.** Inhibition of tumor invasion and metastasis by peptidic mimetics of Arg-Gly-Asp (RGD) derived from the cell recognition site of fibronectin. *Oncol. Res.*, 8, 333-342, 1996
- Galileo, D.S., Majors, J., Horwitz, A.F., and Sanes, J.R.** Retrovirally introduced antisense integrin RNA inhibits neuroblast migration in vivo. *Neuron*, 9, 1117-1131, 1992
- Galliano, M-F, Aberdam, D., Aguzzi, A., Ortonne, J-P., and Meneguzzi, G.** Cloning and complete primary structure of the mouse laminin $\alpha 3$ chain: Distinct expression pattern of the laminin $\alpha 3A$ and $\alpha 3B$ isoforms. *J. Biol. Chem.*, 268, 21820-21826, 1993
- Gao, W.-Q., and Hatten, M.E.** Neuronal differentiation rescued by implantation of weaver granule cell precursors into wild-type cerebellar cortex. *Science*, 260, 367-369, 1993
- Gao, W.Q., Liu, X-L, and Hatten, M.E.** The weaver gene encodes a nonautonomous signal for CNS neuronal differentiation. *Cell*, 68, 841-854, 1992
- Garcia-Alonso, L., Fetter, R.D., and Goodman, C.S.** Genetic analysis of *Laminin A* in *Drosophila*: extracellular matrix containing laminin A is required for ocellar axon pathfinding. *Development (Camb)*, 122, 2611-2621, 1996
- Garwood, J., Schnadelbach, O., Clement, A., Schutte, K., Bach, A., and Faissner, A.** DSD—proteoglycan is the mouse homolog of phosphacan and displays opposing effects on

neurite outgrowth dependent on neuronal lineage. *J. Neurosci.*, 19, 3888-3899, 1999

Geddes, J.W., Anderson, K.J., and Cotman, C.W. Senile plaques and aberrant-sprout stimulating structures. *Exp. Neurol.* 94, 767-776, 1986

Gee, S.H., Blacher, R.W., Douville, P.J., Provost, P.R., Yurchenco, P.D., and Carbonetto, S. Laminin-binding protein 120 from brain is closely related to the dystrophin-associated glycoprotein, dystroglycan, and binds with high affinity to the major heparin binding domain of laminin. *J. Biol. Chem.*, 268, 14972-14980, 1993

Gehlsen, K.R., Sriramaraio, P., Furcht, L.T., and Skubitz, A.P. A synthetic peptide derived from the carboxy terminus of the laminin A chain represents a binding site for the alpha 3 beta 1 integrin. *J. Cell Biol.*, 117, 449-459, 1992

Gegelashvili, G., Andersson, A-M., Schousboe, A. and Bock, E. Characterization of N-CAM diversity in cultured neurons. *FEBS Lett.* 324, 337-340, 1993

Gerecke, D.R., Wagman, D.W., Champlaud, M.F., and Burgeson, R.E. The complete primary structure for a novel laminin chain, the laminin-B1K chain. *J. Biol. Chem.*, 269, 11073-11080, 1994

Giger, R.J., Cloutier, J.-F., Sahay, A., Prinjha, R.K., Levengood, D.V., Moore, S.E., Pickering, S., Simmons, D., Rastan, S., Walsh, F.S., Kolodkin, A.L., Ginty, D.D., and Geppert, M. Neuropilin-2 is required in vivo for selective axon guidance responses to secreted semaphorins. *Neuron*, 25, 29-41, 2000

Giménez, A., Hostench, J., Stamatoglou, S.C., and Enrich, C. Differential expression of A and B laminin chains during rat liver regeneration. *Hepatology*, 22, 1259-1262, 1995

Giulian, D., and Lachman, L.B. Interleukin-1 stimulation of astroglial proliferation after brain injury. *Science*, 228, 497-499, 1985

Gleeson, J.G., Lin, P.T., Flanagan, L.A., and Walsh, C.A. Doublecortin is a microtubule-associated protein and is expressed widely by migrating neurons. *Neuron*, 23, 257-271, 1999

Gloor, S., Antonicek, H., Sweadner, K.J., Pagliusi, S., Frank, R., Moss, M., and Schachner, M. The adhesion molecule on glia (AMOG) is a homologue of the beta subunit of the Na,K-ATPase. *J. Cell Biol.*, 110, 165-174, 1990

Godsave, S.F., and Slack, J.M.W. Clonal analysis of mesoderm induction in *Xenopus Laevis*. *Dev. Biol.*, 134, 486-490, 1989

Goke, M., Zuk, A., and Podolsky, D.K. Regulation and function of extracellular matrix in intestinal epithelial restitution in vitro. *Am. J. Physiol.*, 271, 729-740, 1996

Goldberg, D.A., Isolation and partial characterization of the *Drosophila* alcohol dehydrogenase gene. *Proc. Natl. Acad. Sci. USA.* 77, 5794-5798, 1980

Goldowitz, D. The weaver granulo-prival phenotype is due to intrinsic action of the mutant locus in granule cells: evidence from homozygous weaver chimeras. *Neuron*, 2, 1565-1575, 1989

Goldowitz, D., and Hamre, K. The cells and molecules that make a cerebellum. *TINS*, 21, 375-382, 1998

Goldfinger, L.E., Hopkinson, S.B., deHart, G.W., Collawn, S., Couchman, J.R., and Jones, J.C.R. The $\alpha 3$ laminin subunit, $\alpha 6 \beta 4$ and $\alpha 3 \beta 1$ integrin coordinately regulate wound healing in cultured epithelial cells and in the skin. *J. Cell Sci.*, 112, 2615-2629, 1999

Graf, J., Iwamoto, Y., Sasaki, M. Martin, G., Kleinman, H. Robey, F. and Yamada, Y. Identification of an amino acid sequence in laminin mediating cell attachment, chemotaxis and receptor binding. *Cell* 48 989-996, 1987

GrandePré, T., Nakamura, F., Vartanian, T., and Strittmatter, S.M. Identification of the Nogo inhibitor of axon regeneration as a

Reticulon protein. *Nature*, 403, 439-444, 2000

Grant, D.S., Tashiro, K.I., Segui-Real, B., Yamada, Y., Martin, G.R., and Kleinman, H.K. Two different laminin domains mediate the differentiation of human endothelial cells into capillary-like structures in vitro. *Cell*, 58, 933-943, 1989

Graziadei, P., and Monti Graziadei, G.A. The olfactory system: A model for the study of neurogenesis and axon regeneration in mammals. In: *Neuronal Plasticity*, C.W. Cotman, ed. Raven Press, New York, pp. 131-153, 1978

Greensmith, L., Mooney, E.C., Waters, H.J., Houlihan-Burne, D.G., Lowrie, M.B. Magnesium ions reduce motoneuron death following nerve injury or exposure to N-methyl-D-aspartate in the developing rat. *Neuroscience*, 68, 807-812, 1995

Grenningloh, G., Rehm, E.J., and Goodman, C.S. Genetic analyses of growth cone guidance in drosophila fasciclin II functions as a neuronal recognition molecule. *Cell*, 67, 45-57, 1991

Grierson, J.P., Petroski, R.E., Ling, D.S.F., and Geller, H.M. Astrocytes topography and tenascin/ cytotoactin expression: correlation with the ability to support neurite outgrowth. *Dev. Brain Res.*, 55, 11-19, 1990

Grumet, M., Friedlander, D., and Edelman, G. Evidence for the binding of Ng-CAM to laminin. *Cell Adhes. Commun.* 1, 177-190, 1993

Grumet, M., Friedlander, D., and Sakurai, T. Functions of brain chondroitin sulfate proteoglycans during developments: interactions with adhesion molecules. *Perspec. Dev. Neurobiol.*, 3, 319-330, 1996

Guinec, N., Dalet-Fumeron, V., and Pagano, M. "In vitro" study of basement membrane degradation by the cysteine proteinases, cathepsins B, B-like and L. Digestion of collagen IV, laminin, fibronectin, and release of gelatinase activities from basement membrane fibronectin. *Biological Chemistry Hoppe-Seyler*, 374, 1135-1146, 1993

Gullberg, D., Tiger, C.-F., and Velling, T. Laminins during muscle development and in muscular dystrophies. *Cell Mol. Life Sci.*, 56, 442-460, 1999

Gundersen, R.W. Response of sensory neurites and growth cones to patterned substrata of laminin and fibronectin in vitro. *Dev. Biol.*, 121, 423-431, 1987

Gökhan, S., Song, Q., and Mehler, M.F. Generation and regulation of developing immortalized neural cell lines. *Methods*, 16, 345-358, 1998

Hadley, M.A., Weeks, B.S., Kleinman, H.K., and Dym, M. Laminin promotes formation of cord-like structures by cells in vitro. *Dev. Biol.*, 140, 318-327, 1990

Hager, G., Dodt, H.U., Zieglgänsberger, W., and Liesi, P. A novel form of neuronal movement in the rat cerebellum. *J. Neurosci. Res.*, 40, 207-219, 1995

Hager, G., Pawelzik, H., Kreutzberg, G.W., and Zieglgänsberger, W. A peptide derived from a neurite outgrowth-promoting domain of the $\gamma 1$ chain of laminin modulates the electrical properties of neocortical neurons. *Neurosci.*, 86, 1145-1154, 1998

Hagg, T., Portera-Cailliau, C., Jucker, M., and Engvall, E. Laminins of the adult mammalian CNS; laminin-alpha2 (merosin M-) chain immunoreactivity is associated with neuronal processes. *Brain Res.*, 764, 17-27, 1997

Hall, D.E., Neugebauer, K.M., and Reichardt, L.F. Embryonic neural retina cell response to extracellular matrix proteins: developmental changes and effects of the cell substratum attachment antibody CSAT. *J. Cell Biol.*, 104, 623-634, 1987

Hall, H., Carbonetto, S., and Schachner, M. LI/HNK-1 carbohydrate- and beta 1 integrin-dependent neural cell adhesion to laminin-1. *J. Neurochem.*, 68, 544-553, 1997a

Hall, H., Deutzmann, R., Timpl, R., Vaughan, L., Schmitz, B., and Schachner, M. HNK-1 carbohydrate-mediated cell adhesion

to laminin-1 is different from heparin-mediated and sulfatide-mediated cell adhesion. *European J. Biochem.*, 246, 233-242, 1997b

Hamelin, M., Zhou, Y., Su, M.W., Scott, I.M., and Culotti, J.G. Expression of the UNC-5 guidance receptor in the touch neurons of *C. elegans* steers their axons dorsally. *Nature*, 364, 327-330, 1993

Hamill, O.P., Marty, A., Neher, E., Sakmann, B., and Sigworth, F.J. Improved patch-clamp techniques for high-resolution current recording from cells and cell-free membrane patches. *Pflügers Arch.*, 391, 85-100, 1981

Harrelson, A.L., and Goodman, C.S. Growth cone guidance in insects: Fasciclin II is a member of the immunoglobulin superfamily. *Science*, 242, 700-708, 1988

Harrison, S.M., and Roffler-Tarlov, S. Male-sterile phenotype of neurological mouse mutant weaver. *Dev. Dyn.*, 200, 26-38, 1994

Harrison, S.M., and Roffler-Tarlov, S.K. Cell death during development of testis and cerebellum in the mutant mouse weaver. *Dev. Biol.*, 195, 174-186, 1998

Hatten, M.E., Furie, M.B., and Rifkin, D.B. Binding of developing mouse cerebellar cells to fibronectin: A possible mechanism for the formation of the external granule layer. *J. Neurosci.*, 2, 1195-1206, 1982

Hatten, M.E., and Mason, C.A. Mechanisms of glial-guided neuronal migration in vitro and in vivo. *Experientia*, 46, 907-916, 1990

Hawkins, R.L., Fan, J., and Hille, M.B. Gastrulation in the sea urchin, *Strongylocentrotus purpuratus*, is disrupted by the small laminin peptides YIGSR and IKVAV. *Cell Adhesion & Communication*, 3, 163-177, 1995

Hayashi, M., Schlesinger, D.H., Kennedy, D.W., and Yamada, K.M. Isolation and characterization of a heparin-binding domain of cellular fibronectin. *J. Biol. Chem.*, 255, 10017-10020, 1980

Hayashi, Y.K., Ishihara, T., Domen, K., Hori, H., and Arahata, K. A benign allel form of laminin $\alpha 2$ chain deficient muscular dystrophy. *Lancet*, 349, 1147, 1997

Hazarika, E.Z. Oxidised regenerated cellulose: an effective emergency haemostatic in burns surgery. *Br. J. Plast. Surg.* 38, 419-421, 1985

Heaton, M.B., and Swanson, D.J. The influence of laminin on the initial differentiation of cultured neural tube neurons. *J. Neurosci. Res.* 19, 212-218, 1988

Heck, L.W., Blackburn, W.D., Irwin, M.H., and Abrahamson, D.R. Degradation of basement membrane laminin by human neutrophil elastase and cathepsin G. *Am. J. Pathol.*, 136, 1267-1274, 1990

Hedgecock, E.M., Culotti, J.G., and Hall, D.H. The *unc-5*, *unc-6*, and *unc-40* genes guide circumferential migrations of pioneer axons and mesodermal cells on the epidermis in *C. elegans*. *Neuron*, 4, 61-85, 1990

Helbling-Leclerc, A., Zhang, X., Topaloglu, H., Cruaud, C., Tesson, F., Weissenbach, J., Tome, F.M., Schwartz, K., Fardeau, M., Tryggvason, K., and Guicheney, P. Mutations in the laminin $\alpha 2$ -chain gene (LAMA2) cause merosin-deficient congenital muscular dystrophy. *Nat. Genet.*, 11, 216-218, 1995

Hemler, M.E. VLA proteins in the integrin family: Structures, functions, and their role on leukocytes. *Annu. Rev. Immunol.*, 8, 365-400, 1990

Hemler, M.E. Structure and functions of VLA proteins and related integrands. In: Mecham, R.P., McDonald, J.A., eds. Receptors for extracellular matrix proteins (A volume of Biology of Extracellular Matrix). San Diego, CA: Academic Press, INC, 225-299, 1991

Hemler, M.E. Dystroglycan versatility. *Cell*, 97, 543-546, 1999

Hemmati-Brivanlou, A., and Melton, D. Vertebrate embryonic cells will become nerve cells unless told otherwise. *Cell*, 88, 13-17, 1997

Hemmati-Brivanlou, A., Kelly, O.G., and Melton, D.A. Follistatin, an antagonist of activin, is expressed in the Spemann organizer and displays direct neuralizing activity. *Cell*, 77, 283-295, 1994

Henchcliffe, C. Garcia-Alonso, L., Tang, J., and Goodman, C.S. Genetic analysis of laminin A reveals diverse functions during morphogenesis in *Drosophila*. *Development (Camb.)*, 118, 325-337, 1993

Henkemeyer, M., Orioli, D., Henderson, J. Saxton, T., Roder, J., Pawson, T., and Klein, R. Nuk controls pathfinding of commissural axons in the mammalian central nervous systems. *Cell*, 86, 35-46, 1996

Henrique, D., Hirsinger, E., Adam, J., Le Roux, I., Pourquite, O., Ish-Horowicz, D., and Lewis, J. Maintenance of neuroepithelial progenitor cells by Delta-Notch signalling in the embryonic chick retina. *Curr. Biol.*, 7, 661-670, 1997

Herdreen, J.C., Portera-Cailliau, C., Price, D.L., and Koliatsos, V.E. Patterns of TUNEL labeling and non-random DNA fragmentation in Huntington Disease. *Soc. Neurosci. Abstr.*, 20, 114.3, 1994

Hillery, C.A., Du, M.C., Montgomery, R.R., and Scott, J.P. Increased adhesion of erythrocytes to components of the extracellular matrix: isolation and characterization of a red blood cell lipid that binds thrombospondin and laminin. *Blood*, 87, 4879-4886, 1996

Hirose, H., Kitaguchi, T., and Tabira, T. Neurite promoting activity of collagens on embryonic neurons: Decreased effect at the postnatal stage. *Tohoku J. Exp. Med.*, 170, 207-218, 1993

Hirotsune, S., Fleck, M.W., Gambello, M.J., Bix, G.J., Chen, A., Clark, G.D., Ledbetter, D.H., McBain, C.J., and Wynshaw-Boris, A. Graded reduction of Pafah1b1 (Lis1) activity results in neuronal migration defects and early embryonic lethality. *Nat. Genet.*, 19, 333-339, 1998

Hohenester, E., Mauer, P., Hohenadl, C., Timpl, R., Andnsonius, J.N., and Engel, J. Structure of a novel Ca^{2+} -binding module in BM-40. *Nat. Struct. Biol.*, 3, 67-73, 1996

Holm, K., and Isacson, O. Factors intrinsic to the neuron can induce and maintain its ability to promote axonal outgrowth: a role for BCL2? *Trends Neurosci.*, 22, 269-273, 1999

Hopkins, J.M., Ford-Holevinski, T.S., McCoy, J.P. and Agranoff B.W. Laminin and optic nerve regeneration in the goldfish. *J Neurosci* 5, 3030-3038, 1985

Howard, J., and Pilkington, G.J. Antibodies to fibronectin bind to plaques and other structures in Alzheimer's disease and control brains. *Neurosci. Lett.*, 118, 71-76, 1990

Huang, D.Y., Weisgraber, K.H., Strittmatter, W.J., and Matthew, W.D. Interaction of apolipoprotein E with laminin increase neuronal adhesion and alter neurite morphology. *Exp. Neur.* 136, 251-257, 1995

Hunter, D.D., and Brunken, W.J. $\beta 2$ laminins modulate neuronal phenotype in the rat retina. *Mol. Cell Neurosci.*, 1997

Hunter, D.D., Cashman, N., Morris-Valero, R., Bullock, J.W., Adams, S.P. and Sanes, J.R. An LRE (leucine-arginine-glutamate)-dependent mechanism for adhesion of neurons to s-laminin. *J. Neurosci.*, 11, 3960-3971, 1991

Hunter, D., Linas, R., Ard, M., Merlie, J., and Sanes, J. Expression of s-laminin and laminin in the developing rat central nervous system. *J. Comp. Neurol.*, 323, 238-251, 1992a

Hunter, D.D., Murphy, M.D., Olsson, C.V., and Brunken, W.J. S-laminin expression in adult and developing retinae: A Potential cue for photoreceptor morphogenesis. *Neuron*, 8, 399-413, 1992b

Hunter, D.D., Porter, B.E., Bullock, J.W., Adams, S.P., Merlie, J.P., and Sanes, J.R. Primary sequence of a motor neuron-selective adhesive site in the synaptic basal lamina protein S-laminin. *Cell*, 59, 905-913, 1989a

- Hunter, D.D., Shah, V., Merlie, J.P., and Sanes, J.R.** A laminin-like adhesive protein concentrated in the synaptic cleft of the neuromuscular junction. *Nature*, 338, 229-234, 1989b
- Husmann, K., Carbonetto, S., and Schachner, M.** Distinct sites on tenascin-C mediate repellent or adhesive interactions with different neuronal cell types. *Cell Adhes. Commun.* 3, 293-310, 1995
- Husmann, K., Faissner, A., and Schachner, M.** Tenascin promotes cerebellar granule cell migration and neurite outgrowth by different domains in the fibronectin type III repeats. *J. Cell Biol.*, 116, 1475-1486, 1992
- Hutchins, J.B., and Barger, S.W.** Why neurons die: cell death in the nervous system. *Anat. Rec.*, 253, 79-90, 1998
- Hynes, R.** Migration of neuroblast along preexisting axonal tracts during prenatal cerebellar development. *J. Neurosci.*, 6, 867-876, 1986
- Ide, C., Tohyama, K., Yokota, R., Nitatori, T., and Onodera, S.** Schwann cell basal lamina and nerve regeneration. *Brain Res.*, 288, 61-75, 1983
- Iivanainen, A., Morita, T., and Tryggvason, K.** Molecular cloning and tissue-specific expression of a novel murine laminin $\gamma 3$ chain. *J. Biol. Chem.*, 274, 14107-14111, 1999
- Iivanainen, A., Sainio, K., Sariola, H., and Tryggvason, K.** Primary structure and expression of a novel human laminin $\alpha 4$ chain. *FEBS Lett.*, 365, 183-188, 1995
- Iivanainen, A., Vuolteenaho, R., Sainio, K., Eddy, R., Shows, T.B., Sariola, H., and Tryggvason, K.** The human laminin $\beta 2$ chain (S-laminin): Structure, expression in fetal tissues and chromosomal assignment of the LAMB2 gene. *Matrix Biol.*, 14, 489-497, 1994
- Iozzo, R., Cohen, I.R., Grässel, S., and Murdoch, A.D.** The biology of perlecan: the multifaceted heparan sulphate proteoglycan of basement membranes and pericellular matrices. *Biochem. J.*, 302, 625-639, 1994
- Ishii, N., Wadsworth, W.G., Stern, B.D., Culotti, J.G., and Hedgecock, E.M.** UNC-6, a laminin-related protein, guides cell and pioneer axon migrations in *C. elegans*. *Neuron*, 9, 873-881, 1992
- Iwamoto, Y., Nomizu, M., Yamada, Y., Ito, Y., Tanaka, K., and Sugioka, Y.** Inhibition of angiogenesis, tumor growth and experimental metastasis of human fibrosarcoma cells HT1080 by a multimeric form of the laminin sequence Tyr-Ile-Gly-Ser-Arg (YIGSR). *British J. Cancer*, 73, 589-595, 1996
- Iwamoto, Y., Robey, F.A., Graf, J., Sasaki, M., Kleinman, H.K., Yamada, Y., and Martin, G.R.** YIGSR a synthetic laminin pentapeptide, inhibits experimental metastasis formation. *Science* 238, 1132-1134, 1987
- Jacobson, M.** Developmental neurobiology. Third edition, Plenum Press, New York, 1991
- Jensen, P., Surmeier, D.J., and Goldowitz, D.** Rescue of cerebellar granule cells from death in weaver NR1 double mutants. *J. Neurosci.*, 19, 7991-7998, 1999
- Johansson, C.B., Momma, S., Clarke, D.L., Risling, M., Lendahl, U., and Frisén, J.** Identification of a neural stem cell in the adult mammalian central nervous system. *Cell*, 96, 25-34, 1999
- Johansson, S., Kjellen, I., Höök, M., and Timpl, R.** Substrate adhesion of rat hepatocytes: a comparison of laminin and fibronectin as attachment protein. *J. Cell Biol.*, 90, 260-264, 1981
- Johe, K.K., Hazel, T.G., Muller, T., Dugich-Djordjevic, M.M., and McKay, R.D.G.** Single factors direct the differentiation of stem cells from the fetal and adult central nervous systems. *Gen.Dev.*, 10, 3129-3140, 1996
- Jucker, M., Tian, M., Norton, D.D., Sherman, C., and Kusiak, J.W.** Laminin $\alpha 2$ is a component of brain capillary basement membrane: reduced expression in dystrophic *dy* mice. *Neuroscience*, 71, 1153-1161, 1996

- Julien**, J-P., Ramachandran, K., and Grosfeld, F. Cloning of the cDNA encoding the smallest neurofilament protein. *Biochim. Biophys. Acta*, 825, 398-404, 1985
- Kadoya**, Y., Nomizu, M., Sorkin, L.M., Yamashina, S., and Yamada, Y. Laminin alpha 1 chain G domain peptide, RKRLQVQLSIRT, inhibits epithelial branching morphogenesis of cultured embryonic mouse submandibular gland. *Dev. Dyn.*, 212, 394-402, 1998
- Kainulainen**, T., Häkkinen, L., Hamidi, S., Larandva, K., Kallioinen, M., Peltonen, J., Salo, T., Larandva, H., and Oikarainen, A. Laminin-5 expression is independent of the injury and the microenvironment during reepithelialization of wounds. *J. Histochem. Cytochem.*, 46, 353-360, 1998
- Kalaria**, R.N., and Pax, A.B. Increased collagen content of cerebral microvessels in Alzheimer's disease. *Brain. Res.*, 705, 349-352, 1995
- Kalaria**, R.N., and Perry, G. Amyloid P component and other acute-phase proteins associated with cerebellar A beta-deposits in Alzheimer's disease. *Brain Res.*, 631, 151-155, 1993
- Kalderon**, N. Role of the plasmin-generating system in the developing nervous tissue: I. Proteolysis as a mitogenic signal for the glial cells. *J. Neurosci. Res.*, 8, 509-519, 1982
- Kallunki**, T., Ikonen, J., Chow, L.T., Kallunki, P., and Tryggvason, K. Structure of the human laminin B2 chain gene reveals extensive divergence from the laminin B1 chain gene. *J. Biol. Chem.*, 266, 221-228, 1991
- Kallunki**, P., Sainio, K., Eddy, R., Byers, M., Kallunki, T., Sariola, H., Beck, K., Hirvonen, H., Shows, T.B., and Tryggvason, K. A truncated laminin chain homologous to the B2 chain: structure, spatial expression, and chromosomal assignment. *J. Cell. Biol.*, 1992
- Kanemoto**, T., Reich, R., Royce, L., Greatorex, D., Adler, S.H. Shiraishi, N., Martin, G.R., Yamada, Y., and Kleinman, H.K. Identification of amino acid sequence from the laminin A chain that stimulates metastasis and collagenase IV production. *Proc. Natl. Acad. Sci. U.S.A.* 87, 2279-2283, 1990
- Kang**, J., Leimaire, H., Unterbeck, A., Salbaum, J.M., Masters, C.L., Grzeschik, K., Multhaup, G., Beyreuther, K., and Muller-Hill, B. The precursor of Alzheimer's disease amyloid A4 protein resembles a cell-surface receptor. *Nature* 325, 733-736, 1987
- Kato**, S., Otsu, K. Ohtake, K., Kimura, Y., Yashiro, T., Suzuki, T., and Akamatsu, N. Concurrent changes in sinusoidal expression of laminin and affinity of hepatocytes to laminin during liver regeneration. *Exp. Cell Res.*, 198, 59-68, 1992
- Kauppila**, T. Nerve resuturation decrease autotomy. *Exp. Neurol.*, 124, 106-110, 1994
- Kauppila**, T., Jyväsjärvi, E., Huopaniemi, T. Huandnen, E. and Liesi, P. A laminin graft replaces neuroorrhaphy in the restorative surgery of the rat sciatic nerve. *Exp. Neur.*, 123, 181-191, 1993
- Kauppila**, T., and Pertovaara, A. Effects of different sensory and behavioural manipulations on autotomy caused by sciatic lesion in rats. *Exp. Neur.*, 111, 128-130, 1991
- Kennedy**, T.E., Serafini, T., de la Torre, J.R., and Tessier-Lavigne, M. Netrins are diffusible chemotropic factors for commissural axons in the embryonic spinal cord. *Cell*, 78, 425-435, 1994
- Kerr**, J.F.R., Wyllie, A.H., and Currie, A.R. Apoptosis: a basic biological phenomenon with wide ranging implications in tissue kinetics. *Br. J. Cancer*, 26, 239-257, 1972
- Kibbey**, M.C. Role of the SIKVAV site of laminin in promotion of angiogenesis and tumor growth: an in vivo matrigel model. *J. Natl. Cancer Inst.* 84, 1633-1638, 1992
- Kibbey**, M.C., Jucker, M., Weeks, B.S., Neve, R.L., Van Nostrand, W.E. and Kleinman, H.K. The β -amyloid precursor protein binds to the neurite-promoting

- IKVAV sequence of laminin. *Proc. Natl. Acad. Sci. U.S.A.* 90, 10150-10153, 1993
- Kim, C.J., Kim, H.O., Choe, Y.J., Lee, Y.A., and Kim, C.W.** Bcl-2 expression in neuroblastoma is differentially regulated by differentiation inducers. *Anticancer Res.*, 15, 1997-200, 1995
- Kim, D.H., Connolly, S.E., Kline, D.G., Voorhies, R.M., Smith, A., Powell, M., Yoes, T., and Daniloff, J.K.** Labeled Schwann cell transplants versus sural nerve grafts in nerve repair. *J. Neurosurg.*, 80, 254-260, 1994
- Kim, W.H., Nomizu, M., Song, S.Y., Tanaka, K., Kuratomi, Y., Kleinman, H.K., and Yamada, Y.** Laminin.alpha1-chain sequence Leu-Gln-Val-Gln-Leu-Ser-Ile-Arg (LQVQLSIR) enhances murine melanoma cell metastases. *Int. J. Cancer*, 77, 632-639, 1998
- Kitsukawa, T., Shimizu, M., Sanbo, M., Hirata, T., Tanguchi, M., Bekku, Y., Yagi, T., and Fujisawa, H.** Neurophilin-semaphorin III/D-mediated chemorepulsive signals play a crucial role in peripheral nerve projection in mice. *Neuron*, 19, 995-1005, 1997
- Klein, G., Langeegger, M., Timpl, R., and Ekblom, P.** Role of laminin A chain in the development of epithelial cell polarity. *Cell*, 55, 331-341, 1988
- Kleinman, H.K., Graft, J., Iwamoto, Y., Sasaki, M., Schasteen, C.S., Yamada, Y., Martin, G.R., and Robey, F.A.** Identification of a second active site in laminin for promotion of cell adhesion and migration and inhibition in vivo melanoma lung colonization. *Arch. Biochem. Biophys.*, 272, 39-45, 1989
- Kleinman, H.K., Weeks, B.S., Cannon, F.B., Sweeney, T.M., Sephel, G.C., Clement, B., Zain, M., Olson, M.O., Jucker, M., and Burrous, B.A.** Identification of a 110-kDa nonintegrin cell surface laminin-binding protein which recognizes an A chain neurite-promoting peptide. *Arch. Biochem. Biophys.*, 290, 320-325, 1991
- Kliot, M., Smith, G., Siegal, J., and Silver, J.** Astrocyte-polymer implants promote regeneration of dorsal root fibers into the adult mammalian spinal cord. *Exp. Neurol.*, 109, 57-69, 1990
- Knibbs, R.N., Perini, F., and Goldstein, I.J.** Structure of the major concanavalin A reactive oligosaccharides of the extracellular matrix component laminin. *Biochemistry*, 28, 6379-6392, 1989
- Kobayashi, T., Honke, K., Kuramitsu, Y., Hosokawa, M., Miyazaki, T., Murata, J., Saiki, I., Ishizuka, I., and Makita, A.** Cell-surface sulfoglycolipids are involved in the attachment of renal-cancer cells to laminin. *Int. J. Cancer*, 56, 281-285, 1994
- Koch, M., Olson, P.F., Albus, A., Jin, W., Hunter, D.D., Brunken, W.J., Burgeson, R.E., and Champlaud, M-F.** Characterization and expression of the laminin $\gamma 3$ chain: A novel, non-basement membrane-associated, laminin chain. *J. Cell Biol.*, 145, 605-617, 1999
- Koh, J-Y., Yang, L.L., and Cotman C.W.** β -Amyloid protein increases the vulnerability of cultured cortical neurons in excitotoxic damage. *Brain Res.*, 533, 315-320, 1990
- Kohfeldt, E., Sasaki, T., Gohring, W., and Timpl, R.** Nidogen-2: a new basement membrane protein with diverse binding properties. *J. Mol. Biol.*, 282, 99-109, 1998
- Komuro H., and Rakic, P.** Selective role of N-type calcium channels in neuronal migration. *Science*, 257, 806-809, 1992
- Komuro H., and Rakic, P.** Modulation of neuronal migration by NMDA receptors. *Science*, 260, 95-97, 1993
- Komuro H., and Rakic, P.** Intracellular Ca^{2+} fluctuations modulate the rate of neuronal migration. *Neuron*, 17, 275-285, 1996.
- Koo, E.H., Park, L., and Selkoe, D.J.** Amyloid β -protein as a substrate interacts with extracellular matrix to promote neurite outgrowth *Proc. Natl. Acad. Sci. U.S.A.* 90, 4748-4752, 1993
- Kopyov, O.V., Andcques, D.S., Lieberman, A., Duma, C.M., and Rogers, R.L.** Clinical study of fetal mesencephalic intracerebral

transplants for the treatment of Parkinson's disease. *Cell Transplant.*, 5, 327-337, 1996

Kouzi-Koliakos, K., Koliakos, G.G., Tsilibary, E.C., Furcht, L.T., and Charonis, A.S. Mapping of three major heparin-binding sites on laminin and identification of a novel heparin-binding site on the B1 chain. *J. Biol. Chem.*, 264, 17971-17978, 1989

Kovacs, D.M., Fausett, H.J., Page, K.J., Kim, T.-W., Moir, R.D., Merriam, D.E., Hollister, R.D., Hallmark, O.G., Mancini, R., Felsentein, K.M., Hyman, B.T., Tanzi, R., and Wasco, W. Alzheimer-associated presenilins 1 and 2. Neuronal expression in brain and localization to intracellular membranes in mammalian cells. *Nature Med.*, 2, 224-229, 1996

Kowall, N.W., Beal, M.F., Busciglio, J., Duffy, L.K., and Yankner, B.A. An in vivo model for the neurodegenerative effects of amyloid and protection by substance P. *Proc. Natl. Acad. Sci. U.S.A.*, 88, 7247-7251, 1991

Kramer, R.H., McDonald, K.A., and Vu, M.P. Human melanoma cells express a novel integrin receptor for laminin. *J. Biol. Chem.*, 264, 15642-15649, 1989

Kromer, L.F., Björklund, A., and Stenevi, U. Regeneration of the septohippocampal pathways in adult rats is promoted by utilizing embryonic hippocampal implants as bridges. *Brain Res.*, 210, 173-200, 1981

Krystosek, A. and Seeds, N. Plasminogen activator secretion by granule neurons in cultures of developing cerebellum. *Proc. Natl. Acad. Sci. U.S.A.* 78, 7810-7814, 1981

Kubo, Y. Development of ion channels and neurofilaments during neuronal differentiation of mouse embryonal carcinoma cell lines. *J. Physiol.* 409, 497-523, 1989

Kubota, S., Tashiro, K., and Yamada, Y. Signaling site of laminin with mitogenic activity. *J. Biol. Chem.*, 267, 4285-4288, 1992

Kuff, E. and Fewell, J. Induction of neural like cells and acetylcholinesterase activity in

cultures of F9 teratocarcinoma treated with retinoic acid and dibutyryl cyclic adenosine monophosphate. *Dev. Biol.* 77, 103-115, 1980

Kuratomi, Y., Nomizu, M., Nielsen, P.K., Tanaka, K., Song, S.Y., Kleinman, Y.K., and Yamada, Y. Identification of metastasis-promoting sequences in the mouse laminin. *Exp. Cell Res.*, 249, 386-395, 1999

Labrador, R.O., Buti, M., and Navarro, X. Influence of collagen and laminin gels concentration on nerve regeneration after resection and tube repair. *Exp. Neur.*, 149, 243-252, 1998

Landolt, R.M., Vaughan, L., Winterhalter, K.H., and Zimmermann, D.R. Versican is selectively expressed in embryonic tissues that act as barriers to neural crest cell migration and axon outgrowth. *Development*, 121, 2303-23-12, 1995

Langone, F., Lora, S., Veronese, F.M., Caliceti, P., Parnigotto, P.P., Valenti, F., and Palma, G. Peripheral nerve repair using a poly(organo)phosphazene tubular prosthesis. *Biomaterials*, 16, 347-353, 1995

Laurie, G.W., Bing, J.T., Kleinman, H.K., Hassell, J.R., Aumailley, M., Martin, G.R., and Feldmann R.J. Localization of binding sites for laminin, heparan sulfate proteoglycan and fibronectin on basement membrane (type IV) collagen. *J. Mol. Biol.*, 188, 1-12, 1986

Lazarov-Spiegler, O., Solomon, A.S., Zeev-Brann, A.B., Hirschberg, D.L., Lavie, V., and Schwartz, M. Transplantation of activated macrophages overcomes central nervous system regrowth failure. *Faseb J.*, 10, 1296-1302, 1996

Lazarov-Spiegler, O., Solomon, A.S., and Schwartz, M. Peripheral nerve-stimulated macrophages stimulate a peripheral nerve-like regenerative response in rat transected optic nerve. *Glia*, 24, 329-337, 1998

Le Beau, J.M., Ellisman, M.H., Powell, H.C. Ultrastructural and morphometric analysis of long-term peripheral nerve regeneration

through silicone tubes. *J. Neurocytol.*, 17, 161-172, 1988

Lee, K.L., Dietrich, P., Jessel, T.M. Genetic ablation reveals that the root plate is essential for dorsal interneuron specification. *Nature*, 403, 734-740, 2000

Lee, M.K., Tuttle, J.B.; Rebhun, L.I.; Cleveland, D.W., and Frankfurter, A. The expression and posttranslational modification of a neuron-specific β -tubulin isotype during chick embryogenesis. *Cell Motil. Cytoskeleton* 17, 118-132, 1990

Lein, P.J., Banker, G.A., and Higgins, D. Laminin selectively enhances axonal growth and accelerates the development of polarity by hippocampal neurons in culture. *Brain Res. Dev. Brain Res.*, 69, 191-197, 1992

Leivo, I., Vaheri, A., Timpl, R., and Wartiovaara, J. Appearance and distribution of collagens and laminin in the early mouse embryo. *Dev. Biol.*, 76, 100-114, 1980

Leivo, I., and Engvall, E. Merosin, a protein specific for basement membranes of Schwann cells, striated muscle, and trophoblast, is expressed late in nerve and muscle development. *Proc. Natl. Acad. Sci. USA*, 85, 1544-1548, 1988

LeMosy, E.K., Lightner, V.A., and Erickson, H.P. Structural analysis of human glial variant laminin. *Exp. Cell Res.*, 227, 80-88, 1996

Lemmon, V., Farr, K.L., and Lagenaur, C. L1-mediated axon outgrowth occurs via a homophilic binding mechanism. *Neuron*, 2, 1597-1603, 1989

Lesage, F., Duprat, F., Fink, M., Guillemare, T., Coppola, T., Lazdunski, M., and Hugnot J.-P. Cloning provides evidence for a family of inward rectifier and G-protein coupled K^+ channels in the brain. *FEBS Lett.*, 353, 37-42, 1994

Letourneau, P., Madsen, A., Palm, S. and Furcht, L. Immunoreactivity for laminin in the developing longitudinal pathway of the brain. *Dev Biol.* 125, 135-144, 1988

Levavasseur, F., Mayer, U., Guillouzo, A., and Clement, B. Influence of nidogen complexed or not with laminin on attachment, spreading, and albumin and laminin B2 mRNA levels of rat hepatocytes. *J. Cell Phys.*, 161, 257-266, 1994

Levine, J. and Flynn, P. Cell surface changes accompanying the neural differentiation of an embryonal carcinoma cell line. *J. Neurosci.* 6, 3374-3384, 1986

Levy-Lahad, E., Wasco, W., Poorkaj, P., Romano, D.M., Oshima, J., Pettingell, W.H., Yu, C., Jondro, P.D., Schmidt, S.D., Wang, K., Crowley, A.C., Fu, Y.-H., Guenette, S.Y., Galas, D., Nemens, E., Wijsman, E.M., Bird, T.D., Schellenberg, G.D., and Tanzi, R.E. Candidate gene for the chromosome 1 familial Alzheimer's disease locus. *Science*, 269, 973-977, 1995

Li, D., Field, P.M., and Raisman, G. Failure of axon regeneration in postnatal rat entorhinohippocampal slice coculture is due to maturation of the axon, not that of the pathway or target. *Eur. J. Neurosci.*, 7, 1164-1171, 1995

Li, J., Xu, M., Zhou, H., Ma, J., Potter, H. Alzheimer presenilins in the nuclear membrane, interphase kinetochores, and centrosomes suggest a role in chromosome segregation. *Cell*, 90, 917-927, 1997

Li, R., Thode, S., Zhou, J., Richard, N., Pardinas, J., Rao, M.S., and Sah, D.W. Motoneuron differentiation of immortalized human spinal cord cell lines. *J. Neurosci. Res.*, 59, 342-352, 2000

Li, Y., and Raisman, G. Schwann cells induce sprouting in motor and sensory axons in the adult rat spinal cord. *J. Neurosci.*, 14, 4050-4063, 1994

Libby, R.T., Lavalley, C.R., Balkema, G.W., Brunken, W.J., and Hunter, D.D. Disruption of laminin β 2 chain production causes alterations in morphology and function in the CNS. *J. Neurosci.*, 19, 9399-9411, 1999

Libby, R.T., Xu, Y., Selfors, L.M., Brunken, W.J., and Hunter, D.D. Identification of the cellular source of laminin β 2 in adult and

developing vertebrate retinae. *J. Comp. Neurol.*, 389, 655-667, 1997

Liem, K., Tremmi, G., Roelink, H., and Jessel, T.M. Dorsal differentiation of neural plate cells induced by BMP-mediated signals from epidermal ectoderm. *Cell*, 82, 969-979, 1995

Liesi, P. Laminin and fibronectin in normal and malignant neuroectodermal cells. *Med. Biol.*, 62, 163-180, 1984

Liesi, P. Do neurons in the vertebrate CNS migrate on laminin. *EMBO J.*, 4, 1163-1170, 1985a

Liesi, P. Laminin immunoreactive glia distinguish regenerative adult CNS systems from nonregenerating ones. *EMBO J* 4, 2505-2511, 1985b

Liesi, P. Extracellular matrix and neuronal movement. *Experientia* 46, 900-907, 1990

Liesi, P. Laminin and fibronectin and their cellular receptors during neural development. *Seminars in the Neurosci.*, 3, 331-340, 1991

Liesi, P. Neuronal migration on laminin involves neuronal contact formation followed by nuclear movement inside a preformed process. *Exp. Neurol.*, 117, 103-113, 1992

Liesi, P., Dahl, D., and Vaheri, A. Neurons cultured from developing rat brain attach and spread preferentially to laminin. *J. Neurosci. Res.*, 11, 241-251, 1984a

Liesi, P., Hager, G., Dodt, H-U., Seppälä, I. and Zieglgänsberger, W. Domain specific antibodies against a neurite outgrowth domain of the B2 chain of laminin inhibit neuronal migration in neonatal rat cerebellum. *J. Neurosci. Res.*, 1995

Liesi, P., Kaakkola, S., Dahl, D. and Vaheri, A. Laminin is induced in adult rat brain by injury. *EMBO J.*, 3, 683-686, 1984b

Liesi, P., Närviäinen, A., Soos, J., Sariola, H. and Snounou, G. Identification of a neurite outgrowth- promoting domain of laminin using synthetic peptides. *FEBS* 224, 1, 141-148, 1989

Liesi, P., Rechartdt, L. and Wartiovaara, J. Nerve growth factor induces adrenergic neuronal differentiation in F9 teratocarcinoma cells. *Nature*, 306, 265-267, 1983

Liesi, P., and Risteli, L. Glial cells of mammalian brain produce a variant form of laminin. *Exp. Neurol.* 105, 86-92, 1989.

Liesi, P., Seppälä, I., and Trenkner, E. Neuronal migration in cerebellar microculture is inhibited by antibodies against a neurite outgrowth domain of laminin. *J. Neurosci. Res.* 33, 170-176, 1992

Liesi, P., and Silver, J. Is glial laminin involved in axon guidance in the mammalian CNS ? *Dev Biol.* 130, 775-785. 1988

Liesi, P., Stewart, R.R., Akinshola, B.E., and Wright, J.M. Weaver cerebellar granule neurons show altered expression of NMDA receptor subunits both in vivo and in vitro. *J. Neurobiol.*, 38, 441-454, 1999

Liesi, P., and Wright, J. M. Weaver Granule neurons are rescued by calcium channel antagonists and antibodies against a neurite outgrowth domain of B2 chain of laminin. *J. Cell Biol.*, 134, 447-486, 1996

Liesi, P., Wright, J., and Krauthamer, V. BAPTA-AM and ethanol protect cerebellar granule neurons from the destructive effect of the weaver gene. *J. Neurosci. Res.*, 48, 571-579, 1997

Liétard, J., Musso, O., Thérêt, N., L'Helgoualc'h, A., Campion, J.P., Yamada, Y., and Clément, B. Sp1-mediated transactivation of lamC1 promoter and coordinated expression of laminin-γ1 and Sp1 in human hepatocellular carcinomas. *Am. J. Pathol.*, 151, 1663-1672, 1997

Lightner, V.A., and Erickson, H.P. Binding of hexabrachion (tenascin) to the extracellular matrix and substratum and its effect on cell adhesion. *J. Cell Sci.*, 95, 263-277, 1990

Lin, D.M., Fetter, R.D., Kopczynski, C., Grenningloh, G., and Goodman, C.S. Genetic analysis of Fasciclin II in *Drosophila*:

defasciculation, refasciculation, and altered fasciculation. *Neuron*, 13, 1055-1069, 1994

Linder, J., Rathjen, F., and Schachner, M. L1 mono- and polyclonal antibodies modify cell migration in early postnatal mouse cerebellum. *Nature*, 305, 427-430, 1986

Lindner, W., and Robey, F.A. Automated synthesis and use of N-chloroacetyl-modified peptides for the preparation of synthetic peptide polymers and peptide-protein immunogens. *Int. J. Pept. Prot. Res.*, 30, 794-800, 1987

Lindvall, O., Sawle, G., Winder, H., Rothwell, J.C., Björklund, A., Brooks, D., Brundin, P., Frackowiak, R., Marsden, C.D., Odin, P., and Rehnström, S. Evidence for long-term survival and function of dopaminergic grafts in progressive Parkinson's disease. *Ann. Neurol.*, 35, 172-180, 1994

Lin-Lee, Y.C., Kao, F.T., Cheung, P., Chan, L. Apolipoprotein E gene and expression: localization of the structural gene to human chromosome 19 and expression of ApoE mRNA in lipoprotein- and non-lipoprotein-producing tissues. *Biochemistry*, 24, 3751-3756, 1985

Liu, J., and Mayne, R. The complete cDNA coding sequence and tissue-specific expression of the mouse laminin alpha 4 chain. *Matrix Biol.*, 15, 433-437, 1996

Livesey, F.J. Netrins and netrin receptors. *CMLS*, 56, 62-68, 1999

Lochter, A., Vaughan, L., Kaplony, A., Prochiantz, A., Schachner, M., and Faissner, A. J1/ tenascin in substrate-bound and soluble form displays contrary effects on neurite outgrowth. *J. Cell Biol.*, 113, 1159-1172, 1991

Lois, C., and Alvarez-Buylla, A. Proliferative subventricular zone cells in the adult mammalian forebrain can differentiate into neurons and glia. *Proc. Natl. Acad. Sci. USA*, 90, 2074-2077, 1993

Loo, D.T., Copani, A., Pike, C.J., Whittemore, E.R., Walencewicz, A.J., and

Cotman, C.W. Apoptosis is induced by β -amyloid in culture central nervous system neurons. *Proc. Natl. Acad. Sci. U.S.A.*, 90, 7951-7955, 1993

Lu, X., and Richardson, P.M. Inflammation near the nerve cell body enhances axonal regeneration. *J. Neurosci.*, 11, 972-978, 1991

Luckenbill-Edds, L. Laminin and the mechanism of neuronal outgrowth. *Brain Res. Rev.*, 23, 1-27, 1997

Luckenbill-Edds, L., Kaiser, C.A., Rodgers, T.R., and Powell, D.D. Localization of the 110 kDa receptor for laminin in brains of embryonic and postnatal mice. *Cell Tissue Res.*, 279, 371-377, 1995

Luebke, A.E., Dickerson, I.M., and Muller, K.J. In situ hybridization reveals transient laminin B-chain expression by individual glial and muscle cells in embryonic leech central nervous systems. *J. Neurobiol.*, 27, 1-14, 1995

Lumsden, A., and Krumlauf, R. Patterning the vertebrate neuraxis. *Science*, 274, 1109-1115, 1996

Lundborg, G., Dahlin, L.B., Danielsen, N., Gelberman, R.H., Longo, F.M., Powell, H.C., and Varon, S. Nerve regeneration in silicone chambers: influence of gap length and of distal stump components. *Exp. Neurol.*, 76, 361-375, 1982

Lundborg, G., Dahlin, L.B., Dohi, D., Kanje, M., and Terada, N. A new type of "bioartificial" nerve graft for bridging extended defects in nerves. *J. Hand Surg.*, 22, 299-303, 1997

Ma, J., Yee, A., Brewer, H.B. Jr., Das, S., and Potter, H. Amyloid-associated proteins α 1-antichymotrypsin and apolipoprotein E promote assembly of Alzheimer β -protein into filaments. *Nature*, 372, 92-94, 1994

Madison, R.S., Archibald, S.J., and Krarup, C. Peripheral nerve injury. In *wound healing: biochemical and clinical aspects* (I.K. Cohen, R.F. Diegelmann, and W.J. Lindblad Eds.), pp450-487, 1992

- Madison, R.,** Da Silva, C.F., Dijkers, P., Sidman, R.L. and Chiu, T.-H. Increased rate of peripheral nerve regeneration using bioresorbable nerve guides and laminin containing gel. *Exp. Neurol.* 88, 767-772, 1985
- Madison, R.,** Da Silva, C.F., Dijkers, P., Sidman, R.L. and Chiu, T.-H. Peripheral nerve regeneration with entubulation repair: comparison of biodegradable nerve guides versus polyethylene tubes and the effects of a laminin-containing gel. *Exp. Neurol.* 95, 378-390, 1987
- Maggi, R.,** Poletti, A., Casulari, L.A., Pimpinelli, F., Piva, F., Zanisi, M.R., and Martini, L. Effects and metabolism of steroid hormones in human neuroblastoma cells. *Steroids*, 63, 257-262, 1998
- Mahley, R.W.,** and Huang, Y. Apolipoprotein E: from atherosclerosis to Alzheimer's disease and beyond. *Curr. Opin. Lipidol.* 10, 207-217, 1999
- Malinoff, H.L.,** and Wicha, M.S. Isolation of a cell surface receptor protein for laminin from murine fibrosarcoma cells. *J. Cell Biol.* 96, 1475-1479, 1983
- Mansikka, H.,** and Pertovaara, A. Influence of selective alpha 2-adrenergic agents on mustard oil-induced central hyperalgesia in rat. *Eur. J. Pharmacol.* 281, 43-48, 1995
- Manthorpe, M.,** Engvall, E., Ruoslahti, E., Longo, F.M., Davis, G.E., and Varon, S. Laminin promotes neuritic regeneration from cultured peripheral and central neurons. *J. Cell Biol.* 97, 1882-1890, 1983
- Marin, F.,** and Puelles, L. Patterning of the embryonic avian midbrain after experimental inversions: a polarizing activity from the isthmus. *Dev. Biol.* 163, 19-37, 1994
- Marinkovich, M.P.,** Lunstrum, G.P., Keene, D.R., and Burgeson, R.E. The dermal-epidermal junction of human skin contains a novel laminin variant. *J. Cell Biol.* 119, 695-703, 1992
- Maricich, S.M.,** Soha, J., Trenkner, E., and Herrup, K. Failed cell migration and death of Purkinje cells and deep nuclear neurons in the weaver cerebellum. *J. Neurosci.* 17, 3675-3683, 1997
- Mark, M.D.,** Lohrum, M., and Puschel, A.W. Patterning neuronal connections by chemorepulsion: the semaphorins. *Cell Tissue Res.* 290, 299-306, 1997
- Marti, E.,** Bumcrot, D.A., Takada, R., and McMahon, A.P. Requirement of 19K form of Sonic hedgehog for induction of distinct ventral cell types in CNS explants. *Nature*, 375, 322-323, 1995
- Martin, G.R.** Teratocarcinomas and mammalian embryogenesis. *Science* 209, 768-776, 1980
- Martin, G.R.** and Evans, M. Differentiation of clonal cell lines of teratocarcinoma cells: formation of embryonal bodies in vitro. *Proc. Natl. Acad. Sci. U.S.A.* 72, 1441-1445, 1975
- Martin, G.,** and Timpl, R. Laminin and other basement membrane components. *Ann. Rev. Cell Biol.* 3, 57-85, 1987
- Martinez-Serrano, A.,** and Björklund, A. Immortalized neural progenitor cells for CNS gene transfer and repair. *Trends Neurosci.* 20, 530-538, 1997
- Masters, C.L.,** Simms, G., Weinman, N.A., Multhapu, G., McDonald, B.L., and Beyreuther, K. Amyloid plaque core protein in Alzheimer disease and Down syndrome. *Proc. Natl. Acad. Sci. U.S.A.* 82, 4245-4249, 1985
- Matsushita, K.,** Yoshioka, H., Fushiki, S., Hirai, K., Kihara, M., Kadono, K., Goma, H., and Sawada, T. Tissue culture study on neural migration in the rat cerebral cortex: effects of low dose radiation. *Brain Dev.* 19, 452-458, 1997
- Matsuzawa, M.,** Tokumitsu, S., Knoll, W., and Liesi, P. Molecular gradient along the axon pathway is not required for directional axon growth. *J. Neurosci. Res.* 53, 114-124, 1998
- Matsuzawa, M.,** Weight, F.F., Potember, R.S., and Liesi, P. Directional neurite

outgrowth and axonal differentiation of embryonic hippocampal neurons are promoted by a neurite outgrowth domain of the B2-chain of laminin. *Internat. J. Dev. Neurosci.*, 14, 283-295, 1996

Matter, M.L., and Laurie, G.W. A novel laminin E8 cell adhesion site required for lung alveolar formation in vitro. *J. Cell Biol.*, 124, 1083-1090, 1994

Mayer, U., Nischt, R., Pöschl, E., Mann, K., Fukuda, K., Gerl, M., Yamada, Y., and Timpl, R. A single EGF-like motif of laminin is responsible for high affinity nidogen binding. *EMBO J.*, 12, 1879-1885, 1993

McBurney, M., Jones-Villeneuve, M., Edwards, M. and Anderson, P. Control of muscle and neuronal differentiation in a cultured embryonal cell line. *Nature*, 299, 165-167, 1982

McCarthy, R.A., Beck, K., and Burger, M. Laminin is structurally conserved in the sea urchin basal lamina. *EMBO J.*, 6, 1587-1593, 1987

McDonnell, J.M., Jones, G.E., White, T.K., and Tanzer, M.L. Calreticulin binding affinity for glycosylated laminin. *J. Biol. Chem.*, 271, 7891-7894, 1996

McGarvey, Baron van Evercooren, A., Kleinman, H.K., and Dubois-Dalcq, M. Synthesis and effects of basement membrane components in cultured rat Schwann cells. *Dev. Biol.*, 105, 18-28, 1984

McGrath, J.A., Pulkkinen, L., Christiano, A.M., Leigh, I.M., Eady, R.A.J., and Uitto, J. Altered laminin 5 expression due to mutations in the gene encoding the $\beta 3$ chain (LAMB3) in generalized atrophic benign epidermolysis bullosa. *J. Invest. Dermatol.*, 104, 467-474, 1995

McKeon, R.J., Höke, A., and Silver, J. Injury-induced proteoglycans inhibit the potential for laminin-mediated axon growth on astrocytic scars. *Exp. Neurol.*, 136, 32-43, 1995

McKerracher, L., Chamoux, M., and Arregui, C.O. Role of laminin and integrin

interactions in growth cone guidance. *Mol. Neurobiol.*, 12, 95-116, 1996

McKerracher, L., David, S., Andckson, D.L., Kottis, V., Dunn, R.J., and Braun, P.E. Identification of myelin-associated glycoprotein as a major myelin-derived inhibitor of neurite outgrowth. *Neuron*, 13, 805-811, 1994

McMahon, A.P., and Bradley, A. The Wnt-1 (Wint-1) proto-oncogene is required for development of a large region of the mouse brain. *Cell*, 62, 1073-1085, 1990

McMahon, A.P., Joyner, A.L., Bradley, A., and McMahon, J.A. The midbrain-hindbrain phenotype of Wnt-1/Wnt-1-mice results from stepwise deletion of engrailed-expression cells by 9.5 days postcoitum. *Cell*, 69, 581-595, 1992

Mecenas, P.E., Tsirka, S.E., Sallés, F., and Strickland, S. Removal of tissue plasminogen activator does not protect against neuronal degeneration in the cerebellum of the weaver mouse. *Brain Res.*, 772, 233-238, 1997

Mecham, R.P. Receptors for laminin. *FASEB J.*, 5, 2538-2546, 1991

Mecham, R.P., Hinek, A., Griffin, G.L., Senior, R.M., and Liotta, L.A. The elastin receptor shows structural and functional similarities to the 67-kDa tumor cell laminin receptor. *J. Biol. Chem.*, 264, 16652-16657, 1989

Menard, S., Tagliabue, E., and Colnaghi, M.I. The 67 kDa laminin receptor as a prognostic factor in human cancer. *Breast Cancer Res. Treat.*, 52, 137-145, 1998

Meyerhardt, J.A., Caca, K., Eckstrand, B.C., Hu, G., Lengauer, C., Banavali, S., Look, A.T., and Fearon, E.R. Netrin-1 interaction with deleted in colorectal cancer (DCC) and alternations in brain tumor and neuroblastomas. *Cell Growth Differ.*, 10, 35-42, 1999

Miale, I.L., and Sidman, R.L. An autoradiographic analysis of histogenesis in the mouse cerebellum. *Exp. Neurol.*, 4, 277-296, 1961

- Migheli, A., Attanasio, A., Lee, W.-H., Bayer, S.A., and Ghetti, B.** Detection of apoptosis in weaver cerebellum by electron microscopic in situ end-labeling of fragmented DNA. *Neurosci. Lett.*, 199, 53-56, 1995
- Miner, J.H., Cunningham, J., and Sanes, J.R.** Roles for laminin in embryogenesis: exencephaly, syndactyly, and placentopathy in mice lacking the laminin alpha 5 chain. *J. Cell Biol.*, 143, 1713-1723, 1998
- Miner, J.H., Lewis, R.M., and Sanes, J.R.** Molecular cloning of a novel laminin chain, alpha 5, and widespread expression in adult mouse tissues. *J. Biol. Chem.*, 270, 28523-28526, 1995
- Miner, J.H., Patton, B.L., Lentz, S.I., Gilgert, D.J., Snider, W.D., Jenkins, N.A., Copeland, N.G., and Sanes, J.R.** The laminin α chains: Expression, developmental transitions, and chromosomal locations of $\alpha 1$ -5, identification of heterotrimeric laminins 8-11, and cloning of a novel $\alpha 3$ isoform. *J. Cell. Biol.*, 3, 685-701, 1997
- Minty, A., Alonso, S., Caravatti, M., and Buckingham, M.** A fetal skeletal muscle actin mRNA in the mouse and its identity with cardiac actin mRNA. *Cell*, 30, 185-192, 1982
- Miyagoe, Y., Hanaoka, K., Nonaka, I., Hayasaka, M., Arahata, K., Nabeshima, Y., and Takeda, S.** Laminin alpha2 chain-null mutant mice by targeted disruption of the Lama2 gene: a new model of merosin (laminin 2)-deficient congenital muscular dystrophy. *FEBS Lett.*, 415, 33-39, 1997
- Mjaatved, A., Cabin, D., Cole, S., Long, L., Breitwieser, G., and Reeves, R.** Assessment of a mutation in the H5 domain of GIRK2 as a candidate for the weaver mutation. *Genome Research*, 5, 453-463, 1995
- Monji, A., Tashiro, K., Yoshida, I., Hayashi, Y., and Tashiro, N.** Laminin inhibits A beta 40 fibril formation promoted by apolipoprotein E4 in vitro. *Brain Res.*, 796, 171-175, 1998a
- Monji, A., Tashiro, K., Yoshida, I., Hayashi, Y., and Tashiro, N.** Laminin inhibits A β 42 fibril formation in vitro. *Brain Res.*, 788, 187-190, 1998b
- Monschau, B., Kremoser, C., Ohta, K., Tanaka, H., Kaneko, T., Yamada, T., Handwerker, C., Hornberger, M.R., Loschinger, J., Pasquale, E.B., Siever, D.A., Verderame, M.F., Muller, B.K., Bonhoeffer, F., and Drescher, U.** Shared and distinct functions of RAGS and ELF-1 in guiding retinal axons. *EMBO J.*, 16, 1258-1267, 1997
- Moonen, G., Grau-Wagemans, M.P., and Selak, I.** Plasminogen activator-plasmin system and neuronal migration. *Nature*, 298, 753-755, 1982
- Morest, D.K.** A study of neurogenesis in the forebrain of opossum pouch young. *Z. Anat. Entwicklungsgesch* 130, 265-305, 1970
- Morissette, N., and Carbonetto, S.** Laminin $\alpha 2$ chain (M chain) is found within the pathway of avian and murine retinal projections. *J. Neurosci.*, 15, 8067-8082, 1995
- Morrow, D.R., Campbell, G., Lieberman, A.R., and Anderson, P.N.** Differential regenerative growth of CNS axons into tibial and peroneal nerve grafts in the thalamus of adult rats. *Exp. Neurol.*, 120, 60-69, 1993
- Morshead, C.M., Reynolds, B.A., Craig, C.G., McBurney, M.W., Staines, W.A., Morassutti, D., Weiss, S., and van der Kooy, D.** Neural stem cells in the adult mammalian forebrain: A relatively quiescent subpopulation of subependymal cells. *Neuron*, 13, 1071-1082, 1994
- Mueller, B.K.** Growth cone guidance: first steps towards a deeper understanding. *Annu Rev. Neurosci.*, 22, 351-388, 1999
- Mukhopadhyay, G., Doherty, P., Walsh, F.S., Crocker, P.R., and Fill-in, M.T.** A novel role for myelin-associated glycoprotein as an inhibitor of axonal regeneration. *Neuron*, 13, 757-767, 1994
- Munz, M., Rasminsky, M., Aguayo, A.J., Vidal-Sanz, M., and Devor, M.** Functional

- activity of rat brainstem neurons regenerating axons along peripheral nerve grafts. *Brain Res.*, 340, 115-125, 1985
- Mutisya**, E.M., Bowling, A.C., and Beal, M.F. Cortical cytochrome oxidase activity is reduced in Alzheimer's disease. *J. Neurochem.*, 63, 2179-2184, 1994
- Mönning**, U. Sandbrink, R., Weidemann, A., Banati, R.B., Masters, C.L., and Beyreuther, K. Extracellular matrix influence the biogenesis of amyloid precursor protein in microglial cells. *J. Biol. Chem.* 270, 7104-7110, 1995
- Nagai**, N., Urano, T., Endo, A., Takahashi, H., Takada, Y., and Takada, A. Neuronal degeneration and a decrease in laminin-like immunoreactivity is associated with elevated tissue-type plasminogen activator in the rat hippocampus after kainic acid injection. *Neurosci. Res.*, 33, 147-154, 1999
- Nakai**, M., Mundy, G.R., Williams, P.J., Boyce, B. and Yoneda, T. A synthetic antagonist to laminin inhibits the formation of osteolytic metastases by human melanoma cells in nude mice. *Cancer Res.* 52, 5395-5399, 1992
- Nakatsuji**, N., and Nagata, I. Paradoxical perpendicular contact guidance displayed by mouse cerebellar granule neurons in vitro. *Development*, 106, 441-447, 1989
- Namba**, Y., Tomonga, M., Kawasaki, H., Otomo, E., and Ikeda, K. Apolipoprotein E immunoreactivity in cerebral amyloid deposits and neurofibrillary tangles in Alzheimer's disease and kuru plaque amyloid in Creutzfeld-Jakob disease. *Brain Res.* 541, 163-166, 1991
- Narakas**, A. The use of fibrin glue in repair of peripheral nerves. *Orthopedic. Clinics of North America*, 19, 187-199, 1988
- Narindrasorasak**, S., Lowery, D.E., Altman, R.A., Gonzalez-DeWhitt, P.A., Greenberg, B.D., and Kisilevsky, R. Characterization of high affinity binding between laminin and Alzheimer's disease amyloid precursor protein. *Lab. Invest.* 67, 643-652, 1992
- Narindrasorasak**, S., Lowery, D., Gonzalez-DeWhitt, P. Poorman, R.A., Greenberg, B., and Kisilevsky, R. High affinity interactions between Alzheimer's β -amyloid precursor protein and the basement membrane form of heparan sulfate proteoglycan. *J. Biol. Chem.*, 266, 12878-12883, 1991
- Navarro**, B., Kennedy, M.E., Velimirovic, B., Bhat, D., Peterson, A.S., and Clapham, D.E. Nonselective and $G_{\beta\gamma}$ -insensitive weaver K^+ Channels. *Science* 272, 1950-1953, 1996
- Neill**, D., Hughes, D., Edwardson, J.A., Rima, B.K., and Allsop, D. Human IMR-32 neuroblastoma cells as a model cell line in Alzheimer's research. *J. Neurosci. Res.*, 39, 482-493, 1994
- Neugebauer**, K.M., Emmet, C.J., Venstrom, K.A., and Reichardt, L.F. Vitronectin and Thrombospondin promote retinal neurite outgrowth: developmental regulation and role of integrins. *Neuron*, 6, 345-358, 1991
- Nissinen**, M., Vuolteenaho, R., Boot-Handfort, R., Kallunki, P. and Tryggvason, K. The primary structure of human laminin A chain. Limited expression in human tissues. *Biochem. J.*, 276, 369-379, 1991
- Noakes**, P.G., Gautam, M., Mudd, J., Sanes, J.R., and Merlie, J.P. Aberrant differentiation of neuromuscular junctions in mice lacking s-laminin/ laminin $\beta 2$. *Nature*, 374, 258-262, 1995a
- Noakes**, P.G., Miner, J.H., Gautam, M., Cunningham, J.M., Sanes, J.R., and Merlie, J.P. The renal glomerulus of mice lacking s-laminin/laminin $\beta 2$: nephrosis despite molecular compensation by laminin $\beta 1$. *Nature Gen.*, 10, 400-406, 1995b
- Nomizu**, M., Utani, A., Beck, K., Otaka, A., Roller, P.P., and Yamada, Y. Mechanism of laminin chain assembly into a triple-stranded coiled-coil structure. *Biochemistry* 35, 2885-2893, 1996
- Nomizu**, M., Yamamura, K., Kleinman, H.K., and Yamada, Y. Multimeric forms of Tyr-Ile-Gly-Ser-Arg (YIGSR) peptide enhance the

inhibition of tumor growth and metatasis. *Cancer Res.*, 53, 3459-3461, 1993

Oberley, T.D., and Steinert, B.W. Effect of the extracellular matrix molecules fibronectin and laminin on the adhesion and growth of primary renal cortical epithelial cells. *Virchows Arch.*, 44, 337-354, 1983

Ohki, K., and Kohasi, O. Laminin promotes proliferation of bone marrow-derived macrophages and macrophage cell lines. *Cell Struct. Funct.*, 19, 63-71, 1994

O'Leary, D.D.M., and Cowan, W.M. Survival of isthmo-optic neurons after early removal of one eye. *Brain Res.*, 314, 293-310, 1984

Olson, L. Regeneration in the adult central nervous systems: Experimental repair strategies. *Nature Med.*, 3, 1329-1335, 1997

Omar, R., and Pappolla, M. Oxygen free radicals as inducers of heat shock protein synthesis in cultured human neuroblastoma cells: relevance to neurodegenerative disease. *Eur. Arch. Psychiatry Clin. Neurosci.*, 242, 262-267, 1993

Ono, K., Tomasiewics, H., Magnuson, T., and Rutishauser, U. N-CAM Mutation Inhibits tangential migration and is phenocopied by enzymatic removal of polysialic acid. *Neuron*, 13, 595-609, 1994

Oo, T.F., Blazeski, R., Harrison, S.M., Henchcliffe, C., Mason, C.A., Roffler-Tarlov, S.K., and Burke, R.E. Neuronal death in the substantia nigra of weaver mouse occurs late in development and is not apoptotic. *J. Neurosci.*, 16, 6134-6145, 1996

Oppenheim, R.W. Cell death during the development of the nervous system. *Ann. Rev. Neurosci.*, 14, 1356-1362, 1991

O'Rourke, N.A., Dailey, M.E., Smith, S.J., and McConnell, S.K. Diverse migratory pathways in the developing cerebral cortex. *Science*, 258, 299-302, 1992

O'Rourke, N.A., Sullivan, D.P., Kaznowski, C.E., Andcobs, A.A., and McConnell, S.K. Tangential migration in the developing

cerebral cortex. *Development*, 121, 2165-2176, 1995

O'Shea, K.S., Reinheimer, J.S., and Dixit, V.M. Deposition and role of thrombospondin in the histogenesis of the cerebellar cortex. *J. Cell Biol.*, 110, 1275-1283, 1990

Ott, U., Odermatt, E., Engel, J., Furtmayr, H., and Timpl, R. Protease resistance and conformation of laminin. *Eur. J. Biochem.*, 173, 629-635, 1982

Ozeki, Y., Matsui, T., Yamamoto, Y., Funahashi, M., Hamako, J., and Titani, K. Tissue fibronectin is an endogenous ligand for galectin-1. *Glycobiology*, 5, 255-261, 1995

Paino, C.L., and Bunge, M.B. Induction of axon growth into Schwann cell implants grafted into lesioned adult rat spinal cord. *Exp. Neurol.*, 114, 254-257, 1991

Palmer, T.D., Takahashi, J., and Gage, F.H. The adult rat hippocampus contains primordial neuronal stem cells. *Mol. Cell Neurosci.*, 8, 389-404, 1997

Palva, T., Korva-, nenä- and kurkkutaudit, 2nd ed., Korvatautien tutkimussäätiö, Helsinki 1982

Pan, T.-C., Kluge, M., Zhang, R.-Z., Mayer, U., Timpl, R., and Chu, M.-L. Sequence of extracellular mouse protein BM-90/ fibulin and its calcium-dependent binding to other basement membrane ligands. *Eur. J. Biochem.*, 215, 733-740, 1993

Panayotou, G., End, P., Aumailley, M., Timpl, R., and Engel, J. Domains of laminin with growth activity. *Cell*, 56, 93-101, 1989

Parker, W.D.Jr., Filley, C.F., and Parks, J.K. Cytochrome oxidase deficiency in Alzheimer's disease. *Neurology*, 40, 1302-1303, 1990

Patil, N., Cox, D.R., Bhat, D., Faham, M., Myers, R.M., and Peterson, A.S. A potassium channel mutation in weaver mice implicates membrane excitability in granule cell differentiation *Nat. Genet.*, 11, 126-129, 1995

- Pattaramalai**, S., Skubitz, K.M., and Skubits, A.P.N. A novel recognition site on laminin for the $\alpha 3\beta 1$ integrin. *Exp. Cell Res.*, 222, 281-290, 1996
- Paulsson**, M. Basement membrane proteins: Structure, assembly, and cellular interactions. *Crit. Rev. Biochem. Mol. Biol.*, 27, 93-127, 1992
- Paulsson**, M., Aumailley, M., Deutzmann, R., Timpl, R., Beck, K., and Engel, J. Laminin-nidogen complex. Extraction with chelating agents and structural characterization. *Eur. J. Biochem.*, 166, 11-19, 1987
- Pearson**, V.L., Rothwell, N.J., and Toulmond, S. Excitotoxic brain damage in the rat induces interleukin-1 β in microglia and astrocytes: Correlation with the progression of cell death. *Glia*, 25, 311-323, 1999
- Pepys**, M.B., Rademacher, T.W., Amataykul, C.S., Williams, P., Noble, G.E., Hutchinson, W.L., Hawkins, P.N., Nelson, S.R., Gallimore, J.R., Herbert, J., Hutton, T., and Dwek, R.A. Human serum amyloid P component is an invariant constituent of amyloid deposits and has a uniquely homogeneous glycostructure. *Proc. Natl. Acad. Sci. U.S.A.*, 91, 5602-5606, 1994
- Pereztur**, J., Froelich, S., Prihar, G., Crook, R., Baker, M., Duff, K., Wragg, M., Busfield, F., Lendon, C., Clark, R.F., Roques, P., Fuldner, R.A., Johnston, J., Cowburn, R., Forsell, C., Axelman, K., Lilius, L., Houlden, H., Karran, E., Roberts, G.W., Rossor, M., Adams, M.D., Hardy, J., Goate, A., Lannfelt, L., and Hutton, M. A mutation in Alzheimer's disease destroying a splice acceptor site in the presenilin-1 gene. *Neuroreport*, 7, 297-301, 1995
- Pericak-Vance**, M.A., Bebout, J.L., Gaskell, P.C. Jr., Yamaoka, L.H., Hung, W.Y., Alberts, M.J., Walker, A.P., Bartlett, R.J., Haynes, C.A., Welsh, K.A., Earl, N.L., Heyman, A., Clarh, C.M., and Roses, A.D. Linkage studies in familial Alzheimer disease: evidence for chromosome 19 linkage. *Am. J. Hum. Genet.*, 48, 1034-1050, 1991
- Perlmutter**, L.S. and Chui, H.D. Microangiopathy, the vascular basement membrane and Alzheimer's disease. A review. *Brain Res. Bull.* 24, 677-686, 1990
- Peters**, B.P., Hartle, R.J., Krzesicki, R.F., Kroll, T.G., Perini, F., Balun, J.E., Goldstein, I.J., and Ruddon, R.W. The biosynthesis, processing and secretion of laminin by human choriocarcinoma cells. *J. Biol. Chem.*, 260, 14732-14742, 1985
- Pfeiffer**, S., Andkob, H., Mikoshiba, K., Dubois, B., Guenet, J., Nicolas, J-F., Gaillard, J., Chevance, G. and Andcob, F. Differentiation of a teratocarcinoma line: preferential development of cholinergic neurons. *J. Cell Biol.* 88, 57-66, 1981
- Piccolo**, S., Sasai, Y., Lu, B., and De Robertis, E. Dorsoroventral patterning in *Xenopus*: inhibition of ventral signals by direct binding of chordin to BMP-4. *Cell*, 86, 589-598, 1996
- Pikkarainen**, T., Eddy, R. Fukushima, Y. Byeres, M., Shows, T., Pihlajaniemi, T., Saraste, M. and Tryggvason, K. Human laminin B1 chain: A Multidomain protein with gene (LAMB1) locus in the g22 region of chromosome 7. *J. Biol. Chem.* 262, 10454-10462, 1987
- Pikkarainen**, T., Kallunki, T., and Tryggvason K. Human laminin B2 chain: Comparison of the complete amino acid sequence with the B1 chain reveals variability in sequence homology between different structural domains. *J. Biol. Chem.* 263, 6751-6758, 1988
- Pilar**, G., Landmesser, L., and Burstein, L. Competition for survival among developing ciliary ganglion cells. *J. Neurophysiol.* 43, 233-254, 1980
- Pilz**, D.T., Matsumoto, N., Minnerath, S., Mills, P., Gleeson, J.G., Allen, K.M., Walsh, C.A., Barkovich, A.J., Dobyns, W.B., Ledbetter, D.H., and Ross, M.E. *LIS1* and *XLIS* (DCX) mutation cause most classical lissencephaly, but different patterns of malformation. *Hum. Mol. Genet.*, 7, 2092-2037, 1998

- Porter**, B.E., and Sanes, J.R. Gated migration: Neurons migrate on but not onto substrates containing S-laminin. *Dev. Biol.*, 167, 609-616, 1995
- Porter**, B.E., Wels, J., and Sanes, J.R. A motoneuron-selective stop signal in the synaptic protein S-laminin. *Neuron*, 14, 549-559, 1995
- Powell**, S.K., Williams, C.C., Nomizu, M., Yamada, Y., and Kleinman, H.K. Laminin-like proteins are differentially regulated during cerebellar development and stimulate granule cell neurite outgrowth in vitro. *J. Neurosci. Res.*, 54, 233-247, 1998
- Prinjha**, R., Moore, S.E., Vinson, M., Blake, S., Morrow, R., Christie, G., Michalovich, D., Simmons, D.L., and Walsh, F.S. Inhibitor of neurite outgrowth in humans. *Nature*, 403, 383-384, 2000
- Påhlman**, S., Hoehner, J.C., Nåberg, E., Hedborg, F., Fagerström, S., Gestblom, C., Johansson, I., Larsson, U., Lavenius, E., Örtoft, E., and Söderholm, H. Differentiation and survival influences of growth factors in human neuroblastoma.
- Quarles**, R.H., and Trapp, B.D. Localization of myelin-associated glycoprotein. *J. Neurochem.*, 43, 1773-1777, 1984
- Raabe**, E.H., Yoshida, K., and Schwarting, G.A. Differential laminin isoform expression in the developing rat olfactory system. *Dev. Brain Res.*, 101, 187-196, 1997
- Rabchevsky**, A.G., and Streit, W.J. Grafting of cultured microglial cells into the lesioned spinal cord of adult rats enhances neurite outgrowth. *J. Neurosci. Res.*, 47, 34-48, 1997
- Raff**, M.C., Barres, B.A., Burne, J.F., Coles, H.S., Ishizaki, Y., and Andocobson, M.D. Programmed cell death and the control of cell survival: Lessons from the nervous system. *Science*, 262, 695-700, 1993
- Rakic**, P. Principles of neural cell migration. *Experientia* 46, 882-891, 1990
- Rakic**, P., and Sidman, R. Weaver mutant mouse cerebellum: Defective neuronal migration secondary to abnormality of Bergmann glia. *Proc. Natl. Acad. Sci. U.S.A.*, 70, 240-244, 1973a
- Rakic**, P., and Sidman, R. Sequence of developmental abnormalities leading to granule cell deficit in cerebellar cortex of weaver mutant mice. *J. Comp. Neurol.* 152, 103-132, 1973b
- Rakic**, P., and Sidman, R. Organization of cerebellar cortex secondary to deficit of granule cells in weaver mutant mouse. *J. Comp. Neurol.* 152, 133-162, 1973c
- Rao**, C.N., Barsky, S.H., Terranova, V.P., and Liotta, L.A. Isolation of a tumor cell laminin receptor. *Biochem. Biophys. Res. Commun.* 11, 804-808, 1983
- Rapalino**, O., Lazarov-Spiegler, O., Agranov, E., Velan, G.J., Yoles, E., Fraidakis, M., Solomon, A., Gepstein, R., Katz, A., Belkin, M., Hadani, M., and Schwartz, M. Implantation of stimulated homologous macrophages results in partial recovery of paraplegic rats. *Nat. Med.*, 4, 814-821, 1998
- Rathjen**, F., and Schachner, M. Immunocytological and biochemical characterization of a new neuronal cell surface component (L1 antigen) which is involved in cell adhesion. *EMBO J.*, 3, 461-465, 1984
- Reeves**, R., Crowley, M., Lorenzon, N., Paven, W., Smeyne, R. and Goldowitz, R. The mouse neurological mutant weaver maps within the region of chromosome 16 that is homologous to human chromosome 21. *Genomics* 5, 522-526, 1989
- Reh**, T. A., Nagy, T. and Gretton, H. Retinal pigmented epithelial cells induced to transdifferentiate to neurons by laminin. *Nature* 330, 68-71, 1987
- Reynolds**, B.A., and Weiss, S. Generation of neurons and astrocytes from isolated cells of the adult mammalian central nervous systems. *Science*, 255, 1707-1710, 1992
- Reynolds**, B.A., and Weiss, S. Clonal and population analyses demonstrate that an EGF-

responsive mammalian embryonic precursor is a stem cell. *Dev. Biol.*, 175, 1-13, 1996

Richard, B.L., Nomizu, M., Yamada, Y., and Kleinman, H.K. Identification of synthetic peptides derived from laminin $\alpha 1$ and $\alpha 2$ chains with cell type specificity for neurite outgrowth. *Exp. Cell Res.*, 228, 98-105, 1996

Richardson, C.A., Gordon, K.L., Couser, W.G., and Bomsztyk, K. IL-1 β increases laminin B2 chain mRNA levels and activates NF κ B in rat glomerular epithelial cells. *Am. Physiol. Soc.*, 268, F273-F278, 1995

Risteli, L., and Risteli, J. Analysis of extracellular matrix proteins in biological fluids. *Methods. Enzymol.*, 145, 391-411, 1987

Risteli, L., and Timpl, R. Isolation and characterization of pepsin fragments of laminin from human placental and renal basement membranes. *Biochem. J.*, 193, 749-755, 1981

Rizzino, A., Terranova, V.P., Rohrbach, D., Crowley, C., and Rizzino, H. The effects of laminin on the growth and differentiation of embryonal carcinoma cells in defined media. *J. Supramol. Struct.*, 13, 243-253, 1980

Rober, A.E., Ball, M.J., Bhawe, S.V. and Wakade, A.B. β - Amyloid from Alzheimer's disease brains inhibits sprouting and survival of sympathetic neurons. *Biochem. Biophys. Res. Commun* 174, 572-579. 1991

Roberts, D.D., Rao, C.N., Magnani, J.L., Spitalnik, S.L., Liotta, L.A., and Ginsburg, V. Laminin binds specifically to sulfated glycolipids. *Proc. Natl. Acad. Sci. USA*, 82, 1306-1310, 1985

Roffler-Tarlov, S. and Graybiel, A. Expression of the weaver gene in dopamine-containing neural systems is dose-dependent and affects both striatal and nonstriatal regions. *J. Neurosci.*, 6, 3319-3330, 1986

Rogaev, E.I., Sherrington, R., Rogaeva, E.A., Levesque, G., Ikeda, M., Liang, Y., Chi, H., Lin, C., Holamn, K., Tsuda, T., Mar, L., Sorbi, S., Nacmias, B., Piacentini, S., Amaducci, L., Chumakov, I., Cohen, D.,

Lannfelt, L., Fraser, P.E., Rommens, J.M., and St. George-Hyslop, P.H. Familial Alzheimer's disease is kindreds with missense mutation in a gene on chromosome 1 related to the Alzheimer's disease type 3 gene. *Nature*, 376, 775-778, 1995

Rothstein, J.D., Bristol, L.A., Hosler, B., Brown, R.H., and Kuncl, R.W. Chronic inhibition of superoxide dismutase produces apoptotic death of spinal neurons. *Proc. Natl. Acad. Sci. USA*, 91, 4155-4159, 1994

Rousselle, P., Lundstrum, G.P., Keene, D.R., and Burgeson, R.E. Kalin: an epithelium-specific basement membrane adhesion molecule that is a component of anchoring filaments. *J. Cell Biol.*, 114, 567-576, 1991

Rovasio, R.A., Delouvé, A., Yamada, K.M., Timpl, R., and Thiery, J.P. Neural crest cell migration: Requirements for exogenous fibronectin and high cell density. *J. Cell Biol.*, 96, 462-473, 1983

Ryan, M.C., Lee, K., Miyashita, Y., Carter, W.G. Targeted distribution of the LAMA3 gene in mice reveals abnormalities in survival and late stage differentiation of epithelial cells. *J. Cell Biol.*, 145, 1309-1323, 1999

Ryan, M.C., Tizard, R., VanDevanter, D.R., and Carter, W.G. Cloning of the LamA3 gene encoding the $\alpha 3$ chain of adhesive ligand epiligrin. *J. Biol. Chem.*, 269, 2279-22787, 1994

Rønn, L.C.B., Hartz, B.P., Bock, E. The neural adhesion molecule (NCAM) in development and plasticity of the nervous system. *Exp. Geront.*, 33, 853-864, 1998

Saarela, S., Westerlund-Wikstrom, B., Rhen, M., and Korhonen, T.K. The GafD protein of the G (F17) fimbrial complex confers adhesiveness of Escherichia coli to laminin. *Infection & Immunity*, 64, 2857-2860, 1996

Saiki, I., Murata, J., Makabe, T., Nishi, N., Tokura, S., and Azuma, I. Inhibition of tumor angiogenesis by a synthetic cell-adhesive polypeptide containing the Arg-Gly-Asp (RGD) sequence of fibronectin, poly(RGD). *Jnp. J. Cancer Res.*, 81, 668-675, 1990

- Saiki, R.K.,** Scharf, S., Faloona, F., Mullis, K.B., Horn, T., Erlich, H.A., and Arnheim, N. Enzymatic amplification of β -globin genomic sequence and restriction site analysis for diagnostics of sickle cell anemia. *Science*, 230, 1350-1354, 1985
- Sakamoto, N.,** Iwahana, M., Tanaka, N.G., and Osada, Y. Inhibition of angiogenesis and tumor growth by a synthetic laminin peptide, CDPGYIGSR-NH₂. *Cancer Res.*, 51, 903-906, 1991
- Sanger, F.,** Nicklen, S., and Coulson, A.R. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA*, 74, 5463-5467, 1977
- Sanes J.R.** Extracellular matrix molecules that influence neural development *Annu Rev. Neurosci.* 12, 491-516, 1989
- Sanes, J.R.,** Engvall, E., Butkowski, R., and Hunter, D.D. Molecular heterogeneity of basal laminae: Isoforms of laminin and collagen IV at the neuromuscular junction and elsewhere. *J. Cell Biol.* 111, 1685-1699, 1990
- Sapir, T.,** Elbaum, M., and Reiner, O. Reduction of microtubule catastrophe events by LIS1, platelet-activating factor acetylhydrolase subunit. *EMBO J.*, 16, 6977-6984
- Sarthy, P.V.,** and Fu, M. Localization of laminin B1 mRNA in retinal ganglion cells by in situ hybridization. *J. Cell Biol.*, 110, 2099-2108, 1996
- Sasai, Y.** Identifying the missing links: genes that connect neural induction and primary neurogenesis in vertebrate embryos. *Neuron*, 21, 455-458, 1998
- Sasai, Y.,** Lu, B., Steinbeisser, H., and De Robertis, E.M. Regulation of neural induction by the *chc* and BMP-4 antagonistic patterning signals in *Xenopus*. *Nature*, 376, 333-336, 1995
- Sasai, Y.,** Lu, B., Steinbeisser, H., Geissert, D., Gont, L.K., and De Robertis, E.M. *Xenopus* chordin: a novel dorsalizing factor activated by organizer-specific homeobox genes. *Cell*, 79, 779-790, 1994
- Sasaki, M.,** Kato, S., Kohno, K. Martin, G.R., and Yamada, Y. Sequence of the cDNA encoding the laminin B1 chain reveals a multidomain protein containing cysteine-rich repeats. *Proc. Natl. Sci. USA* 84, 935-939, 1987
- Sasaki, M.,** Kleinman, H.K., Huber, H. Deutzmann, R. and Yamada, Y. Laminin, a multidomain protein. *J. Biol. Chem.* 263, 16536-16544, 1988
- Sasaki, M.,** and Yamada, Y. The laminin B2 chain has a multidomain structure homologous to the B1 chain. *J. Biol. Chem.* 262, 17111-17117, 1987
- Sasaki, T.,** Göhring, W., Pan, T.C., Chu, M.L., and Timpl, R. Binding of mouse and human fibulin-2 to extracellular matrix ligands. *J. Mol. Biol.*, 245, 892-899, 1995a
- Sasaki, T.,** Kostka, G., Göhring, W., Wiedemann, H., Mann, K., Chu, M.L., and Timpl, R. Structural characterization of two variants of fibulin-1 that differ in nidogen affinity. *J. Mol. Biol.*, 245, 241-250, 1995b
- Sato, S.M.,** and Sargent, T.D. Development of neural inducing capacity in dissociated *Xenopus* embryos. *Dev. Biol.*, 143, 263-266, 1989
- Sauer, F** Mitosis in the neural tube. *J. Comp. Neurol.*, 62, 377-405, 1935
- Scheibel, A.B,** and Tomiyasu, U. Dendritic sprouting in Alzheimer's presenile dementia. *Exp. Neurol.*, 60, 1-8, 1978
- Schachner, M.G.,** Schoonmaker, G., and Hynes, R.O. Cellular and subcellular localization of LETS protein in the nervous system. *Brain Res.*, 158, 149-158, 1978
- Schittny, J.C.,** and Yurchenco, P.D. Terminal short arm domains of basement membrane laminin are critical for its self-assembly. *J. Cell Biol.*, 110, 825-832, 1990
- Schmidt, M.J.,** Sawyer, B.D., Perry, K.W., Fuller, R.W., Foreman, M.M., and Ghetti, B.

Dopamine deficiency in the weaver mutant mouse. *J. Neurosci.*, 2, 376-380, 1982

Schnell, L., Schneider, R., Kolbeck, R., Barde, Y.A., and Schwab, M.E. Neurotrophin-3 enhances sprouting of corticospinal tract during development and after adult spinal cord lesion. *Nature*, 367, 170-173, 1994

Schnell, L., and Schwab, M.E. Axonal regeneration in the rat spinal cord produced by an antibody against myelin-associated neurite growth inhibitors. *Nature*, 343, 269-272, 1990

Schwab, M.E., and Caroni, P. Oligodendrocytes and myelin are nonpermissive substrates for neurite growth and fibroblast spreading in vitro. *J. Neurosci.*, 8, 2381-2393, 1988

Schwarzman, A.L., Singh, N., Tsiper, M., Gregori, L., Dranovsky, A., Vitek, M.P., Glabe, C.G., St. George-Hyslop, P.H., and Goldgaber, D. Endogenous presenilin 1 redistributes to the surface of lamellipodia upon adhesion of Jurkat cells to a collagen matrix. *Proc. Natl. Acad. Sci. USA*, 96, 7932-7937, 1999

Seeds, N.W., Haffke, S., Christensen, K., and Schoonmaker, J. Cerebellar granule cell migration involves proteolysis. *Add. Exp. Biol. Med.*, 265, 169-178, 1990

Seeds, N.W., Siconolfi, L.B., and Haffke, S.P. Neural extracellular proteases facilitate cell migration, axonal growth, and pathfinding. *Cell Tiss. Res.*, 290, 367-370, 1997

Selkoe, D.J. Missense on the membrane. *Nature*, 375, 734-735, 1995

Selkoe, D.J., Abraham, C.R., Podlisny, M.B., and Duffy, L.K. Isolation of low-molecular-weight proteins from amyloid plaque fibers in Alzheimer's disease. *J. Neurochem.*, 46, 1820-1834, 1986

Sell, S. and Pierce, G.B. Maturation arrest of stem cell differentiation is a common pathway for the cellular origin of teratocarcinomas and epithelial cancers. *Lab. Invest.* 70, 6-22, 1994

Semaphorin nomenclature committee, Unified nomenclature for the semaphorins/collapsins. *Cell*, 97, 551-552, 1999

Serafini, T., Colamarino, S.A., Leonardo, E.D., Wang, H., Beddington, R., Skarnes, W.C., Tessier-Lavigne, M. Netrin-1 is required for commissural axon guidance in the developing vertebrate nervous system. *Cell*, 87, 1001-1014, 1996

Serafini, T., Kennedy, T.E., Galko, M.J., Mirzayan, C., Jessell, T.M., and Tessier-Lavigne, M. The netrins define a family of axon outgrowth-promoting proteins homologous to *C. elegans* UNC-6. *Cell*, 78, 409-424, 1994

Sherrington, R., Rogaev, E.I., Liang, Y., Rogaeva, E.A., Levesque, G., Ikeda, M., Chi, H., Lin, C., Li, G., Holman, K., Tsuda, T., Mar, L., Foncin, J.-F., Bruni, A.C., Montesi, M.P., Sorbi, S., Rainero, I., Pinessi, L., Nee, L., Chumakov, I., Pollen, D., Brookes, A., Sanseau, P., Polinsky, R.J., Wasco, W., da Silva, H.A.R., Haines, J.L., Pericak-Vance, M.A., Tanzi, R.E., Roses, A.D., Fraser, P.E., Rommens, J.M., and St. George-Hyslop, P.H. Cloning of a gene bearing missense mutations in early-onset familial Alzheimer's disease. *Nature*, 375, 754-760, 1995

Sidman, R., Green, M. and Appel, S. "Catalogue of neurological mutants of the mouse." Harvard Univ. Press. Cambridge. 1965

Sidman, R. and Rakic, P. Neuronal migration, with special reference to developing human brain. A review. *Brain Res.* 62, 1-35, 1973

Signorini, S., Liao, Y.J., Duncan, S.A., Andn, L.Y., and Stoffel, M. Normal cerebellar development but susceptibility to seizures in mice lacking G protein-coupled, inwardly rectifying K⁺ channels. *Proc. Natl. Acad. Sci.*, 94, 923-927, 1997

Siman, R.M., Card, J.P., Nelson, R.B., and Davis, L.G. Expression of β -amyloid precursor protein in reactive astrocytes

following neuronal damage. *Neuron*, 3, 275-285, 1989

Skubitz, A.P.N., McCarthy, J.B., Charonis, A.S., and Frucht, L.T. Localization of three distinct heparin-binding domains of laminin by monoclonal antibodies. *J. Biol. Chem.*, 263, 4861-4868, 1988

Skubitz, A.P.N., Letourneau, P.C., Wayner, E., and Furcht, L.T. Synthetic peptides from the carboxy-terminal globular domain of the A chain of laminin: Their ability to promote cell adhesion and neurite outgrowth, and interact with heparin and the $\beta 1$ integrin subunit. *J. Cell Biol.*, 115, 1137-1148, 1991

Slesinger, P.A., Patil, N., Liao, Y.J., Andn, Y.N., and Cox, D.R. Functional effects of the mouse weaver mutation on G protein-gated inwardly rectifying K⁺. *Neuron*, 16, 321-331, 1996

Smale, G., Nichols, N.R., Brady, D.R., Finch, C.E., and Horton, W.E. Jr. Evidence for apoptotic cell death in Alzheimer's disease. *Exp. Neurol.*, 133, 225-230, 1995

Smalheiser, N.R., Cranin interacts specifically with the sulfatide-binding domain of laminin. *J. Neurosci. Res.*, 36, 528-538, 1993

Smalheiser, N.R. and Kim, E. Purification of cranin, a laminin binding membrane protein. Identification with dystroglycan and reassessment of its carbohydrate moieties. *J. Biol.Chem.*, 270, 15425-15433, 1995

Smalheiser, N.R., and Schwartz, N.B. Cranin: a laminin-binding protein of cell membranes. *Proc. Natl. Sci. USA*, 84, 6457-6461, 1987

Smeyne, R.J. and Goldowitz, D. Purkinje cell loss is due to a direct action of the weaver gene in Purkinje cells: evidence from chimeric mice. *Brain Res. Dev. Brain Res.*, 52, 211-218, 1990

Smith, G., Miller, R., and Silver, J. Changing role of forebrain astrocytes during development, regenerative failure and induced regeneration upon transplantation. *J. Comp. Neurol.*, 251, 23-43, 1986

Smith, W.B., and Harland, R.M. Expression cloning of noggin, a new dorsalizing factor localized to the Spemann organizer in *Xenopus* embryos. *Cell*, 70, 829-840, 1992.

Smyth, N., Vatansever, H.S., Murray, P., Meyer, M., Frie, C., Paulsson, M., and Edgar, D. Absence of basement membranes after targeting the LAMC1 gene results in embryonic lethality due to failure of endoderm differentiation. *J. Cell Biol.*, 144, 151-160, 1999

Snow, A., and Wight, T.N. Proteoglycans in the pathogenesis of Alzheimer's disease and other amyloidoses. *Neurobiol. Aging*, 10, 481-497, 1989

Snow, A., Mar, H., Nochlin, D., Kresse, H., and Wight, T.N. Peripheral distribution of dermatan sulfate proteoglycans (Decorin) in amyloid-containing plaques and their presence in neurofibrillary tangles of Alzheimer's disease. *J. Histochem. Cytochem.*, 40, 105-113, 1992

Snow, A., Mar, H., Nochlin, D., Kimata, K., Kato, M., Suzuki, S., Hassell, J., and Wight, T.N. The presence of heparan sulfate proteoglycan in the neurite plaques and congophilic angiopathy in Alzheimer's disease. *Am. J. Pathol.*, 133, 456-463, 1988

Snow, A., Mar, H., Nochlin, D., Sekiguchi, R.T. Kimata, K., Koike, Y. and Wight, T.N. Early accumulation of heparan sulfate in neurons and in the β -amyloid protein-containing lesions of Alzheimer's disease and Down's syndrome. *Am. J.Pathol.*, 137, 1253-1269, 1990

Snow, A., Nochlin, D., Sekiguchi, R.T., and Carlson, S.S. Identification and immunolocalization of a new class of proteoglycans (keratan sulfate) to the neuritic plaques of Alzheimer's disease. *Exp. Neurol.*, 138, 305-317, 1996a

Snow, A., Sekiguchi, R., Nochlin, D., Fraser, P., Kimata, K., Mizutani, A., Arai, M., Schreier, W., and Morgan, D. An Important role of heparan sulfate proteoglycan (Perlecan) in a model system for the deposition and persistence of fibrillar A β -

amyloid in rat brain. *Neuron*, 12, 219-234, 1994

Snow, D.M., Brown, E.M. and Letourneau, P.C. Growth cone behaviour in the presence of soluble chondroitin sulfate proteoglycan (CSPG), compared to behaviour on CSPG bound to laminin or fibronectin. *Int. J. Dev. Neuroscience*, 14, 331-349, 1996b

Song, D-L, Chalepakis, G., Gruss, P., and Joyner, A.L. Two Pax-binding sites are required for early embryonic expression of an *Engrailed-2* transgene. *Development*, 122, 627-635, 1996

Sotelo, C. Anatomical, physical and biochemical studies of the cerebellum from mutant mice. II. Morphological study of cerebellar cortical neurons and circuits in the weaver mouse. *Brain Res.* 94, 19-44, 1975

Sotelo, C., and Changeux, J-P. Bergmann fibres and granular cell migration in the cerebellum of homozygous weaver mutant mouse. *Brain Res.*, 77, 484-491, 1974

Sotelo, C., and Rio, J. Cerebellar malformation obtained in rats by early postnatal treatment with 6-aminonicotinamide. Role of neuron-glia interactions in cerebellar development. *Neuroscience*, 5, 1737-1759, 1980

Sporn, M., and Roberts, A. Role of retinoids in differentiation and carcinogenesis. *Cancer Res.* 43, 3034-3040, 1983

Stack, S., Gray, R.D., and Pizzo, S.V. Modulation of plasminogen activation and type IV collagenase activity by a synthetic peptide derived from the laminin A chain. *Biochemistry*, 30, 2073-2077, 1991

Steadman, R., Irwin, M.H., St John, P.L., Blackburn, W.D., Heck, L.W., and Abrahamson, D.R. Laminin cleavage by activated human neutrophils yields proteolytic fragments with selective migratory properties. *J. Leukocyte Biology*, 53, 354-365, 1993

St George-Hyslop, P.H., Tanzi, R.E., Polinsky, R.J., Haines, J.L., Nee, L., Watkins, P.C., Myers, R.H., Feldman, R.G., Pollen, D.,

and Drachman, D. The genetic defect causing familial Alzheimer's disease maps on chromosome 21. *Science*, 235, 885-890, 1987

Stoeckli, E.T., and Landmesser, L.T. Axonin-1, Nr-CAM, and Ng-CAM play different roles in the in vivo guidance of chick commissural neurons. *Neuron*, 14, 1165-1229, 1995

Stopa, E.G., Gonzales, A-M., Chorsky, R., Corona, R.J., Alvasrez, J., Bird, E.D. , and Baird, A. Basic fibroblast growth factor in Alzheimer's disease. *Biochem. Biophys. Res. Commun.*, 171, 690-696, 1990

Streit, A., Nolte, C., Rásony, and Schachner, M. Interaction of astrochondrin with extracellular matrix components and its involvement in astrocyte process formation and cerebellar granule cell migration. *J. Cell Biol.*, 120, 799-814, 1993

Strickland, S. and Mahdavi, V. The induction of differentiation in teratocarcinoma cells by retinoic acid. *Cell*, 15, 393-403, 1978

Sugiyama, S., Utani, A., Yamada, S., Kozak, C.A., and Yamada, Y. Cloning and expression of the mouse laminin $\gamma 2$ (B2t) chain, a subunit of epithelial cell laminin. *Eur. J. Biochem.*, 228, 120-128, 1995

Sunada, Y., Edgar, T.S., Lotz, B.P., Rust, R.S., and Campbell, K.P. Merosin-negative congenital muscular dystrophy associated with extensive brain abnormalities. *Neurology*, 45, 2084-2089, 1995

Sunderland, S. Nerve injuries and their repair, 1st ed. Churchill Livingstone, London, 1991

Surmeier, D.J., Mermelstein, P.G., and Goldowitz, D. The weaver mutation of GIRK2 results in a loss of inwardly rectifying K^+ current in cerebellar granule cells. *Proc. Natl. Acad. Sci. U.S.A.*, 93, 11191-11195, 1996

Svendsen, C.N., Caldwell, M.A., Shen, J., Borg, M.G., Rosser, A.E., Tyers, P., Karmiol, S., and Dunnet, S.B. Long-term survival of human central nervous system progenitor

cells transplanted into a rat model of Parkinson's disease. *Exp. Neurol.*, 148, 135-146, 1997

Sweeney, T.M., Ogle, R.C., and Little, C.D. Laminin potentiates differentiation of PCC4uva embryonal carcinoma into neurons. *J. Cell Sci.*, 97, 23-31, 1990

Takei, K., Chan, T.A., Wang, F.S., Deng, H., Rutishauser, U., and Andy, D.G. The neural cell adhesion molecules L1 and NCAM-180 act in different steps of neurite outgrowth. *J. Neurosci.*, 19, 9469-9479, 1999

Tanabe, Y., and Jessel, T.M. Diversity and pattern in the developing spinal cord. *Science*, 274, 1115-1123, 1996

Tanzi, R.E., Gusella, J.F., Watkins, P.C., Bruns, G.A.P., St. George-Hyslop, P.H., Van Keuren, M.L., Patterson, D., Pagan, S., Kurnit, D.M., and Neve, R.L. Amyloid β protein gene: c-DNA, mRNA distribution, and genetic linkage near the Alzheimer locus. *Science*, 235, 880-884, 1987

Tashiro, K., Nagata, I., Yamashita, N., Okazaki, K., Ogomori, K., Tashiro, N., and Anai, M. A synthetic peptide deduced from the sequence in the cross-region of laminin A chain mediates neurite outgrowth, cell attachment and heparin binding. *Biochemical. Journal*, 302, 73-79, 1994

Tashiro, K., Sephel, G.C., Grotorex, D., Sasaki, M., Shirashi, N., Martin, G.R., Kleinman, H.K., and Yamada, Y. The RGD containing site of the mouse laminin A chain is active for cell attachment, spreading, migration and neurite outgrowth. *J. Cell Physiol.*, 146, 451-459, 1991

Tashiro, K., Sephel, G., Weeks, B., Sasaki, M., Kleinman, H. and Yamada, Y. A synthetic peptide containing the IKVAV sequence from the A chain of laminin mediates cell attachment, migration and neurite outgrowth. *J. Biol. Chem.*, 264, 16174-16182, 1989

Terranova, V.P., Liotta, L.A., Russo, R.G., and Martin, G.R. Role of laminin in the attachment and metastasis of murine tumor cells. *Cancer Res.*, 42, 2265-2269, 1982

Terranova, V.P., Rao, C.N., Kalebic, T., Margulies, I.M., and Liotta, L.A. Laminin receptor on human breast carcinoma cells. *Proc. Natl. Acad. Sci. USA*, 80, 444-448, 1983

Terzis, J.K. and Smith, K.J. *The peripheral nerve, structure, function and reconstruction*, Raven Press, New York, 1991

Tessier-Lavigne, M., and Goodman, C.S. The molecular biology of axon guidance. *Science*, 274, 1123-1133, 1996

Tian, M., Hagg, T., Denisova, N., Knusel, B., Engvall, E., and Jucker, M. Laminin alpha2 chain-like antigen in CNS dendritic spines. *Brain Res.*, 764, 28-38, 1997

Tienari, J., Alanko, T., Lehtonen, E., and Saksela, O. The expression and localization of urokinase-type plasminogen activator and its type 1 inhibitor are regulated by retinoic acid and fibroblast growth factor in human teratocarcinoma cells. *Cell Regul.*, 2, 285-297, 1991

Tienari, J., Virtanen, I., Soinila, S., and Lehtonen, E. Neuron-like derivatives of cultured F9 embryonal carcinoma cells express characteristics of parietal endoderm. *Dev. Biol.*, 123, 566-573, 1987

Timpl, R. Structure and biological activity of basement membrane proteins. *Eur. J. Biochem.*, 180, 487-502, 1989

Timpl, R. Proteoglycans of basement membranes. *EXS*, 70, 123-144, 1994

Timpl, R. Macromolecular organization of basement membranes. *Current Opinion in Cell Biology*, 9, 618-624, 1996

Timpl, R., and Brown, C.J. The laminins. *Matrix Biol.*, 14, 275-281, 1994

Timpl, R., and Martin, G. Laminin and other basement membrane components. *Annu. Rev. Cell Biol.*, 3, 57-85, 1987

Timpl, R., and Risteli, L. Radioimmunoassays in studies of connective tissue proteins: In: Furthmar H (ed):

Immunocytochemistry of the extracellular matrix. Boca Raton, FL: CRC Press, Vol. 1, pp 119-235, 1982

Timpl, R., Rohde, H., Robey, P.G., Rennard, S.I., Foidart, J-M. and Martin, G.R. Laminin, a glycoprotein from basement membranes. *J.Biol. Chem.* 254, 9933-9937, 1979

Towbin, H., Staehelin, T., Gordon, J. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: Procedure and some applications. *Proc. Natl. Acad. Sci. USA.* 76, 4350-4354, 1979

Tomaselli, K.J., Reichardt, L.F., and Bixby, J.L. Distinct molecular interactions mediate neuronal process outgrowth on non-neuronal cell surfaces and extracellular matrices. *J. Cell Biol.*, 103, 2659-2672, 1986

Tomasiewicz, H., Ono, K., Yee, D., Thompson, C., Goridis, C., Rutishauser, U., and Magnuson, T. Genetic deletion of a neural cell adhesion molecule variant (N-CAM-180) produces distinct defects in the central nervous system. *Neuron*, 11, 1163-1174, 1993

Tomé, F.M., Evangelista, T., Leclerc, A., Sunada, Y., Manole, E., Estournet, B., Barois, A., Campbell, K.P., and Fardeau, M. Congenital muscular dystrophy with merosin deficiency. *C.R. Acad. Sci. III*, 317, 351-357, 1994

Tong, X.J., Hirai, K.I., Shimada, H., Mizutani, Y., Izumi, T., Toda, N., and YU, P. Sciatic nerve regeneration navigated by laminin-fibronectin double coated biodegradable collagen grafts in rats. *Brain Res.*, 663, 155-162, 1994

Trapp, B.D. Myelin-associated glycoprotein. Location and potential functions. *Ann. N.Y. Acad. Sci.*, 605, 29-43, 1990

Trenkner, E. Cerebellar cells in culture. In "Culturing Nerve Cells" pp. 283-307, (G.Banker and K. Goslin, Eds.), MIT Press, Cambridge, MA, 1991

Trenkner, E., Hatten, M. and Sidman, R. Effect of ether-soluble serum components in vitro on the behavior of immature cerebellar

cells in weaver mutant mice. *Neuroscience* 3, 1093-1100, 1978

Tsirka, S., Rogove, A.D., Bugge, T.H., Degen, J.L., and Strickland, S. An extracellular proteolytic cascade promotes neuronal degeneration in the mouse hippocampus. *J. Neurosci.*, 17, 543-552, 1997

Underwood, P.A., Bennett, F.A., Kirkpatrick, A., Bean, P.A., and Moss, B.A. Evidence for the location of a binding sequence for the alpha 2 beta 1 integrin of endothelial cells, in the beta subunit of laminin. *Biochemical. J.*, 309, 765-771, 1995

Utani, A., Kopp, J.B., Kozak, C.A., Matsuki, Y., Amizuka, N., Sugiyama, S., and Yamada, Y. Mouse kalinin B1 (laminin $\beta 3$ chain): Cloning and tissue distribution. *Lab. Invest.*, 72, 300-310, 1995a

Utani, A., Nomizu, M., Sugiyama, S., Miyamoto, S., Roller, P.P., and Yamada, Y. A specific sequence of the laminin alpha 2 chain critical for the initiation of heterotrimer assembly. *J. Biol. Chem.*, 270, 3292-3298, 1995b

Utani, A., Nomizu, M., Timpl, R., Roller, P.P., and Yamada, Y. Laminin chain assembly. Specific sequences at the C terminus of the long arm are required for the formation of specific double- and triple-stranded coiled-coil structures. *J. Biol. Chem.*, 269, 19167-19175, 1994

van Raay, T.J., Foskett, S.M., Connors, T.D., Klinger, K.W., Landes, G.M., and Burn, T.C. The NTN2L gene encoding a novel human netrin maps to the autosomal dominant polycystic kidney disease on chromosome 16p13.3. *Genomics*, 41, 279-282, 1997

Varga, Z.M., Bandtlow, C.E., Erulkar, S.D., Schwab, M.E., and Nicholls, J.G. The critical period for repair of CNS of neonatal opossum (*Monodelphis domestica*) in culture: correlation with development of glial cells, myelin and growth-inhibitory molecules. *Eur. J. Neurosci.*, 7, 2119-2129, 1995

- Vastamäki, M.** Hermovamman hoitoperiaatteet. *Suomen Lääkärilehti*, 45, 544-546, 1990
- Verrall, S., and Seeds, N.W.** Tissue plasminogen activator binding to mouse cerebellar granule neurons. *J. Neurosci. Res.*, 21, 420-425, 1988
- Villegas-Perez, M.P., Vidal-Sanz, M., Bray, G.M., and Aguayo, A.J.** Influences of peripheral nerve grafts on the survival and regrowth of axotomized retinal ganglion cells in adult rats. *J. Neurosci.*, 8, 265-280, 1988
- Vlodavsky, I., and Gospodarowicz, D.** Respective roles of laminin and fibronectin in adhesion of human carcinoma and sarcoma cells. *Nature*, 289, 304-306, 1981
- Vukicevic, S., Luyten, F.P., Kleinman, H.K., and Reddi, A.H.** Differentiation of canalicular cell processes in bone cells by basement membrane matrix components: regulation by discrete domains of laminin. *Cell*, 63, 437-445, 1990
- von der Mark, H., Durr, J., Sonnenberg, A., von der Mark, K., Deutzmann, R., and Goosman, S.L.** Skeletal myoblast utilize a novel beta-1 integrin and not alpha6 beta 1 for binding to the E8 and T8 fragments of laminin. *J. Biol. Chem.*, 266, 23593-23601, 1991
- Vuolteenaho, R., Chow, L.T., and Tryggvason, K.** Structure of the human laminin B1 chain gene. *J. Bio. Chem.*, 265, 15611-15616, 1990
- Vuolteenaho, R., Nissinen, M., Sainio, K., Byers, M., Eddy, R., Hirvonen, H., Shows, T.B., Sariola, H., Engvall, E., and Tryggvason, K.** Human laminin-M chain (merosin) - complete primary structure, chromosomal assignment, and expression of the M-chain and A-chain in human fetal tissues. *J. Cell Biol.*, 124, 381-394, 1994
- Wall, P.D., Devor, M., Imbal, R., Scadding, J.W., Schonfeld, D., Zelzer, Z., and Tomkiewicz, M.M.** Autonomy following peripheral nerve lesions: experimental anesthesia dolorosa. *Pain*, 7, 103-113, 1979
- Wang, H., Copeland, N.G., Gilbert, D.J., Jenkins, N.A., and Tessier-Lavigne, M.** Netrin-3, a mouse homology of human NTN2L, is highly expressed in sensory ganglia and shows differential binding to netrin receptors. *J. Neurosci.*, 19, 493-4947, 1999
- Wartiovaara, J., Leivo, I., Virtanen, I., Vaheri, A., and Graham, C.** Appearance of fibronectin during differentiation of mouse teratocarcinoma *in vitro*. *Nature*, 272, 355-356, 1978
- Watt, J.A., Pike, C.J., Walencewicz Wasserman, A.J., and Cotman, C.W.** Ultrastructural analysis of beta-amyloid-induced apoptosis in cultured hippocampal neurones. *Brain Res.*, 661, 147-156, 1994
- Weeks, B.S., Holloway, E., Klotman, P.E., Akiyama, S.K., Schnaper, H.W., and Kleinman, H.K.** 12-O-tetradecanoylphorbol 13-acetate stimulates human T-lymphocyte adherence to the fibronectin RGD domain and the laminin IKVAV domain. *Cellular Immunology*, 153, 94-104, 1994
- Weeks, B.S., Nomizu, M., Ramchandran, R.S., Yamada, Y., and Kleinman, H.K.** Laminin-1 and the RKRLQVQLSIRT laminin-1 $\alpha 1$ globular domain peptide stimulate matrix metalloproteinase secretion by PC12 cells. *Exp. Cell Res.*, 243, 375-382, 1998
- Weiss, S., Dunne, C., Hewson, J., Wohl, C., Wheatley, M., Peterson, A.C., and Reynolds, B.R.** Multipotent CNS stem cells are present in the adult mammalian spinal cord and ventricular neuroaxis. *J. Neurosci.*, 16, 7599-7609, 1996
- Wewer, U.M., Engvall, E., Paulsson, M., Yamada, Y., and Albrechtsen, R.** Laminin A, B1, B2, S and M subunits in the postnatal rat liver development and after partial hepatectomy. *Lab. Invest.*, 66, 378-389, 1992
- Wewer, U.M., Gerecke, D., Durkin, M.E., Kurtz, K.S., Mattei, M.G., Champlaud, M.F., Burgeson, B., and Albrechtsen, R.** Human $\beta 2$ chain of laminin (formerly S chain): cDNA cloning, chromosomal localization, and

expression in carcinomas. *Genomics*, 24, 243-252, 1994

Whitson, J.S., Selkoe, D.J. and Cotman, C.W. Amyloid β protein enhances the survival of hippocampal neurons in vitro. *Science*, 243, 1488-1490, 1989

Willinger, M. and Margolis, D. Effect of weaver (wv) mutation on cerebellar neuron differentiation. I. Qualitative observations of neuron behavior in culture. *Dev. Biol.* 107, 156-172, 1985

Wisniewski, T., and Frangione, B. Apolipoprotein E: a pathological chaperone protein in patients with cerebral and systemic amyloid. *Neurosci. Lett.* 135, 235-238, 1992

Wilson, P.A., and Hemmati-Brivanlou, A. Induction of epidermis and inhibition of neural fate by Bmp-4. *Nature*, 376, 331-333, 1995

Wong, J.T., Wong, S.T., and O'Connor, T.P. Ectopic semaphorin-1a function as an attractive guidance cue for developing peripheral neurons. *Nat. Neurosci.*, 2, 798-803, 1999

Woo, H.J., Shaw, L.M., Messier, J.M., and Mercurio, A.M. The major non-integrin laminin binding protein of macrophages is identical to carbohydrate binding protein 35 (Mac-2). *J. Cell Biol.*, 265, 7097-7099, 1990

Wright, L.L. and Smolen, A.J. Effects of neonatal castration or treatment with dihydrotestosterone on numbers of neurons in the rat superior cervical sympathetic ganglion. *Dev. Brain Res.*, 20, 314-316, 1985

Wullner, U., Löschmann, P.-A., Weller, M., and Klockgether, T. Apotic cell death in cerebellum of mutant weaver and lurcher mice. *Neurosci. Lett.*, 200, 109-112, 1995

Wurst, W., Auerbach, A.B., and Joyner, A.L. Multiple developmental defects in Engrailed-1 mutant mice: an early mid-hindbrain deletion and patterning defects in forelimbs and sternum. *Development*, 120, 2065-2075, 1994

Wyllie, A.H., Kerr, J.F.R., and Currie, A.R. Cell death: the significance of apoptosis. *Int. Rev. Cytol.*, 68, 251-306, 1980

Xu, H., Wu, X.-R. Wewer, U.M., and Engvall, E. Murine muscular dystrophy caused by a mutation in the laminin $\alpha 2$ (Lama2) gene. *Nature Genetics*, 8, 297-302, 1994

Yamamoto, S., Kaneda, Y., Okada, N., Nakagawa, S., Kubo, K., Inoue, S., Maeda, M., Yamashiro, Y., Kawasaki, K., and Mayumi, T. Antimetastatic effects of synthetic peptides containing the core sequence of the type III connecting segment domain (IIICS) of fibronectin. *Anti-Cancer Drugs*, 5, 424-428, 1994

Yamamura, K., Kibbey, M.C., Jun, S.H., and Kleinman, H.K. Effect of matrigel and laminin peptide YIGSR on tumor growth and metastasis. *Semin. Cancer Biol.*, 4, 259-265, 1993a

Yamamura, K., Kibbey, M.C., and Kleinman, H.K. Melanoma cells selected for adhesion to laminin peptides have different malignant properties. *Cancer Res.*, 53, 423-428, 1993b

Yang, Y., Palmer, K.C., Relan, N., Diglio, C., and Schuger, L. Role of laminin polymerization at the epithelial mesenchymal interface in bronchial myogenesis. *Development*, 125, 2621-2629, 1998

Yankner, B.A., Caceres, A., and Duffy, L.K. Nerve growth factor potentiates the neurotoxicity of β amyloid. *Proc. Natl. Acad. Sci. U.S.A* 87, 9020-9023, 1990a

Yankner, B.A., Dawes, L.R., Fisher, S., Villa-Komaroff, L., Oster-Granite, M.L., and Neve, R.L. Neurotoxicity of a fragment of the amyloid precursor associated with Alzheimer's disease. *Science* 245, 417-420, 1989

Yankner, B.A., Duffy, L.K., and Kirschner D.A. Neurotrophic and neurotoxic effects of amyloid β proteins: Reversal by tachykinin neuropeptides. *Science* 250, 279-282, 1990b

Yannas, I.V., Orgill, D.P., Silver, J., Norregaard, T., Zervas, N.T., and Schoene, W.C. Polymeric templates facilitates regeneration of sciatic nerve across 15-mm gap. *Trans. Soc. Biomater.*, 8, 146-146, 1985

Yarnitzky, T., and Volk, T. Laminin is required for head, somatic muscles and gut development in the *Drosophila* embryo. *Dev. Biol.*, 169, 609-618, 1995

Yurchenco, P.D., and Cheng, Y.-S. Self assembly and calcium-binding sites in laminin. A three-arm interaction model. *J. Biol. Chem.*, 268, 17286-17299, 1993

Yurchenco, P.D., Cheng, Y.-S., and Colonato, H. Laminin forms an independent network in basement membranes. *J. Cell Biol.*, 117, 1119-1133, 1992

Zahedi, K. Characterization of the binding of serum amyloid P to laminin. *J. Biol. Chem.*, 272, 2143-2148, 1997

Zak, N.B., Harel, A., Bawnik, Y., Benbasat, S., Vogel, Z., and Schwartz, M. Laminin-immunoreactive sites are induced by growth-associated triggering factors in injured rabbit optic nerve. *Brain Res.*, 408, 263-266, 1987

Zeev-Brann, A.B., Lazarov-Spiegler, O., Brenner, T., and Schwartz, M. Differential effects of central and peripheral nerves on macrophages and microglia. *Glia*, 23, 181-190, 1998

Zhang, L., and Goldman, J.E. Generation of cerebellar interneurons from dividing progenitors in white matter. *Neuron*, 16, 47-54, 1996

Zhang, L., Zhao, B., Yew, D.T., Kusiak, J.W., and Roth, G.S. Processing of Alzheimer's amyloid precursor protein during H₂O₂-induced apoptosis in human neuronal cells. *Biochemical & Biophysical Research Communications.*, 235, 845-848, 1997

Zheng, C., Heintz, N., and Hatten, M.E. CNS gene encoding astrotactin, which supports neuronal migration along glial fibers. *Science*, 272, 417-419, 1996

Zimmerman, L., De Jesús-Escobar, J.M., and Harland, R.M. The Spemann organizer signal noggin binds and inactivates bone morphogenetic protein 4. *Cell*, 599-606, 1996